


CIPROFLOXACIN HYDROCHLORIDE- ciprofloxacin hydrochloride solution

Advagen Pharma Limited

Ciprofloxacin Ophthalmic Solution, USP 0.3% Sterile

DESCRIPTION:

Ciprofloxacin ophthalmic solution is a synthetic, sterile, multiple dose, antimicrobial for topical ophthalmic use. Ciprofloxacin is a fluoroquinolone antibacterial active against a broad spectrum of gram-positive and gram-negative ocular pathogens. It is available as the monohydrochloride monohydrate salt of 1-cyclopropyl-6-fluoro-1,4-dihydro-4-oxo-7-(1-piperazinyl)-3-quinoline-carboxylic acid. It is a faint to light yellow crystalline powder with a molecular weight of 385.8. Its empirical formula is $C_{17}H_{18}FN_3O_3 \cdot HCl \cdot H_2O$ and its chemical structure is as follows:



Ciprofloxacin differs from other quinolones in that it has a fluorine atom at the 6-position, a piperazine moiety at the 7-position, and a cyclopropyl ring at the 1-position.

Each mL of ciprofloxacin ophthalmic solution contains:

Active:Ciprofloxacin hydrochloride 3.5 mg equivalent to 3mg base.

Preservative:Benzalkonium chloride 0.006%.

Inactives:Acetic acid, edetate disodium 0.05%, mannitol 4.6%, Sodium acetate, hydrochloric acid and/or sodium hydroxide (to adjust pH) and water for injection. The pH is between 3.5 to 5.5 and the osmolality is between 290 to 300 mOsm/kg.

CLINICAL PHARMACOLOGY:

Systemic Absorption:A systemic absorption study was performed in which ciprofloxacin ophthalmic solution was administered in each eye every two hours while awake for two days followed by every four hours while awake for an additional 5 days. The maximum reported plasma concentration of ciprofloxacin was less than 5 ng/mL. The mean concentration was usually less than 2.5 ng/mL.

Microbiology: Ciprofloxacin has *in vitro* activity against a wide range of gram-negative and gram-positive organisms. The bactericidal action of ciprofloxacin results from interference with the enzyme DNA gyrase which is needed for the synthesis of bacterial

DNA.

Ciprofloxacin has been shown to be active against most strains of the following organisms both *in vitro* and in clinical infections (SEE **INDICATIONS AND USAGE** section):

Gram-Positive:

Staphylococcus aureus

Staphylococcus epidermidis

Streptococcus pneumoniae

Streptococcus (Viridans Group)

Gram-Negative:

Haemophilus influenzae

Pseudomonas aeruginosa

Serratia marcescens

Ciprofloxacin has been shown to be active *in vitro* against most strains of the following organisms, however, *the clinical significance of these data is unknown*:

Gram-Positive:

Enterococcus faecalis (Many strains are only moderately susceptible)

Staphylococcus haemolyticus

Staphylococcus hominis

Staphylococcus saprophyticus

Streptococcus pyogenes

Gram-Negative:

Acinetobacter calcoaceticus

subsp. anitratus

Aeromonas caviae

rettgeri

Aeromonas hydrophila

Brucella melitensis

enteritidis

Campylobacter coli

Campylobacter jejuni

Citrobacter diversus

Citrobacter freundii

Edwardsiella tarda

parahaemolyticus

Enterobacter aerogenes

Enterobacter cloacae

enterocolitica

Escherichia coli

Haemophilus ducreyi

Haemophilus parainfluenzae

Klebsiella pneumoniae

Klebsiella oxytoca

Legionella pneumophila

Moraxella (Branhamella)

catarrhalis

Morganella morganii

Neisseria gonorrhoeae

Neisseria meningitidis

Pasteurella multocida

Proteus mirabilis

Proteus vulgaris

Providencia

Providencia stuartii

Salmonella

Salmonella typhi

Shigella sonnei

Shigella flexneri

Vibrio cholerae

Vibrio

Vibrio vulnificus

Yersinia

Other Organisms: *Chlamydia trachomatis* (only moderately susceptible) and *Mycobacterium tuberculosis* (only moderately susceptible).

Most strains of *Pseudomonas cepacia* and some strains of *Pseudomonas maltophilia* are resistant to ciprofloxacin as are most anaerobic bacteria, including *Bacteroides*

fragilis and *Clostridium difficile*. The minimal bactericidal concentration (MBC) generally does not exceed the minimal inhibitory concentration (MIC) by more than a factor of 2. Resistance to ciprofloxacin *in vitro* usually develops slowly (multiple-step mutation).

Ciprofloxacin does not cross-react with other antimicrobial agents such as beta-lactams or aminoglycosides; therefore, organisms resistant to these drugs may be susceptible to ciprofloxacin.

Clinical Studies:

Following therapy with ciprofloxacin ophthalmic solution, 76% of the patients with corneal ulcers and positive bacterial cultures were clinically cured and complete re-epithelialization occurred in about 92% of the ulcers.

In 3 and 7 day multicenter clinical trials, 52% of the patients with conjunctivitis and positive conjunctival cultures were clinically cured and 70-80% had all causative pathogens eradicated by the end of treatment.

In a randomized, double-masked, multicenter, parallel-group clinical trial of pediatric patients with bacterial conjunctivitis, between birth and 31 days of age, patients were dosed with ciprofloxacin or another anti-infective agent. Clinical outcomes for the trial demonstrated a clinical cure rate of 80% at Day 9 and a microbiological eradication success rate of 85% at Day 9.

Please note that microbiologic eradication does not always correlate with clinical outcome in anti-infective trials.

INDICATIONS AND USAGE:

Ciprofloxacin ophthalmic Solution is indicated for the treatment of infections caused by susceptible strains of the designated microorganisms in the conditions listed below:

Corneal Ulcers: *Pseudomonas aeruginosa*
*Serratia marcescens**
Staphylococcus aureus
Staphylococcus epidermidis
Streptococcus pneumoniae
Streptococcus (Viridans Group) *

Conjunctivitis: *Haemophilus influenzae*
Staphylococcus aureus
Staphylococcus epidermidis
Streptococcus pneumoniae

*Efficacy for this organism was studied in fewer than 10 infections.

CONTRAINDICATIONS:

A history of hypersensitivity to ciprofloxacin or any other component of the medication

is a contraindication to its use. A history of hypersensitivity to other quinolones may also contraindicate the use of ciprofloxacin.

WARNINGS:

NOT FOR INJECTION INTO THE EYE.

Serious and occasionally fatal hypersensitivity (anaphylactic) reactions, some following the first dose, have been reported in patients receiving systemic quinolone therapy. Some reactions were accompanied by cardiovascular collapse, loss of consciousness, tingling, pharyngeal or facial edema, dyspnea, urticaria, and itching. Only a few patients had a history of hypersensitivity reactions. Serious anaphylactic reactions require immediate emergency treatment with epinephrine and other resuscitation measures, including oxygen, intravenous fluids, intravenous antihistamines, corticosteroids, pressor amines and airway management, as clinically indicated.

Remove contact lenses before using.

PRECAUTIONS:

General:As with other antibacterial preparations, prolonged use of ciprofloxacin may result in overgrowth of nonsusceptible organisms, including fungi. If superinfection occurs, appropriate therapy should be initiated. Whenever clinical judgment dictates, the patient should be examined with the aid of magnification, such as slit lamp biomicroscopy and, where appropriate, fluorescein staining.

Ciprofloxacin should be discontinued at the first appearance of a skin rash or any other sign of hypersensitivity reaction.

In clinical studies of patients with bacterial corneal ulcer, a white crystalline precipitate located in the superficial portion of the corneal defect was observed in 35 (16.6%) of 210 patients. The onset of the precipitate was within 24 hours to 7 days after starting therapy. In one patient, the precipitate was immediately irrigated out upon its appearance. In 17 patients, resolution of the precipitate was seen in 1 to 8 days (seven within the first 24 to 72 hours), in five patients, resolution was noted in 10 to 13 days. In nine patients, exact resolution days were unavailable; however, at follow-up examinations, 18 to 44 days after onset of the event, complete resolution of the precipitate was noted. In three patients, outcome information was unavailable. The precipitate did not preclude continued use of ciprofloxacin, nor did it adversely affect the clinical course of the ulcer or visual outcome. (SEE **ADVERSE REACTIONS**).

Information for patients:Do not touch dropper tip to any surface, as this may contaminate the solution.

Drug Interactions:Specific drug interaction studies have not been conducted with ophthalmic ciprofloxacin. However, the systemic administration of some quinolones has been shown to elevate plasma concentrations of theophylline, interfere with the metabolism of caffeine, enhance the effects of the oral anticoagulant, warfarin, and its derivatives, and has been associated with transient elevations in serum creatinine in patients receiving cyclosporine concomitantly.

Carcinogenesis, Mutagenesis, Impairment of Fertility:Eight *in vitro* mutagenicity tests have been conducted with ciprofloxacin and the test results are listed below:

Salmonella/Microsome Test (Negative)

E. coli DNA Repair Assay (Negative)

Mouse Lymphoma Cell Forward Mutation Assay (Positive)

Chinese Hamster V79 Cell HGPRT Test (Negative)

Syrian Hamster Embryo Cell Transformation Assay (Negative)

Saccharomyces cerevisiae Point Mutation Assay (Negative)

Saccharomyces cerevisiae Mitotic Crossover and Gene Conversion Assay (Negative)

Rat Hepatocyte DNA Repair Assay (Positive)

Thus, two of the eight tests were positive, but the results of the following three *in vivo* test systems gave negative results:

Rat Hepatocyte DNA Repair Assay

Micronucleus Test (Mice)

Dominant Lethal Test (Mice)

Long term carcinogenicity studies in mice and rats have been completed. After daily oral dosing for up to two years, there is no evidence that ciprofloxacin had any carcinogenic or tumorigenic effects in these species.

Pregnancy: Reproduction studies have been performed in rats and mice at doses up to six times the usual daily human oral dose and have revealed no evidence of impaired fertility or harm to the fetus due to ciprofloxacin. In rabbits, as with most antimicrobial agents, ciprofloxacin (30 and 100 mg/kg orally) produced gastrointestinal disturbances resulting in maternal weight loss and an increased incidence of abortion. No teratogenicity was observed at either dose. After intravenous administration, at doses up to 20 mg/kg, no maternal toxicity was produced and no embryotoxicity or teratogenicity was observed. There are no adequate and well controlled studies in pregnant women.

Ciprofloxacin Ophthalmic Solution should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers: It is not known whether topically applied ciprofloxacin is excreted in human milk; however, it is known that orally administered ciprofloxacin is excreted in the milk of lactating rats and oral ciprofloxacin has been reported in human breast milk after a single 500 mg dose. Caution should be exercised when Ciprofloxacin Ophthalmic Solution is administered to a nursing mother.

Pediatric Use:

The safety and effectiveness of ciprofloxacin ophthalmic solution 0.3% have been established in all ages. Use of ciprofloxacin is supported by evidence from adequate and well controlled studies of ciprofloxacin in adults, children and neonates [see *Clinical Studies*]. Although ciprofloxacin and other quinolones cause arthropathy in immature animals after oral administration, topical ocular administration of ciprofloxacin to immature animals did not cause any arthropathy and there is no evidence that the ophthalmic dosage form has any effect on the weight bearing joints.

Geriatric Use: No overall differences in safety or effectiveness have been observed between elderly and younger patients.

ADVERSE REACTIONS:

The most frequently reported drug related adverse reaction was local burning or discomfort. In corneal ulcer studies with frequent administration of the drug, white crystalline precipitates were seen in approximately 17% of patients (SEE **PRECAUTIONS**). Other reactions occurring in less than 10% of patients included lid margin crusting, crystals/scales, foreign body sensation, itching, conjunctival hyperemia and a bad taste following instillation. Additional events occurring in less than 1% of patients included corneal staining, keratopathy/keratitis, allergic reactions, lid edema, tearing, photophobia, corneal infiltrates, nausea and decreased vision.

To report SUSPECTED ADVERSE REACTIONS, contact Advagen Pharma Ltd, at 866-488-0312 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

OVERDOSAGE:

A topical overdose of ciprofloxacin ophthalmic solution may be flushed from the eye(s) with warm tap water.

DOSAGE AND ADMINISTRATION:

Corneal Ulcers:The recommended dosage regimen for the treatment of **corneal ulcers** is two drops into the affected eye every 15 minutes for the first six hours and then two drops into the affected eye every 30 minutes for the remainder of the first day. On the second day, instill two drops in the affected eye hourly. On the third through the fourteenth day, place two drops in the affected eye every four hours. Treatment may be continued after 14 days if corneal re-epithelialization has not occurred.

Bacterial Conjunctivitis:The recommended dosage regimen for the treatment of **bacterial conjunctivitis** is one or two drops instilled into the conjunctival sac(s) every two hours while awake for two days and one or two drops every four hours while awake for the next five days.

How Supplied:As a sterile ophthalmic solution: 2.5 mL, 5 mL and 10 mL in translucent LDPE bottle, translucent LDPE nozzle and tan coloured polypropylene closure.

2.5 mL - NDC 72888-102-21

5 mL - NDC 72888-102-22

10 mL - NDC 72888-102-23

STORAGE:Store at 2°C to 25°C (36°F to 77°F). Protect from light.

ANIMAL PHARMACOLOGY:

Ciprofloxacin and related drugs have been shown to cause arthropathy in immature animals of most species tested following oral administration. However, a one-month topical ocular study using immature Beagle dogs did not demonstrate any articular lesions.

Distributed by:


Advagen Pharma Ltd.,

East Windsor, NJ 08520, USA.

Rev. 01, 09/2024

PACKAGE LABEL-PRINCIPAL DISPLAY PANEL

Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-21 - 2.5 mL Carton



Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-21 - 2.5 mL Container

N 3 72888 10221 4

Each mL contains:
Active: Ciprofloxacin hydrochloride 3.5 mg equivalent to 3 mg base.
Preservative: Benzalkonium Chloride 0.006%.
Store at 2° to 25°C (36° to 77°F). Protect from light.
Usual Dosage: See accompanying package insert for dosage information.
FOR OPHTHALMIC USE ONLY
KEEP CONTAINER TIGHTLY CLOSED

NDC 72888 -102-21

Ciprofloxacin Ophthalmic Solution, USP

0.3%

Sterile Ophthalmic Solution
Rx Only 2.5 mL

ADVAGEN

Distributed by:
Advagen Pharma Ltd.,
East Windsor, NJ, 08520, USA
Code No.: MH/DRUGS/28-KD/74

Rev. 01, 09/24 PLB0252

Overprinting Area
18.6 mm X 8 mm

Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-22 - 5 mL Carton



Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-22 - 5 mL Container



Each mL contains:
Active: Ciprofloxacin hydrochloride 3.5 mg equivalent to 3 mg base.
Preservative: Benzalkonium Chloride 0.006%.
Store at 2° to 25°C (36° to 77°F). Protect from light.
Usual Dosage: See accompanying package insert for dosage information.
FOR OPHTHALMIC USE ONLY
KEEP CONTAINER TIGHTLY CLOSED

NDC 72888 -102-22

Ciprofloxacin Ophthalmic Solution, USP

0.3%

Sterile Ophthalmic Solution
 Rx Only 5 mL



Distributed by:
Advagen Pharma Ltd.,
 East Windsor, NJ, 08520, USA
 Code No.: MH/DRUGS/28-KD/74

Rev. 02, 10/24 PLB0253



Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-23 - 10 mL Carton



Ciprofloxacin Ophthalmic Solution, 0.3% - NDC 72888-102-23 - 10 mL Container



Each mL contains:
Active: Ciprofloxacin hydrochloride 3.5 mg
 equivalent to 3 mg base. **Preservative:**
 Benzalkonium Chloride 0.006%.

Store at 2° to 25°C (36° to 77°F).
 Protect from light.

Usual Dosage: See accompanying
 package insert for dosage information.

FOR OPHTHALMIC USE ONLY
KEEP CONTAINER TIGHTLY CLOSED

NDC 72888 -102-23

Ciprofloxacin Ophthalmic Solution, USP

0.3%

Sterile Ophthalmic Solution

Rx Only

10 mL



Distributed by: **Advagen Pharma Ltd.**,
 East Windsor, NJ 08520, USA

Code No.: MH/DRUGS/28-KD/74

Rev 01, 09/24

PLB0254



Overprinting Area
 18.6 X 8 mm



CIPROFLOXACIN HYDROCHLORIDE

ciprofloxacin hydrochloride solution

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:72888-102
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CIPROFLOXACIN HYDROCHLORIDE (UNII: 4BA73M5E37) (CIPROFLOXACIN - UNII:5E8K9I004U)	CIPROFLOXACIN	3 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ACETIC ACID (UNII: Q40Q9N063P)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
MANNITOL (UNII: 3OWL53L36A)	
SODIUM ACETATE (UNII: 4550K0SC9B)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72888-102-21	2.5 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/2022	
2	NDC:72888-102-22	5 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/2022	
3	NDC:72888-102-23	10 mL in 1 BOTTLE; Type 0: Not a Combination Product	03/16/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075928	03/16/2022	

Labeler - Advagen Pharma Limited (051627256)

Revised: 3/2025

Advagen Pharma Limited