

**STOMACH RELIEF- bismuth subsalicylate suspension**  
**Consumer Value Products, Inc.**

-----  
**STOMACH RELIEF**

***Drug Facts***

***Active ingredient***  
***(in each 30 mL dose cup or 2 tablespoons)***

Bismuth subsalicylate 525 mg

***Purposes***

Bismuth subsalicylate ..... Upset stomach reliever  
and antidiarrheal

***Uses*** relieves ■ travelers' diarrhea ■ diarrhea ■ upset  
stomach due to overindulgence of food and drink including:  
■ heart burn ■ indigestion ■ nausea ■ gas ■ belching

***Warnings***

**Reye's syndrome:** Children and teenagers who have or are  
recovering from chicken pox or flu-like symptoms should not use  
this product. When using this product, if changes in behavior with  
nausea and vomiting occur, consult a doctor because these  
symptoms could be an early sign of Reye's syndrome, a rare but  
serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use** if you have ■ an ulcer ■ a bleeding problem  
■ black or bloody stool

**Ask a doctor before use if you have**

- fever ■ mucus in stool

**Ask a doctor or pharmacist before use if you are**

taking any drug for ■ anticoagulation (thinning of the  
blood) ■ diabetes ■ gout ■ arthritis

**When using this product** a temporary, but harmless,  
darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

**If pregnant or breast feeding,** ask health professional before use.

**Keep out of reach of children.** In case of overdose, get  
medical help or contact a Poison Control Center right away.

***Directions***

- Shake well before use
- use dose cup or tablespoon (TBSP)
- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1/2 to 1 hour as needed
- do not exceed 8 doses (240 mL or 16 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information ■ each 30 mL dose**

**contains:** sodium 13 mg, salicylate 256 mg

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- Low sodium

**Inactive ingredients** benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xantham gum

**Questions?** 1-877-798-5944

**Principal Display Panel**

**CVP®**

**HEALTH**

**\*COMPARE TO  
PEPTO-BISMOL®  
ACTIVE INGREDIENT**

**Stomach  
Relief**

**PROTECTIVE COATING ACTION  
Bismuth Subsalicylate**

**SOOTHING RELIEF FOR:**

**UPSET STOMACH  
HEARTBURN  
INDIGESTION  
NAUSEA  
COMMON DIARRHEA**

**8 FL OZ (237 mL)**

**TAMPER EVIDENT: Do not use if imprinted  
shrinkband is missing or broken**

\*This product is not manufactured or distributed by Procter & Gamble,

Inc., the distributor of Pepto-Bismol™.

**DISTRIBUTED BY  
CONSUMER VALUE PRODUCTS, INC.  
P.O. BOX 6115 TEMPLE, TX 76502**

**Made in USA  
LR-102 REV01**

CVPproducts.com

7 61706 16202 6

**PEEL**

**HERE**

**TAMPER EVIDENT: Do not use if imprinted shrinkband is missing or broken**

**CVP HEALTH**

**COMPARE TO PEPTO-BISMOL® ACTIVE INGREDIENT**

**STOMACH RELIEF**

PROTECTIVE COATING ACTION  
BISMUTH SUBSALICYLATE

SOOTHING RELIEF FOR:  
UPSET STOMACH  
HEARTBURN  
INDIGESTION  
NAUSEA  
COMMON DIARRHEA

8 FL OZ (237mL)

**Drug Facts (continued)**

**When using this product** a temporary, but harmless, darkening of the stool and/or tongue may occur

**Stop use and ask a doctor if**

- diarrhea lasts more than 2 days
- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs

**If pregnant or breast feeding**, ask health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol™.

**DISTRIBUTED BY  
CONSUMER VALUE PRODUCTS, INC.  
P.O. BOX 6115 TEMPLE, TX 76502  
CVPproducts.com**

**Made in USA  
LR-102 REV01**

7 61706 16202 6

**PEEL HERE**

<b>Drug Facts</b>	<b>Purposes</b>
<b>Active ingredient</b> (in each 30 mL dose cup or 2 tablespoons) Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal
<b>Uses</b> relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence of food and drink including: ■ heart burn ■ indigestion ■ nausea ■ gas ■ belching	
<b>Warnings</b> <b>Reye's syndrome:</b> Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.	
<b>Allergy alert:</b> Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products	
<b>Do not use</b> if you have ■ an ulcer ■ a bleeding problem ■ black or bloody stool	
<b>Ask a doctor before use</b> if you have ■ fever ■ mucus in stool	
<b>Ask a doctor or pharmacist before use</b> if you are taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis	

**Drug Facts (continued)**

**Directions**

- Shake well before use
- use dose cup or tablespoon (TBSP)
- adults and children 12 years and over: 1 dose (30 mL or 2 TBSP) every 1/2 to 1 hour as needed
- do not exceed 8 doses (240 mL or 16 TBSP) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL dose contains: sodium 13 mg, salicylate 256 mg

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- low sodium

**Inactive ingredients** benzoic acid, D&C red #22, D&C red # 28, flavor, purified water, saccharin sodium, salicylic acid, sodium salicylate, xanthan gum

**Questions?** 1-877-798-5944

Stomach Relief by Consumer Value Products, Inc.

res

# STOMACH RELIEF

bismuth subsalicylate suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:57243-557
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>SODIUM SALICYLATE</b> (UNII: WQ1H85SYP)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	pink	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MINT	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:57243-557-26	237 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M008	10/01/2018	

**Labeler** - Consumer Value Products, Inc. (009830555)

**Registrant** - AptaPharma Inc. (790523323)

**Establishment**

Name	Address	ID/FEI	Business Operations
AptaPharma Inc.		790523323	manufacture(57243-557)

Revised: 12/2023

Consumer Value Products, Inc.