NICADAN- ascorbic acid, niacinamide, pyridoxine hydrochloride, folic acid, magnesium citrate, zinc gluconate, copper gluconate, and .alpha.-lipoic acid tablet, film coated Medimetriks Pharmaceuticals, Inc.

NICADANTM

Rx Only

Niacinamide, Vitamin B6, Vitamin C, Folic Acid, Copper, Magnesium, Zinc & Alpha Lipoic Acid

DESCRIPTION

Nicadan[™] tablets are a specially formulated dietary supplement containing natural ingredients with antiinflammatory properties. Each pink-coated tablet is oval shaped, scored and embossed with "MM". **Nicadan**[™] is for oral administration only.

Nicadan[™] should be administered under the supervision of a licensed medical practitioner.

INGREDIENTS

Each tablet of **Nicadan**TM contains:

Vitamin C (as Ascorbic Acid)	100 mg
Niacinamide (Vitamin B-3)	800 mg
Vitamin B-6 (as Pyridoxine HCI)	10 mg
Folic Acid	500 mcg
Magnesium (as Magnesium Citrate)	5 mg
Zinc (as Zinc Gluconate)	20 mg
Copper (as Copper Gluconate)	2 mg
Alpha Lipoic Acid	50 mg

Other Ingredients: Microcrystalline cellulose, Povidone, Hypromellose, Croscarmellose Sodium, Polydextrose, Talc, Magnesium Sterate Vegetable, Vegetable Stearine, Red Beet Powder, Titanium Dioxide, Maltodextrin and Triglycerides.

Nicadan[™] is free of yeast, wheat, dairy, egg, soy, gluten, artificial color or flavors, added sugar, starch and preservatives.

USES

As a dietary supplement, **Nicadan**TM tablets are beneficial in maintaining overall skin health, especially in patients with inflammatory skin conditions.¹

¹ This statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

WARNINGS

If you are pregnant, nursing or taking medication, consult your doctor before use.

CONTRAINDICATIONS

Use of **Nicadan™** is contraindicated in patients who are hypersensitive to any ingredient in this tablet.

PRECAUTIONS

There have been reports of GI upset, nausea, vomiting, heartburn, diarrhea, dizziness, drowsiness or headache with large doses of niacinamide. Rarely, niacinamide in doses of 3 grams/day or more has been associated with elevated liver function tests. There are case reports in the literature indicating that niacinamide may reduce the hepatic metabolism of primidone and carbamazepine. Individuals taking these medications should consult their physician. Individuals taking antidiabetes medications should have their blood glucose levels monitored.

Allergic sensitization has been reported rarely following oral administration of folic acid. Folic acid above 1 mg daily may obscure pernicious anemia in that hematologic remission may occur while neurological manifestations remain progressive.

DOSAGE AND ADMINISTRATION

Take one tablet daily with food or as directled by a physician. **Nicadan™** tablets are scored, so they may be broken in half if required.

HOW SUPPLIED

Nicadan[™] is available in a bottle containing 60 tablets. 43538-440-60

Store at 15°C to 30°C (59°F to 86°F). Keep bottle tightly closed. Store in cool dry place.

KEEP THIS AND ALL MEDICATIONS OUT OF THE REACH OF CHILDREN.

You may report side effects by calling 1-973-882-7512.

Manufactured for:

MEDIMETRIKS PHARMACEUTICALS, INC.

383 Route 46 West, Fairfield, NJ 07004-2402 USA

www.medimetriks.com

Iss: 2/2014

IP038

PRINCIPAL DISPLAY PANEL - 60 Tablet Bottle Carton

43538-440-60

Rx Only NICADAN™

Niacinamide, Vitamin B6, Vitamin C, Folic Acid, Copper, Magnesium, Zinc & Alpha Lipoic Acid

Helps Maintain Overall Skin Health[†]


Dermatologist Developed

60 Tablets

Dietary Supplement

MEDIMETRIKS

PHARMACEUTICALS, INC.



SUGGESTED USE: Take one tablet daily with food or as directed by a physician. Keep bottle tightly closed. Store in cool dry place. WARNING: If you are pregnant, nursing or taking medication, consult your doctor before use.

Keep out of reach of children. DO NOT USE IF SEAL UNDER CAP IS BROKEN.

Supplement Facts

Serving Size: One (1) Tablet

	Amount Per Tablet	% Daily Value
Vitamin C (as Ascorbic Acid)	100 mg	167%
Niacinamide (Vitamin B-3)	800 mg	4000%
Vitamin B-6 (as Pyridoxine HCI)	10 mg	500%
Folic Acid	500 mcg	125%
Magnesium (as Magnesium Citra	tte) 5 mg	1%
Zinc (as Zinc Gluconate)	20 mg	133%
Copper (as Copper Gluconate)	2 mg	100%
Alpha Lipoic Acid	50 mg	*
* Daily Value not established		- 8

Other Ingredients: Microcrystalline Cellulose, Povidone, Hypromellose, Croscarmellose Sodium, Polydextrose, Talc, Magnesium Sterate Vegetable, Vegetable Stearine, Red Beet Powder, Titanium Dioxide, Maltodextrin and Triglyerides.

Nicadan™ is free of yeast, wheat, dairy, egg, soy, gluten, artificial color or flavors, added sugar, starch and preservatives.

tThis statement has not been evaluated by the Food and Drug Administration. This product is not intended to diagnose, treat, cure or prevent any disease.

Store between 15°-30°C (59°-86°F).

Manufactured for: MEDIMETRIKS PHARMACEVITICALS. INC. 383 Route 46 West, Fairfield, NJ 07004-2402 USA www.medimetriks.com

IC142

Rx Only NICADAN[™] Niacinamide, Vitamin B6, Vitamin C, Folic Acid, Copper, Magnesium, Zinc & Alpha Lipoic Acid


Gluten-Free Sodium-Free Sugar-Free Vegan No Artifical Colors No Artifical Flavors

60 Tablets

Dietary Supplement

MEDIMETRIKS PHARMACEUTICALS, INC.

BAR CODE 3 43538 44060 (#)



NICADAN

т

ascorbic acid, niacinamide, pyridoxine hydrochloride, folic acid, magnesium citrate, zinc gluconate, copper gluconate, and .alpha.-lipoic acid tablet, film coated

Product Information						
Product Type	DIETARY SUPPLEMENT Item Code (Source) NHRI			NHRIC:4353	RIC:43538-440	
Route of Administration	ORAL					
Active Ingredient/Active Mo	biety					
Ir	igredient Name		Basis of S	trength	Strength	
Ascorbic Acid (UNII: PQ6CK8PD0R) (Ascorbic Acid - UNII:PQ6CK8PD0R)			Ascorbic Acid		100 mg	
Niacinamide (UNII: 25X5118RD4) (Niacinamide - UNII:25X5118RD4)			Niacinamide		800 mg	
pyridoxine hydrochloride (UNII: 68Y4CF58BV) (pyridoxine - UNII:KV2JZ1BI6Z)			oyridoxine hyd	ro chlo ride	10 mg	
folic acid (UNII: 935E97BOY8) (folic acid - UNII:935E97BOY8)folic acid			500 ug			
magnesium citrate (UNII: RHO26O1T9V) (magnesium cation - UNII:T6V3LHY838) magnesium c			nagnesium citi	rate	5 mg	
zinc gluconate (UNII: U6WSN5SQ1Z) (zinc cation - UNII:13S1S8SF37)			zinc gluconate		20 mg	
copper gluconate (UNII: RV823G6G67) (cupric cation - UNII:8CBV67279L) cu			cupric cation		20 mg	
copper graconate (orth: htto25606		K73Y) .	alphalipoic a	_	50 mg	

Inactive Ingredients				
Ingredient Name	Strength			
cellulose, microcrystalline (UNII: OP1R32D61U)				
povidones (UNII: FZ989GH94E)				
hypromelloses (UNII: 3NXW29V3WO)				
croscarmellose sodium (UNII: M28OL1HH48)				
polydextrose (UNII: VH2XOU12IE)				
talc (UNII: 7SEV7J4R1U)				
titanium dioxide (UNII: 15FIX9V2JP)				
maltodextrin (UNII: 7CVR7L4A2D)				

# It	em Code	Package Description	Marketing Start Date	Marketing End Date
1 NHRIC:43	538-440-60	1 in 1 CARTON		
1		60 in 1 BOTTLE		

mation		
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	02/22/2014	
ľ		Application Number or Monograph Citation Marketing Start Date

Supplement F	acts	
Serving Size :		Serving per Container :
	Amount Per Serving	% Daily Value
color		
scoring	2	
shape		
size (solid drugs)	22 mm	
imprint		

Labeler - Medimetriks Pharmaceuticals, Inc. (019903816)

Establishment			
Name	Address	ID/FEI	Business Operations
Contract Pharmacal		968335112	MANUFACTURE(43538-440)

Revised: 2/2014

Medimetriks Pharmaceuticals, Inc.