

ERYTHROMYCIN- erythromycin gel
Aidarex Pharmaceuticals LLC

Erythromycin Topical Gel USP, 2%


For Dermatologic Use Only – Not for Ophthalmic Use

Rx Only

DESCRIPTION

Erythromycin Topical Gel USP, 2% contains erythromycin (3R*, 4S*, 5S*, 6R*, 7R*, 9R*, 11R*, 12R*, 13S*, 14R*)-4-[(2,6-Dideoxy-3-C-methyl-3-O-methyl- α -L-ribo-hexopyranosyl)oxy]-14-ethyl-7, 12, 13-trihydroxy-3, 5, 7, 9, 11, 13-hexamethyl-6-[[3, 4, 6,-trideoxy-3-(dimethylamino)- β -D-xylo-hexopyranosyl]oxy]oxacyclotetradecane-2,10-dione), for topical dermatological use. Erythromycin is a macrolide antibiotic produced from a strain of *Saccaropolyspora erythraea* (formerly *Streptomyces erythreus*). It is a base and readily forms salts with acids.

Chemically, erythromycin is C₃₇H₆₇NO₁₃. It has the following structural formula:



Erythromycin has a molecular weight of 733.94. It is a white or slightly yellow, odorless or practically odorless, bitter crystalline powder. Erythromycin is very soluble in very polar organic solvents such as alcohols, acetone, chloroform, acetonitrile and ethyl acetate. It is moderately soluble in less polar solvents such as ether, dichloroethylene and amyl acetate. It is slightly soluble in nonpolar solvents such as hexane. It is very poorly soluble in water.

Each gram of Erythromycin Topical Gel USP, 2% contains 20 mg of erythromycin USP in a vehicle consisting of dehydrated alcohol and hydroxypropyl cellulose.

CLINICAL PHARMACOLOGY

The exact mechanism by which erythromycin reduces lesions of acne vulgaris is not fully known; however, the effect appears to be due in part to the antibacterial activity of the drug.

MICROBIOLOGY

Erythromycin acts by inhibition of protein synthesis in susceptible organisms by reversibly binding to 50S ribosomal subunits, thereby inhibiting translocation of aminoacyl transfer-RNA and inhibiting polypeptide synthesis. Antagonism has been demonstrated *in vitro* between erythromycin, lincomycin,

chloramphenicol, and clindamycin.

INDICATIONS AND USAGE

Erythromycin Topical Gel USP, 2% is indicated for the topical treatment of acne vulgaris.

CONTRAINDICATIONS

Erythromycin Topical Gel USP, 2% is contraindicated in those individuals who have shown hypersensitivity to any of its components.

WARNINGS

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including erythromycin, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of “antibiotic-associated colitis”.

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

PRECAUTIONS

General -

For topical use only; not for ophthalmic use. Concomitant topical acne therapy should be used with caution because a possible cumulative irritancy effect may occur, especially with the use of peeling, desquamating or abrasive agents.

Avoid contact with eyes and all mucous membranes.

The use of antibiotic agents may be associated with the overgrowth of antibiotic-resistant organisms. If this occurs, discontinue use and take appropriate measures.

Information For Patients -

Patients using Erythromycin Topical Gel USP, 2% should receive the following information and instructions:

1. This medication is to be used as directed by the physician. It is for external use only. Avoid contact with the eyes, nose, mouth, and all mucous membranes.
2. This medication should not be used for any disorder other than that for which it was prescribed.
3. Patients should not use any other topical acne medication unless otherwise directed by their physician.
4. Patients should report to their physician any signs of local adverse reactions.

Carcinogenesis, Mutagenesis, Impairment of Fertility -

No animal studies have been performed to evaluate carcinogenic and mutagenic potential, or effects on

fertility of topical erythromycin. However, long-term (2-year) oral studies in rats with erythromycin ethylsuccinate and erythromycin base did not provide evidence of tumorigenicity. There was no apparent effect on male or female fertility in rats fed erythromycin (base) at levels up to 0.25% of diet.

Pregnancy:

Teratogenic Effects:

Pregnancy Category B -

There was no evidence of teratogenicity or any other adverse effect on reproduction in female rats fed erythromycin base (up to 0.25% of diet) prior to and during mating, during gestation and through weaning of two successive litters. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used in pregnancy only if clearly needed. Erythromycin has been reported to cross the placental barrier in humans, but fetal plasma levels are generally low.

Nursing Women -

It is not known whether erythromycin is excreted in human milk after topical application. However, erythromycin is excreted in human milk following oral and parenteral erythromycin administration. Therefore, caution should be exercised when erythromycin is administered to a nursing woman.

Pediatric Use -

Safety and effectiveness in pediatric patients have not been established.

KEEP OUT OF REACH OF CHILDREN.

ADVERSE REACTIONS

In controlled clinical trials, the incidence of burning associated with erythromycin topical gel USP, 2% was approximately 25%. The following additional local adverse reactions have been reported occasionally: peeling, dryness, itching, erythema, and oiliness. Irritation of the eyes and tenderness of the skin have also been reported with the topical use of erythromycin. A generalized urticarial reaction, possibly related to the use of erythromycin, which required systemic steroid therapy has been reported.

DOSAGE AND ADMINISTRATION

Erythromycin Topical Gel USP, 2% should be applied sparingly as a thin film to affected area(s) once or twice a day after the skin is thoroughly cleansed and patted dry. If there has been no improvement after 6 to 8 weeks, or if the condition becomes worse, treatment should be discontinued, and the physician should be reconsulted. Spread the medication lightly rather than rubbing it in. There are no data directly comparing the safety and efficacy of b.i.d. versus q.d. dosing.

HOW SUPPLIED

Erythromycin Topical Gel USP, 2% is available as follows:

60 g sealed metal tube (NDC 53217-**0021**-01)

STORAGE

Note - FLAMMABLE. Keep away from heat and flame. Store and dispense in original container. Keep tube tightly closed.

Store at 20-25°C (68-77°F) [see USP Controlled Room Temperature].

Made in Israel

Manufactured by Perrigo

Yeruham 80500, Israel

Distributed By



Allegan, MI 49010 • www.perrigo.com

Repackaged By :

Aidarex Pharmaceuticals LLC,

Corona, CA 92880

Rev. 09/11

6J400 RC J4

Principal Display Panel - 60 g Tube

Rx Only

Erythromycin Topical Gel USP, 2%

For Dermatologic Use Only - Not for Ophthalmic Use

CAUTION: FEDERAL LAW PROHIBITS DISPENSING WITHOUT PRESCRIPTION. SEE PACKAGE INSERT. KEEP OUT OF REACH OF CHILDREN. STORE AT CONTROLLED ROOM TEMP 15-30C (59-86F)

Packaged and Distributed by: **AIDAREX PHARMACEUTICALS LLC.**

ERYTHROMYCIN GEL

2%

60 G

EACH GRAM CONTAINS THE FOLLOWING ACTIVE INGREDIENTS:
ERYTHROMYCIN20MG

RX 1000966176

GENERIC FOR : ERYTHROCIN APPLY _____ TIMES DAILY
NDC: 53217-0021-01 APLIQUE _____ VECES AL DIA

MFG: PERRIGO, ISRAEL

ERYTHROMYCIN GEL 2%
NDC: 53217-0021-01
RX 1000966176

ERYTHROMYCIN GEL 2%
NDC: 53217-0021-01
RX 1000966176

ERYTHROMYCIN GEL 2% 60 G
NDC: 53217-0021-01
RX 1000966176

ERYTHROMYCIN GEL 2% 60 G
NDC: 53217-0021-01
RX 1000966176

PATIENT CHART BILL

PEEL HERE

PEEL HERE

ERYTHROMYCIN

erythromycin gel

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:53217-021(NDC:45802-966)
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ERYTHROMYCIN (UNII: 63937KV33D) (ERYTHROMYCIN - UNII:63937KV33D)	ERYTHROMYCIN	20 mg in 1 g

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
HYDROXYPROPYL CELLULOSE (TYPE H) (UNII: RFW2ET671P)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53217-021-01	60 g in 1 TUBE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA063211	06/12/2008	

Labeler - Aidarex Pharmaceuticals LLC (801503249)

Revised: 9/2013

Aidarex Pharmaceuticals LLC