LAXATIVE- bisacodyl tablet, delayed release WinCo Foods, LLC

Winco 44-676A

Active ingredient (in each tablet)

Bisacodyl USP, 5 mg

Purpose

Stimulant laxative

Uses

- for relief of occasional constipation (irregularity)
- this product generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

if you cannot swallow without chewing.

Ask a doctor before use if you have

- a sudden change in bowel habits that lasts more than 2 weeks
- stomach pain, nausea or vomiting

When using this product

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness, and cramps

Stop use and ask a doctor if

- you have rectal bleeding or fail to have a bowel movement after use of a laxative.
 These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

do not take more than directed

take with a glass of water

adults and children 12 years	take 1 to 3 tablets in a single
	daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
crilidien o to under 12 years	dose
	ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

Inactive ingredients

carmine, colloidal silicon dioxide, corn starch, FD&C red #40 aluminum lake, hypromellose, lactose anhydrous, magnesium stearate, methacrylic acid, microcrystalline cellulose, polydextrose, polyethylene glycol, shellac wax, simethicone, sodium bicarbonate, sodium lauryl sulfate, stearic acid, talc, titanium dioxide, triacetin, triethyl citrate

Questions?

Call 1-800- 824-1706 Monday - Friday 9am - 4pm MST

Principal Display Panel

Winco® Foods

Laxative

BISACODYL USP 5 mg/STIMULANT LAXATIVE

30

Enteric

Coated

Tablets

Gentle, Overnight Relief

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

BISACODYL USP 5 mg/STIMULANT LAXATIVE

100% SATISFACTION GUARANTEE

Comments & Questions? www.wincofoods.com

70552¹¹24133

magnesium stearate, methacrylic acid, micro crystalline #40 aluminum lake, hypromellose, lactose anhydrous,

carmine, colloidal silicon dioxide, com starch, FD&C red

■ see end flap for expiration date and lot number

OPENED OR BLISTER IS TORN OR BROKEN

■ store at 25°C (77°F); excursions permit ted between

■ TAM PER EVIDENT: DO NOT USE IF OUTER PACKAGE IS

sak a doctor

take 1 tablet in a single

single daily dose

daily dose

Inactive ingredients

12°-30°C (59°-86°F)

Other information

children under 6 years

children 6 to under 12 years

12M mq4 - ms9 Questions? Call 1-800-824-1706 Monday - Friday

stearic acid, talc, titanium dioxide, triacetin, triethyl citrate simethicone, so dium bicarbo nate, so dium la uryl sulfate, cell ulose, polydextrose, polyethylene glycol, shellac wax,

Drug Facts (continued)

DISTRIBUTED BY: WINCO FOODS, LLC, BOISE, ID 83704

100% SATISFACTION GUARANTEE Comments & Questions? www.wincofoods.com

medical help or contact a Poison Control Centerright Keep out of reach of children. In case of overdose, get

If pregnant or breast-feeding, ask a health professional

- you need to use a laxative for more than 1 week of a serious condition.
- movement after use of a laxative. These could be signs
- you have rectal bleeding or fail to have a bowel Stop use and ask a doctor if
- it may cause stomach discomfort, faintness, and cramps
- do not use within 1 hour after taking an antacid or milk
 - do not chew or crush tablet(s) When using this product
 - sto mach pain, nause a or vomiting
- a sudden change in bowel habits that lasts more than Aska doctor before use if you have

Do not use if you cannot swallow without chewing.

Drug Facts (continued)

No Print Area Lot & Exp. Date

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

B-2150-676A-01-R REV0923A67601

Drug Facts (continued)

take with a glass of water

adults and children 12 years | take 1 to 3 tablets in a

Directions • do not take more than directed

■ this product generally produces bowel movement in 6 to ■ for relief of occasional constipation (irregularity)

sasn

Purpose

Stimulant laxative

COMPLETE PRODUCT INFORMATION KEEP OUTER PACKAGE FOR

Active ingredient Drug Facts

(in each tablet)

Drug Facts (continued)

WinCo FOODS

axative

BISACODYL USP 5 mg/STIMULANT LAXATIVE

Gentle, Overnight Relief

Winco 44-676A

Warnings Bisacodyl USP, 5 mg.....

LAXATIVE

bisacodyl tablet, delayed release

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:67091-452

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
The state of the s		

BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y) BISACODYL 5 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SHELLAC (UNII: 46N107B710)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
WATER (UNII: 059QF0KO0R)	
SODIUM BICARBONATE (UNII: 8MDF5V39QO)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

TRIETHYL CITRATE (UNII: 8Z96QXD6UM)

Color	pink	Score	no score
Shape	ROUND	Size	8mm
Flavor		Imprint Code	В
Contains			

kag

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	-------------------------	-----------------------

	NDC:67091- 452-24	2 in 1 CARTON	07/17/2024	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		
M	arketing	nformation		
M	arketing Marketing Category	nformation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

Labeler - WinCo Foods, LLC (056098817)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(67091-452)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(67091-452) , pack(67091-452)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(67091-452)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(67091-452)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(67091-452)

Revised: 7/2024 WinCo Foods, LLC