

CORICIDIN HBP COLD AND FLU- acetaminophen and chlorpheniramine maleate tablet, film coated
Bayer HealthCare LLC.

Coricidin® HBP Cold & Flu

Drug Facts

Active ingredients (in each tablet)	Purpose
Acetaminophen 325 mg	Pain reliever/ fever reducer
Chlorpheniramine maleate 2 mg	Antihistamine

Uses

- temporarily relieves
 - minor aches and pains
 - headache
 - sneezing
 - runny nose
- temporarily reduces fever

Warnings

Liver Warning

This product contains acetaminophen. Sever liver damage may occur if

- adult takes more than 12 tablets in 24 hours
- child takes more than 5 tablets in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks everyday while using this product.

Do not use with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

Ask a doctor before use if the user has

- liver disease
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if the user is

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days (children 6 to under 12 years) or 10 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning

Taking more than the recommended dose may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)

Adults and children 12 years and over 2 tablets every 4 to 6 hours, not more than 12 tablets in 24 hours

Children 6 to under 12 years of age 1 tablet every 4 to 6 hours, not more than 5 tablets in 24 hours

Children under 6 years of age ask a doctor

Other information

- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

corn starch, FD&C red No. 40 aluminum lake, lactose anhydrous, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, talc, titanium dioxide

Questions or comments?

Call 1-800-317-2165

PRINCIPAL DISPLAY PANEL - 10 Tablet Carton

Coricidin®

HBP

Decongestant-free

COLD SYMPTOM RELIEF for people with

HIGH BLOOD PRESSURE

Chlorpheniramine - Antihistamine,

Acetaminophen - Pain Reliever/Fever Reducer

See New Warnings Information

COLD

& FLU

Relieves:

- **Runny Nose**
- **Sneezing**
- **Body Aches
and Pains**
- **Fever**

10 TABLETS

Coricidin[®] HBP

COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE*

COLD & FLU 20 TABLETS
SAFETY SEALED: Do not use if blister is cut or broken.

Coricidin[®] HBP

COLD RELIEF FOR PEOPLE WITH HIGH BLOOD PRESSURE*

COLD & FLU

Acetaminophen
Pain Reliever/Fever Reducer
Chlorpheniramine Maleate
Antihistamine

- Runny Nose • Fever
- Sore Throat • Sneezing
- Body Aches & Pains

*Decongestant Free

20 TABLETS

Bayer, the Bayer Cross, Coricidin, and Coricidin HBP are registered trademarks of Bayer.
© 2020 Bayer
Mkt. by Bayer Healthcare LLC
Whispery, NJ 07081
Pat. patents (www.bayer.com) 88461517

Bayer



Drug Facts (continued)
Uses
Temporarily relieves these symptoms due to a cold or flu:
■ sore throat ■ headache
■ minor aches and pains ■ sinusitis
■ sneezing ■ temporarily reduces fever

Drug Facts (continued)
Purposes
Active ingredients (in each tablet)
Acetaminophen 325 mg
Chlorpheniramine maleate 2 mg
Pain reliever/fever reducer
Antihistamine

Drug Facts (continued)
Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
■ skin reddening ■ blisters ■ rash
■ hives ■ facial swelling
■ asthma (wheezing) ■ shock
If a skin or general allergic reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.
Do not use to sedate children
Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you have ever had an allergic reaction to this product or any of its ingredients
■ in children under 12 years of age
Ask a doctor before use if you have
■ liver disease ■ glaucoma
■ a breathing problem such as emphysema or chronic bronchitis
■ difficulty in urination due to enlargement of the prostate gland
Ask a doctor or pharmacist before use if you are
■ taking the blood thinning drug warfarin
■ taking sedatives or tranquilizers
When using this product
■ excitability may occur, especially in children

Drug Facts (continued)
■ drowsiness may occur
■ avoid alcoholic beverages
■ alcohol, sedatives and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery
Stop use and ask a doctor if
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions
■ do not take more than the recommended dose
■ adults and children 12 years and over: take 2 tablets with water every 4-6 hours. Do not exceed 10 tablets in 24 hours or as directed by a doctor.
■ children under 12 years of age: do not use
Other information
■ store between 20° to 25°C (68° to 77°F)
Inactive ingredients colloidal silicon dioxide, FD&C red #40 aluminum lake, lactose monohydrate, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, purified water, stearic acid, talc, titanium dioxide
Questions or comments?
1-800-317-2165 (Mon-Fri 9AM - 5PM EST)

88461517

CORICIDIN HBP COLD AND FLU

acetaminophen and chlorpheniramine maleate tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11523-4325
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg	
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg	

Inactive Ingredients		
Ingredient Name		Strength
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)		
STARCH, CORN (UNII: O8232NY3S)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		
ALUMINUM OXIDE (UNII: LMI26O6933)		
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)		
MAGNESIUM STEARATE (UNII: 70097M6130)		
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)		

POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL (UNII: 532B59J990)	
POVIDONE (UNII: FZ989GH94E)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	CF
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-4325-1	1 in 1 CARTON	10/01/2013	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11523-4325-2	2 in 1 CARTON	10/01/2013	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:11523-4325-3	4 in 1 CARTON	02/18/2025	
3		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/01/2013	

Labeler - Bayer HealthCare LLC. (112117283)