

**DAYTIME NIGHTTIME COLD FLU RELIEF- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl  
Safeway, Inc.**

-----  
**DRUG FACTS**

**Active ingredients for Daytime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Active ingredients for Nighttime (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose for Daytime**

**Pain reliever/fever reducer**

Cough suppressant

Nasal decongestant

**Purpose for Nighttime**

**Pain reliever/fever reducer**

Cough suppressant

Antihistamine

**Uses**

**DAYTIME**

- temporarily relieves common cold and flu symptoms:
  - cough due to minor throat and bronchial irritation
  - fever
  - sore throat
  - nasal congestion and sinus pressure
  - headache
  - minor aches and pains

**NIGHTTIME**

- temporarily relieves common cold and flu symptoms:
  - cough due to minor throat and bronchial irritation
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing

## Warnings

### DAYTIME

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### NIGHTTIME

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or following by fever, headache, rash, nausea, vomiting, consult a doctor promptly.

## Do not use

## **DAYTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **NIGHTTIME**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

## **Ask a doctor before use if you have**

### DAYTIME

- liver disease
- heart disease
- diabetes
- thyroid disease
- high blood pressure
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### NIGHTTIME

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to an enlarged prostate gland

## **Ask a doctor or pharmacist before use if you are**

### DAYTIME

taking the blood thinning drug warfarin

### NIGHTTIME

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

## **When using this product,**

### DAYTIME

## **do not exceed recommended dosage**

### **NIGHTTIME**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

#### DAYTIME

- nervousness, dizziness or sleeplessness occur
- pain, cough, and nasal congestion gets worse or lasts more than 7 days
- redness or swelling is present
- new symptoms occur
- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

#### NIGHTTIME

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash headache that lasts.

These could be a signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### DAYTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 4 hours.
- children under 12 years: do not use

## NIGHTTIME

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- swallow whole; do not crush, chew, or dissolve
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- children under 12 years: do not use

## Other information

- store between 15°-30°C (59°-86°F)
- avoid excessive heat

## Inactive ingredients

### DAYTIME

FD&C red #40, FD&C yellow #6, , gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### NIGHTTIME

D&C yellow #10, FD&C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

## Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

## Principal Display Panel

**COMPARE TO** Vicks® DayQuil™ and NyQuil™ Cold & Flu LiquiCaps™ active ingredients\*

DAYTIME MULTI-SYMPTOM

COLD & FLU RELIEF

**ACETAMINOPHEN 325 mg -**

Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 10 mg

Cough Suppressant

PHENYLEPHRINE HCl 5 mg

Nasal Decongestant

Relieves aches, fever, sore throat, cough, nasal congestion & sinus pressure

ALCOHOL-FREE

ANTIHISTAMINE-FREE

SOFTGELS\*\*

(\*\*LIQUID-FILLED CAPSULES)

**NIGHTTIME MULTI-SYMPTOM**

**COLD & FLU RELIEF**

**ACETAMINOPHEN 325 mg -**

Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 15 mg

Cough Suppressant

DOXYLAMINE SUCCINATE 6.25 mg

Antihistamine

Relieves aches, fever, sore throat, cough, runny nose, sneezing

ALCOHOL-FREE

SOFTGELS\*\*

(\*\*Liquid-Filled Capsules)

**WHEN USING DAYTIME AND NIGHTTIME PRODUCTS, CAREFULLY READ THE LABELING TO ENSURE CORRECT DOSING,**

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, DayQuil™, NyQuil™, and LiquiCaps™ are registered trademarks of the Procter and Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

DISTRIBUTED BY

BETTER LIVING BRANDS LLC

P.O. BOX 99 PLEASANTON, CA 94566-0009

**Product Label**

**COMPARE TO**  
Vicks® DayQuil™ and NyQuil™  
Cold & Flu LiquiCaps™  
active ingredients\*

NDC 21130-901-48



**DAYTIME MULTI-SYMP TOM  
COLD & FLU RELIEF**

**ACETAMINOPHEN 325 mg**  
Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 10 mg  
- Cough Suppressant  
PHENYLEPHRINE HCl 5 mg  
- Nasal Decongestant



**NIGHTTIME MULTI-SYMP TOM  
COLD & FLU RELIEF**

**ACETAMINOPHEN 325 mg**  
Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 15 mg.  
- Cough Suppressant  
DOXYLAMINE SUCCINATE 6.25 mg  
- Antihistamine

Purposes	Active ingredients (in each softgel)
Pain reliever/fever reducer	Acetaminophen 325 mg
Cough suppressant	Dextromethorphan HBr 10 mg
Nasal decongestant	Phenylephrine HCl 5 mg

Uses	Warnings
Temporarily relieves common cold and flu symptoms: cough due to minor throat and bronchial irritation, fever, sore throat, nasal congestion and sinus pressure, headache, minor aches and pains	Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: more than 4,000 mg of acetaminophen in 24 hours; with other drugs containing acetaminophen; 3 or more alcoholic drinks every day while using the product

**Warnings** Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: skin redness, blisters, rash. If a skin reaction occurs, stop use and seek medical help right away. **Safe throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. **Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Do not use** with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Daytime Cold & Flu Softgel**

**Drug Facts (continued)**  
Ask a doctor before use if you have liver disease, glaucoma, asthma, chronic bronchitis, or emphysema. Cough that occurs with too much phlegm (mucus) is a breathing problem or chronic cough that lasts such as occurs with smoking. Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. Taking the blood thinning drug warfarin. When using this product, do not exceed recommended dosage. **Stop use and ask a doctor if** pain or cough gets worse or lasts more than 7 days; you get worse or lasts more than 3 days; fever gets worse or lasts more than 3 days; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. **Keep out of reach of children.** Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** Do not take more than directed (see Overdose warning). Adults and children 12 years and over: take 2 softgels with water every 6 hours. Children under 12 years: do not use.

**Drug Facts (continued)**  
Ask a doctor before use if you have liver disease, glaucoma, asthma, chronic bronchitis, or emphysema. Cough that occurs with too much phlegm (mucus) is a breathing problem or chronic cough that lasts such as occurs with smoking. Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. Taking the blood thinning drug warfarin. When using this product, do not exceed recommended dosage. **Stop use and ask a doctor if** pain or cough gets worse or lasts more than 7 days; you get worse or lasts more than 3 days; fever gets worse or lasts more than 3 days; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. **Keep out of reach of children.** Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** Do not take more than directed (see Overdose warning). Adults and children 12 years and over: take 2 softgels with water every 6 hours. Children under 12 years: do not use.

**Drug Facts (continued)**  
Ask a doctor before use if you have liver disease, glaucoma, asthma, chronic bronchitis, or emphysema. Cough that occurs with too much phlegm (mucus) is a breathing problem or chronic cough that lasts such as occurs with smoking. Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. Taking the blood thinning drug warfarin. When using this product, do not exceed recommended dosage. **Stop use and ask a doctor if** pain or cough gets worse or lasts more than 7 days; you get worse or lasts more than 3 days; fever gets worse or lasts more than 3 days; cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. **Keep out of reach of children.** Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. **Directions** Do not take more than directed (see Overdose warning). Adults and children 12 years and over: take 2 softgels with water every 6 hours. Children under 12 years: do not use.

• Relieves aches, fever, sore throat, cough, nasal congestion & sinus pressure

**ALCOHOL-FREE**  
**ANTIHISTAMINE-FREE**

Actual Size

WHEN USING DAYTIME AND NIGHTTIME PRODUCTS, CAREFULLY READ THE LABELING TO ENSURE CORRECT DOSING.

**32 SOFTGELS\*\***  
(\*\*UNDOSE-FILLED CAPSULES)

• Relieves aches, fever, sore throat, cough, runny nose, sneezing

**ALCOHOL-FREE**

Actual Size

WHEN USING DAYTIME AND NIGHTTIME PRODUCTS, CAREFULLY READ THE LABELING TO ENSURE CORRECT DOSING.

**16 SOFTGELS\*\***  
(\*\*UNDOSE-FILLED CAPSULES)

48 TOTAL SOFTGELS



2 21130 76351 3

8025294 52801

Scan here for more information



smartlabel

PRODUCT OF CHINA

+1-888-723-3929

PLEASANTON, CA 94566-0009

**DISTRIBUTED BY:**  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94566-0099



KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

TAMPER EVIDENT. DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.




www.BetterLivingBrands.org

This product is not manufactured or distributed by The Procter & Gamble Company. Vicks<sup>®</sup>, DayQuil<sup>™</sup>, NyQuil<sup>™</sup>, and LiquiCaps<sup>™</sup> are registered trademarks of The Procter and Gamble Company.

<p><b>Daytime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> FD&amp;C red #10, FD&amp;C yellow #6, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b> Call 1-877-753-3929 Monday-Friday 9AM-5PM EST</p>	<p><b>Nighttime Cold &amp; Flu Softgel</b></p> <p><b>Drug Facts (continued)</b></p> <p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ store between 15-30°C (59-86°F)</li> <li>■ avoid excessive heat</li> </ul> <p><b>Inactive ingredients</b> D&amp;C yellow #10, FD&amp;C blue #1, gelatin, glycerin, lecithin, light mineral oil, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink</p> <p><b>Questions or comments?</b> Call 1-877-753-3929 Monday-Friday 9AM-5PM EST</p>
--	--

## SIGNATURE SELECT Daytime Nighttime Multi-Symptom Cold and Flu Relief

### DAYTIME NIGHTTIME COLD FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl kit

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:21130-901
---------------------	----------------	---------------------------	---------------

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-901-48	1 in 1 KIT; Type 1: Convenience Kit of Co-Package	12/27/2019	12/27/2026

#### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	16 BLISTER PACK	16
Part 2	32 BLISTER PACK	32

# NIGHTTIME COLD AND FLU RELIEF

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

## Product Information

**Item Code (Source)** NDC:21130-899

**Route of Administration** ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	

## Product Characteristics

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC10
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		16 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/27/2019	

## Part 2 of 2

### COLD AND FLU DAYTIME

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

## Product Information

Item Code (Source)	NDC:21130-598
Route of Administration	ORAL

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>LECITHIN, SOYBEAN</b> (UNII: 1DI56QDM62)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>LIGHT MINERAL OIL</b> (UNII: N6K5787QVP)	

## Product Characteristics

Color	orange	Score	no score
-------	--------	-------	----------

<b>Shape</b>	CAPSULE	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	PC9
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		32 in 1 CARTON		
1		1 in 1 BLISTER PACK; Type 1: Convenience Kit of Co-Package		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/27/2019	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/27/2019	12/27/2026

**Labeler** - Safeway, Inc. (009137209)

Revised: 12/2025

Safeway, Inc.