BZK ANTISEPTIC - benzalkonium chloride swab Aero-Med Ltd

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BZK Towelettes

Active Ingredient Purpose

Benzalkonium Chloride 0.13% v/v Antiseptic

Use: towelette

- First aid antiseptic to help prevent skin infection in minor cuts, scrapes and burns
- Antiseptic cleansing
- Perineal and maternity care

Warnings

For external use only

Indications and Usage

General antiseptic

Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

Stop use if

Stop Use if:

- irritation and redness develop
- if condition persists for more than 72 hours, consult a physician.

Directions

Directions:

• Tear at notch, remove towelette, use only once

As a first aid antiseptic

- clean affeected area
- apply 1 to 3 times daily
- may be covered with a sterile bandage
- if bandaged let dry first

Keep out of reach of children

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away.

Do not use

Do not use

- as a first aid antiseptic for more than 1 week
- in the eyes
- over large areas of the body

Inactive Ingredient

Inactive Ingredient: Water

Principal Display Panel

Aeromed Benzalkonium Chloride Antiseptic Towelette

Aeromed BZK.jpg



BENZALKONIUM CHLORIDE ANTISEPTIC TOWELETTE 100 COUNT

Distributed by: AEROMED Glastonbury, CT 06033

Drug Facts	
Active ingredient Purpo Benzalkonium Chloride, 0.13% v/v. Antisep	
Uses • first aid antiseptic to help prevent skin infection in minor cuts, scrapes and burns • Antiseptic cleaning • Perineal and mate	rnity care
Warnings • For external use only	
Do not use • as a first aid antiseptic for more than one week • in the eyes • over large areas of the body	
Ask a doctor before use if you have • deep puncture wounds • animal bites • serious burns	
Stop use if irritation and redness develop. If condition persists for more than 72 hours, consult a physician.	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions • Tear at notch • remove towelette • use only once	
As a first aid antiseptic	
 clean affected area apply 1 to 3 times daily may be covered with a sterile bandage if bandaged, let dry first 	
Other Information • store at room temperature • avoid excessive heat	
Inactive ingredients water	

Made in China

BZK ANTISEPTIC

benzalkonium chloride swab

D		TC		. •
Prod	liict	Into	rma	tion

Product Type HUMAN OTC DRUG Item Code (Source) NDC:42449-210

n

TOPICAL

Active ingredictionative workly				
Ingredient Name	Basis of Strength	Strength		
BENZALKO NIUM CHLO RIDE (UNII: F5UM2KM3W7) (BENZALKO NIUM - UNII:7N6 JUD5X6 Y)	BENZALKONIUM CHLORIDE	1 g in 750 g		

Inactive Ingredients

Ingredient Name	Strength

WATER (UNII: 059QF0KO0R)

P	Packaging					
#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:42449-210-13	10 in 1 CASE				
1		100 in 1 BOX				
1		3 g in 1 PACKET				

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC monograph not final	part333A	02/02/2012		

Labeler - Aero-Med Ltd (194550042)

Revised: 5/2012 Aero-Med Ltd