GOOD NEIGHBOR PHARMACY PAIN AND FEVER- acetaminophen suspension Amerisource Bergen

Amerisource Bergen Infants' Pain & Fever Drug Facts

Active ingredient (in each 5 mL)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of it's ingredients

Ask a doctor before use if your child has

liver disease

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- new symptoms occur
- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- push air out of syringe. Insert syringe tip into bottle opening.
- flip bottle upside down. Pull yellow part of syringe to the first dose line and then push product back into bottle.
- pull yellow part of syringe until it reaches and stays at the correct dose
- dispense liquid slowly into child's mouth, toward inner cheek
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours
- replace cap tightly to maintain child resistance

Dosing Chart			
Weight (lb)	Age (yr)	Dose (mL)*	
under 24	under 2 years	ask a doctor	

24-35	2-3 years	5 mL

*or as directed by a doctor

Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other Information

- store at 20-25°C (68-77°F)
- do not use if printed bottle wrap is broken or missing

Inactive ingredients

anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments?

1-800-719-9260

Package/Label Principal Display Panel

Compare to Infants' Tylenol® active ingredient

GOOD NEIGHBOR PHARMACY®

FOR AGES 2 TO 3 YEARS

infants' Pain & Fever

acetaminophen 160 mg per 5 mL

oral suspension

Fever Reducer - Pain Reliever

Aspirin Free • Ibuprofen Free

Alcohol Free • Paraben Free

Use only with enclosed syringe.

See side panel for more information.

Grape Flavor

1 fl oz (30 mL)





■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe live damage may occur if your child takes

more than 5 doses in 24 hours, which is the maximum daily amount

with other drugs containing acetaminophen
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: me skin reddening me blisters me rash if a skin reaction occurs, stop use and seek medical help right away. Sore threat warning: if sore throat is server, persists for more than 2 days, is accompanied or followed by fever, headache, rash, neusea, or vomiting, consult a doctor promptly.

Do not use with any other drug containing acetaminophen Do not use I will ally consulting overalling overalling overalling overalling or operation or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. I if your child has ever had an allergic reaction to this product or any of its ingredients. Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if ■ new symptoms occur

■ pain gets worse or lasts more than 5 days ■ fever gets worse or lasts more than 3 days

■ redness or swelling is present. These could be signs of a serious condition Geep out of reach of children.

Neep out or reach or contraers. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions this product does not contain directions or complete warnings for adult use do not give more than directed (see overdose warning) shake well before using ml = milliliter information from the milliliter formation from the milliliter for

syringe to the first dose line and then push product back into bottle.

Drug Facts (continued)

and practs (commuted)

pull yellow part of syringe until it
reaches and stays at the correct dose
■ dispense liquid stowly into child's
mouth, toward inner cheek
■ repeat
dose every 4 hours white symptoms
last
■ do not give more than 5 times
is 34 hours.
■ replaces on the thirty. in 24 hours replace cap tightly to maintain child resistance

Dosing Chart			
Weight (lb)	Age (yr)	Dose (mL)*	
under 24	Age (yr) under 2 years 2-3 years	ask a doctor	
24-35	2-3 years	5 mL	

* or as directed by a doctor Attention: use only enclosed syringe specifically designed for use with this product. Do not use any other dosing device.

Other information

store at 20-25°C (68-77°F)

do not use if printed bottle wrap is broken or missing

Inactive ingredients

mactive ingredients anhydrous citric acid, calcium sulfate, carrageenan, D&C red #33, FD&C blue #1, flavor, glycerin, high fructose corn syrup, hydroxyethyl callulose, microcrystalline callulose and carbonymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, tribasic sodium phosphate

Questions or comments? 1-900-719-9260

**Good Neighbor Pharmacy® Infants' Pain & Fever is not manufactured or distributed by Johnson & Johnson Consumer Inc... distributor of Consumer Inc., distributor of Infants' Tylen ol®.



infants'

Infants' Tylenol' active ingredient"





acetaminophen 160 mg per 5 mL oral suspension

Fever Reducer – Pain Reliever

Aspirin Free • Ibuprofen Free Alcohol Free • Paraben Free

Use only with enclosed syringe See side panel for more information.

Grape Flavor

1 fl oz (30 m

IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose.

STEP 1. See Directions for the correct dose for your child.

STEP 2. Shake well before using.

STEP 3. Push air out of syringe, insert syringe tip into bottle opening.



STEP 4. Hip bottle upside down. Pull yellow part of syninge to the first dose line and then push product back into bottle

STEP 5. Pull yellow part of syringe until it reaches and stays at the correct dose.



STEP 6. Dispense liquid slowly into child's mouth.

Distributed By AmerisourceBergen 1 West First Avenue Conshohocken, PA 19428 Questions or Concerns?





94610 29 C5

GOOD NEIGHBOR PHARMACY PAIN AND FEVER

acetaminophen suspension

Product Information

HUMAN OTC DRUG **Item Code (Source)** NDC:46122-042 **Product Type**

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name Basis of Strength Strength

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) ACETAMINOPHEN 160 mg in 5 mL

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED FORM (UNII: K6790BS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	

Product Characteristics		
Color	PURPLE	Score
Shape		Size
Flavor		Imprint Code
Contains		

F	Packaging				
#	tem Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:46122-042- 03	1 in 1 CARTON	08/28/2011		
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	08/28/2011	

Labeler - Amerisource Bergen (007914906)

Revised: 4/2024 Amerisource Bergen