

**ACETAMINOPHEN- acetaminophen tablet**  
**Rugby Laboratories**

-----  
**5427-Rugby**

***Drug Facts***

***Active ingredient (in each tablet)***

Acetaminophen 500 mg

***Purpose***

Pain reliever/fever reducer

***Uses***

Temporarily reduces fever and relieves minor aches and pains caused by

- headache
- muscular aches
- the common cold
- toothache
- backache
- minor pain of arthritis
- premenstrual and menstrual cramps

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not take more than directed (see overdose warning)**

**adults and children 12 years and over:**

- take 2 tablets every 6 hours while symptoms last
- do not take more than 6 tablets in 24 hours, unless directed by a doctor
- do not take for more than 10 days unless directed by a doctor

**children under 12 years:** ask a doctor

***Other information***

store in a dry place at 15° – 30°C (59° – 86°F).

Povidone, pregelatinized starch, sodium starch glycolate, stearic acid

***Questions or comments?***

**Call 1-800-231-4670**

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*Rugby Laboratories is not affiliated with the owner of the registered trademark Tylenol®.

**Distributed by:  
RUGBY® LABORATORIES**

Indianapolis, IN 46268

Questions or comments?

Call (800) 616-2471

www.majorpharmaceuticals.com

Rugby

COMPARE TO THE ACTIVE INGREDIENT IN EXTRA STRENGTH TYLENOL®\*

NDC 0536-1172-01

Extra Strength Acetaminophen

500mg each

Pain Reliever/Fever Reducer

Aspirin Free

100 Tablets

FRONT OF LABEL

**Rugby**  
NDC 0536-1172-01

COMPARE TO ACTIVE INGREDIENT IN EXTRA STRENGTH TYLENOL®

EXTRA STRENGTH  
**Acetaminophen**  
500 mg Each  
Aspirin Free  
Extra Strength Pain Reliever /  
Fever Reducer  
**100 TABLETS**

**Drug Facts**  
**Active ingredient (in each tablet)** Purpose  
Acetaminophen 500 mg.....Pain reliever/fever reducer

**Uses** Temporarily reduces fever and relieves minor aches and pains caused by • headache • muscular aches • the common cold • toothache • premenstrual and menstrual cramps • backache • minor pain of arthritis

**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:  
• more than 4,000 mg of acetaminophen in 24 hours  
• with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:  
• skin reddening • blisters • rash  
If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • If you are allergic to acetaminophen or any of the inactive ingredients in this product

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

**Drug Facts (continued under label)**

3 05361 17201 5

Distributed by:  
**RUGBY® LABORATORIES** Call (800) 616-2471  
Indianapolis, IN 46268 www.majorpharmaceuticals.com  
R-29 Re-order No. 370597 Rev. 10/23 R52189

PEEL HERE

6.0"

INSIDE OF LABEL

**Drug Facts (continued)**

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**  
• do not take more than directed (see overdose warning)

**adults and children 12 years and over:**  
• take 2 tablets every 6 hours while symptoms last  
• do not take more than 6 tablets in 24 hours, unless directed by a doctor • do not take for more than 10 days unless directed by a doctor

**children under 12 years:** ask a doctor

**Other information** store in a dry place at 15° – 30°C (59° – 86°F).

**Inactive ingredients** povidone, pregelatinized starch, sodium starch glycolate, stearic acid

**Questions or comments?**  
Call 1-800-231-4670

**NO COPY  
ADHESIVE ZONE**

# ACETAMINOPHEN

acetaminophen tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0536-1172
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

## Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
STARCH, CORN (UNII: O8232NY35J)	
POVIDONE K30 (UNII: U725QWY32X)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	

## Product Characteristics

<b>Color</b>	white	<b>Score</b>	2 pieces
<b>Shape</b>	ROUND	<b>Size</b>	12mm
<b>Flavor</b>		<b>Imprint Code</b>	54;27
<b>Contains</b>			

## Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:0536-1172-10	1000 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	11/30/2025
2	NDC:0536-1172-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	10/01/2018	

## Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M013	10/01/2018	

**Labeler** - Rugby Laboratories (079246066)