# ESOMEPRAZOLE MAGNESIUM- esomeprazole magnesium capsule, delayed release RUGBY LABORATORIES

-----

#### **Drug Facts**

# Active ingredient(s) in each capsule

\*Esomeprazole 20 mg

(Each delayed-release capsule corresponds to 20.7 mg esomeprazole magnesium, USP)

### Purpose

Acid reducer

#### Uses

- treats frequent heartburn (occurs **2 or more** days a week)
- not intended for immediate relief of heartburn; this drug may take 1 to 4 days for full effect

# Warnings

Allergy alert: Do not use if you are allergic to esomeprazole

#### Do not use if you have:

- trouble or pain swallowing food, vomiting with blood, or bloody or black stools
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadness
- frequent chest pain.

These may be signs of a serious condition. See your doctor.

#### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain

## Ask a doctor or pharmacist before use if you are taking

- warfarin, clopidogrel or cilostazol (blood-thinning medicines)
- prescription ant fungal or anti-yeast medicines
- digoxin (heart medicine)
- diazepam (anxiety medicine)
- tacrolimus or mycophenolate mofetil (immune system medicine)
- prescription antiretrovirals (medicines for HIV infection)
- methotrexate (arthritis medicine)

#### Stop use and ask doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

- you need to take more than 1 course of treatment every 4 months
- you get diarrhea
- you develop a rash or joint pain

**If pregnant or breast-feeding,** ask a health professional before use.

# Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

#### **Directions**

- adults 18 years of age and older
- this product is to be used once a day (every 24 hours), every day for 14 days
- may take 1 to 4 days for full effect

#### 14-Day Course of Treatment

- swallow 1 capsule with a glass of water before eating in the morning
- take every day for 14 days
- do not take more than 1 capsule a day
- swallow whole. Do not crush or chew capsules.
- do not use for more than 14 days unless directed by your doctor

#### Repeated 14-Day Courses (if needed)

- you may repeat a 14-day course every 4 months
- do not take for more than 14 days or more often than every 4 months unless directed by a doctor
- children under 18 years of age: ask a doctor before use. Heartburn in children may sometimes be caused by a serious condition.

#### Other information

- each capsule contains: magnesium, 15 mg
- read directions and warnings before use
- keep the carton. It contains important information
- store at 20 to 25°C (68 to 77°F)

#### **Inactive ingredients**

acetyl tributyl citrate, dibutyl sebacate, FD&C Blue #1, ferroso ferric oxide, gelatin, glyceryl monostearate, hypromellose, magnesium oxide, magnesium stearate, methacrylic acid copolymer dispersion, poloxamer, polysorbate 80, povidone, potassium hydroxide, propylene glycol, shellac, sugar, talc, titanium dioxide

#### Questions or comments

call toll-free weekdays 8 AM to 10 PM EST at 1-888-375-3784

#### Tips for Managing Heartburn

- Avoid foods or drinks that are more likely to cause heartburn, such as rich, spicy, fatty and fried foods, chocolate, caffeine, alcohol and even some acidic fruits and vegetables.
- Eat slowly and do not eat big meals.
- Do not eat late at night or just before bedtime.

- Do not lie flat or bend over soon after eating.
- Raise the head of your bed.
- Wear loose-fitting clothing around your stomach.
- If you are overweight, lose weight.
- If you smoke, quit smoking.

#### **Bottle Label**

Container Label:



# Reseal Area

Orug Facts (continued with heartburn = unexplained 3 months. This may be a sign frequent wheezing, particular ou have m had heartburn o chest pain or shoulder pai These may be signs of a ser irms, neck or shoulders; or lsk a doctor before use i condition. See your doctor ifth shortness of breath; a more serious condition. frequent chest pain ghilheadedness

eight loss = nausea or vorni storrach pain

> this product is to be used once a day (every 24 hours), every day for 14 day

14-Day Course of Treatment

adults 18 years of age and older

Orug Facts (continued)

isk a doctor or pharmacis defore use if you are = tak prescription drug. Acid redumay interact with certain

do not také môre than 1 capsule a day water before eating in the morning swallow 1 capsule with a glass of take every day for 14 days

swallow whole. Do not crush or chev

unless directed by your doctor repeated 14-Day Courses (If needed every 4 months ado not take for more than 14 days or more often than every 4 months do not use for more than 14 days you may repeat a 14-day course unless directed by a doctor

Stop use and ask a doctor li

worsens - you need to take

vour heartburn continues

product for more than 14 day you need to take more that

course of treatment every

children under 18 years of age: ask a doctor before use. Heartburn in children

may some times be caused by a serious

you develop a rash or joint

Other Information = each capsule Information. ■ store at 20 to 25°C (68 to 77°F) read the directions and warnings before use keep the carton. It contains important acetyl tributyl cifrate, dibutyl sebaca FD&C Blue #1, ferroso ferric oxide nactive ingredients contains: magnesium, 15 mg

elp or contact a Polson Con

eepout of reach of childr case of overdose, get med

health professional before

Questions or comments? call toil-free weekdays 8 AM to 10 PM EST 1-888-375-3784 hydroxide, propylene glycol, shellac, sugar, talc, titarium dioxide

magnesium stearate, methacrylic add

hypromellose, magnesium oxide

gelatin, glyceryl monostearate

polysórbate 80, povidóne, potassium

copolymer dispersion, poloxamer



#### Carton

Carton



#### **ESOMEPRAZOLE MAGNESIUM**

esomeprazole magnesium capsule, delayed release

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0536-1230(NDC:43598-407)	
Route of Administration	ORAL			

Active Ingredient/Active Molety			
Ingredient Name	Basis of Strength	Strength	
Esomeprazole Magnesium (UNII: R6DXU4WAY9) (Esomeprazole - UNII:N3PA6559FT)	Esomeprazole	20 mg	

Inactive Ingredients		
Ingredient Name	Strength	
acetyltributyl citrate (UNII: 0ZBX0N59RZ)		
Dibutyl Sebacate (UNII: 4W5IH7FLNY)		
Ferrosoferric Oxide (UNII: XM0 M87F357)		
Gelatin (UNII: 2G86QN327L)		
Glyceryl Monostearate (UNII: 230 O U9 XXE4)		
Hypromelloses (UNII: 3NXW29 V3WO)		
Magnesium Oxide (UNII: 3A3U0GI71G)		
Magnesium Stearate (UNII: 70097M6I30)		
Methacrylic Acid - Ethyl Acrylate Copolymer (1:1) Type A (UNII: NX76LV5T8J)		
Poloxamer 188 (UNII: LQA7B6G8JG)		
Polysorbate 80 (UNII: 6OZP39ZG8H)		
Povidone K30 (UNII: U725QWY32X)		
Talc (UNII: 7SEV7J4R1U)		
Titanium Dioxide (UNII: 15FIX9 V2JP)		
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		

Potassium Hydroxide (UNII: WZH3C48M4T)	
Shellac (UNII: 46N107B71O)	
Sucrose (UNII: C151H8 M554)	
Fd&C Blue No. 1 (UNII: H3R47K3TBD)	

Product Characteristics			
Color	BLUE (light blue cap), BLUE (dark blue body)	Score	no score
Shape	CAPSULE	Size	16 mm
Flavor		Imprint Code	RDY;327
Contains			

Packaging				
#	Item Code	Package Description	<b>Marketing Start Date</b>	Marketing End Date
1	NDC:0536-1230-88	1 in 1 CARTON	05/28/2019	
1		14 in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:0536-1230-13	3 in 1 CARTON	05/28/2019	
2		14 in 1 BOTTLE; Type 0: Not a Combination Product		
3	NDC:0536-1230-71	2 in 1 CARTON	05/28/2019	
3		14 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207673	05/28/2019	

# Labeler - RUGBY LABORATORIES (079246066)

Revised: 5/2019 RUGBY LABORATORIES