

MILK OF MAGNESIA- magnesium hydroxide suspension
Cardinal Health 107, LLC

Milk of Magnesia

saline laxative/antacid

DESCRIPTION

Active Ingredient

(in each 5 mL = 1 teaspoonful)

Magnesium Hydroxide.....400 mg

Inactive Ingredients

anhydrous citric acid, benzyl alcohol, methylparaben, propylene glycol, propylparaben, purified water.

FOR USE AS A LAXATIVE

INDICATIONS

Uses

- For relief of occasional constipation (irregularity)
- This product generally produces bowel movement in 1/2 to 6 hours.

WARNINGS

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. This product may interact with certain prescription drugs.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

DIRECTIONS

- **shake well before use**
- drink a full glass (8 oz) of liquid with each dose
- mL = milliliter

Adults and children 12 years of age and older	30 mL to 60 mL
Children 6 to 11 years of age	15 mL to 30 mL
Children under 6 years of age	Consult a physician.

or as directed by a physician.

- The dose may be taken as a single daily dose or in divided doses.

FOR USE AS AN ANTACID

INDICATIONS

Uses

For the relief of

- heartburn
- sour stomach
- acid indigestion
- and upset stomach associated with these symptoms

WARNINGS

Do not take more than 12 teaspoonfuls in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician

May have laxative effect

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

DIRECTIONS

- **shake well before use**
- mL = milliliter
- adults and children 12 years old and older, take 1 to 3 teaspoonfuls (5 to 15 mL) four times a day or as directed by a physician
- do not take more than 12 teaspoonfuls (60 mL) in 24-hour period
- do not use the maximum dosage for more than 2 weeks
- children under 12 years: consult a physician

KEEP OUT OF REACH OF CHILDREN

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

HOW SUPPLIED

Milk of Magnesia USP 400 mg/5 mL is a white opaque liquid supplied as follows:

Overbagged with 5 x 30 mL unit-dose cups per bag, NDC 55154-4061-5

STORAGE

Other Information

- Protect from freezing
- Store at 20° to 25°C (68° to 77°F)

Questions or Comments?

Call 1-888-655-1505 (toll-free)

Distributed by:

VistaPharm, Inc.

Largo, FL 33771, USA

**Distributed by:
Cardinal Health**

Dublin, OH 43017

L55273200519

VP2152R1

09/17

Principal Display Panel

Milk of Magnesia, USP

2400 mg Magnesium Hydroxide per 30 mL

5 Cups



NDC 55154-4061-5

A101

MILK OF MAGNESIA, USP

2400 mg Magnesium Hydroxide per 30 mL

5 CUPS

Delivers 30 mL

Shake Well

Usual Dosage: See product insert for prescribing information, precautions and warnings.

STORAGE: Store at 20°- 25° C (68°-77° F); excursions permitted to 15°-30° C (59°-86° F).

[See USP Controlled Room Temperature]

Protect from Freezing

WARNING: This Unit Dose package is not child resistant and is Intended for Institutional Use Only. Keep this and all drugs out of the reach of children.

XACT-DOSE™

Manufactured by:
VistaPharm
Largo, FL 33771, USA

Distributed by Cardinal Health
Dublin, OH 43017
L55273200519

LOT #: XXXXXXXXXX EXP. DATE: XX/XX/XX

MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55154-4061(NDC:66689-053)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
HYPROMELLOSE 2910 (4000 MPA.S) (UNII: RN3152OP35)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
POTASSIUM CITRATE (UNII: EE90ONI6FF)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
CYCLOMETHICONE 4 (UNII: CZ227117JE)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SORBITOL (UNII: 506T60A25R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor	SPEARMINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55154-4061-5	5 in 1 BAG	04/19/2018	
1		30 mL in 1 CUP, UNIT-DOSE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	04/19/2018	

Labeler - Cardinal Health 107, LLC (118546603)