

**BOVAGUARD BARRIER- iodine solution solution**  
**AgroChem USA, LLC.**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----  
**BOVAGUARD BARRIER**

**BOVAGUARD BARRIER**

**dip cup**

**NPE FREE**

**WARNING**

**KEEP OUT OF REACH OF CHILDREN.**

**AVOID FREEZING.**

**KEEP OUT OF DIRECT SUNLIGHT.**

**CAUSES SERIOUS EYE IRRITATION.**

**FOR FARM AND INDUSTRIAL USE ONLY.**

**PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.**

**FIRST AID / RESPONSE:**

**IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.**

**STORAGE: Store locked up.**

**DISPOSAL: Dispose of contents/ container to an approved waste disposal plant .**

**EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494**

**ACTIVE INGREDIENT: Iodine 1.0 % (10% Polyethoxy Polypropoxy Polyethoxy Ethanol - Iodine Complex) Titration via USP33 Reissue, Topical Solution**

**EMOLLIENTS: 10.0%**

**BovaGuard Barrier is made with IoPlex technology and is NPE free.**

**NET CONTENTS: 55 Gallons**

**NDC: 17307-3170-3**

**LOT:**

**EXP. DATE:**

**Use BovaGuard Barrier to reduce the spread of mastitis causing organisms. BovaGuard Barrier is an effective and economical 1% iodine barrier teat dip which is both non-drip and highly water resistant. Recommended for post milking applications.**

**Directions:**

**General Use**

**Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.**

**Use a teat dip cup to apply. Clean dip cup just prior to use.**

**If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.**

**If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.**

**If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.**

**Teat Sprays can also be foamed or sprayed using recommended equipment.**

**Pre Dipping Use**

**Perform forestripping and inspect teats for clinical mastitis.**

**Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels**

**Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.**

**Allow product to stand for 30 seconds.**

**Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.**

**Post Dipping Use**

**Immediately after each milking, use product.**

**Dip each teat full-length into the teat dip cup containing product. Allow to**

air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

### Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

Manufactured By:

AgroChem USA, LLC 26 Freedom Way • Saratoga Springs, NY 12866 • (518)226-4850



## WARNING

AVOID FREEZING.  
KEEP OUT OF DIRECT SUNLIGHT.  
CAUSES SERIOUS EYE IRRITATION.  
KEEP OUT OF REACH OF CHILDREN.  
FOR FARM AND INDUSTRIAL USE ONLY.



**PREVENTION:** Wash skin thoroughly after handling. Wear eye protection/ face protection.

**FIRST AID/ RESPONSE:**

**IF IN EYES:** Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do.  
Continue rinsing. If eye irritation persists: Get medical advice/ attention.

**STORAGE:** Store locked up.

**DISPOSAL:** Dispose of contents/ container to an approved waste disposal plant.

**EMERGENCY TELEPHONE NUMBER (Chemtrec):** 800-424-9300 (CN205494)

**ACTIVE INGREDIENTS:**

Iodine – 1.0 % (10.0 % Polyethoxy Polypropoxy Polyethoxy Ethanol – Iodine Complex)  
Titration via USP33 Reissue, Topical Solution  
**EMOLLIENTS:** – 10%

BovaGuard Barrier is made with loPlex technology and is NPE free.

NET CONTENTS: 55 GALLON

NDC: 17307-3170-3



LOT:

EXP DATE:

Use BovaGuard Barrier to reduce the spread of mastitis-causing organisms. BovaGuard Barrier is an effective and economical 1% iodine barrier teat dip which is both non-drip and highly water resistant. Recommended for post milking applications.

**Directions:**

General Use

Ready to use. Do not dilute. Do not mix product with water, any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

Teat Sprays can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.



# AgroChem

Manufactured By: AgroChem USA, LLC 26 Freedom Way - Saratoga Springs, NY 12866 - (518)226-4850

## BOVAGUARD BARRIER

iodine solution solution

### Product Information

<b>Product Type</b>	OTC ANIMAL DRUG	<b>Item Code (Source)</b>	NDC:17307-3170	
<b>Route of Administration</b>	TOPICAL			
<b>Active Ingredient/Active Moiety</b>				
<b>Ingredient Name</b>		<b>Basis of Strength</b>	<b>Strength</b>	
IODINE (UNII: 9679TC07X4) (IODINE - UNII:9679TC07X4)		IODINE	0.01 kg in 1 kg	
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:17307-3170-1	19 kg in 1 CONTAINER		
2	NDC:17307-3170-2	57 kg in 1 DRUM		
3	NDC:17307-3170-3	208 kg in 1 DRUM		
4	NDC:17307-3170-4	1041 kg in 1 CONTAINER		
<b>Marketing Information</b>				
<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>	
unapproved drug other		01/03/2022		

**Labeler** - AgroChem USA, LLC. (118336116)

**Registrant** - AgroChem USA, LLC. (118336116)

### Establishment

Name	Address	ID/FEI	Business Operations
AgroChem USA, LLC.		118336116	api manufacture, manufacture, analysis, label, pack

Revised: 1/2022

AgroChem USA, LLC.