

**NIGHTTIME COLD AND FLU MAXIMUM STRENGTH- acetaminophen,
diphenhydramine hci, phenylephrine hci liquid
Dolgenercorp, Inc. (DOLLAR GENERAL & REXALL)**

Drug Facts

Active ingredients (in each 20 mL)

Acetaminophen 650 mg

Diphenhydramine HCl 25 mg

Phenylephrine HCl 10 mg

Purpose

Pain reliever/fever reducer

Antihistamine / cough suppressant

Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor ache and pains
 - sore throat
 - headache
 - runny nose
 - sneezing
- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine even one used on the skin
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash, or headache that lasts These could be signs

of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 6 doses in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter
- dose as follows or as directed by a doctor
- adults and children 12 years and older :
 - 20 mL every 4 hours while symptoms last
- children under 12 years of age do not use

Other information

- **each 20 mL contains:** sodium 12 mg
- store between 20-25°C (68-77F). Do not refrigerate

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

Principal Display Panel

Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu*

Maximum Strength

Fast Acting

Night Time Cold & Flu

Multi-Symptom Relief

Acetaminophen

Diphenhydramine HCl

Phenylephrine HCl

Pain Reliever/Fever Reducer

Antihistamine/Cough Suppressant

Nasal Decongestant

- For ages 12 years and over

FL OZ (mL)

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY OLD EAST MAIN CO.

100 MISSION RIDGE

GOODLETTSVILLE, TN 37072

Product Label

DG™ health

Compare to the active ingredients of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu*

Maximum Strength Fast Acting Night Time Cold & Flu

Multi-Symptom Relief Acetaminophen

Diphenhydramine HCl
Phenylephrine HCl

Pain Reliever/Fever Reducer
Antihistamine/Cough Suppressant
Nasal Decongestant

• For ages 12 years and over

DISTRIBUTED BY OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100* Satisfaction Guaranteed! (888) 309-9030

6 fl oz (177 mL)

PLD-B377D LB008187

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Night Time Cold & Flu.

PLD-C377D
LB003148

Drug Facts

Active ingredients (in each 20 mL)

Active ingredients (in each 20 mL)	Purposes
Acetaminophen 650 mg.....	Pain reliever/fever reducer
Diphenhydramine HCl 25 mg.....	Antihistamine/cough suppressant
Phenylephrine HCl 10 mg.....	Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms
 - cough
 - nasal congestion
 - minor aches and pains
 - sore throat
 - headache
 - runny nose
 - sneezing

PEEL CORNER FOR MORE DRUG FACTS ▲

Drug Facts (continued)

- temporarily reduces fever
- controls cough to help you get to sleep

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- with any other drug containing diphenhydramine, even one used on the skin

Drug Facts (continued)

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if

- you are taking the blood thinning drug warfarin
- you are taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children

Drug Facts (continued)

- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse, or lasts more than 7 days
- fever gets worse, or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- dose as follows or as directed by a doctor
- adults and children 12 years and older: 20 mL every 4 hours while symptoms last
- children under 12 years of age: do not use

Other information

- each 20 mL contains: sodium 12 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

PEEL CORNER FOR MORE DRUG FACTS ▲

DOLLAR GENERAL HEALTH Mucus Relief Night Time Cold and Flu

NIGHTTIME COLD AND FLU MAXIMUM STRENGTH

acetaminophen, diphenhydramine hci, phenylephrine hci liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-460
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 20 mL
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	25 mg in 20 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
PROPYL GALLATE (UNII: 8D45NN7V92)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
EDETATE CALCIUM DISODIUM (UNII: 25IH6R4SGF)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-460-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/31/2018	03/31/2027

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/31/2018	03/31/2027

Revised: 7/2025

Dolgencorp, Inc. (DOLLAR GENERAL & REXALL)