LORATADINE- loratadine tablet, orally disintegrating Topcare Associates LLC

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Loratadine USP, 10 mg

PURPOSE

Antihistamine

USES

Temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

WARNINGS

Do not use

If you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have

Liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product

Do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if

An allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

• place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

OTHER INFORMATION

- Phenylketonurics: Contains Phenylalanine 0.6 mg Per Tablet.
- TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF TAMPERING.
- store between 20° to 25° C (68° to 77° F). Protect from excessive moisture.
- keep in a dry place.
- use tablet immediately after opening individual blister.

INACTIVE INGREDIENTS

Aspartame, croscarmellose sodium, magnesium stearate, mannitol, mint flavor, sodium stearyl fumarate, strawberry cream flavor, tutti-frutti flavor

QUESTIONS?

call **1-888-423-0139**

PRINCIPAL DISPLAY PANEL

Topcare_®

NDC 36800-527-69

MELT-IN-YOUR-MOUTH TABLETS. NO WATER NEEDED

Original Prescription Strength

24 HOUR NON-DROWSY*

Allergy Relief

LORATADINE ORALLY DISINTEGRATING TABLETS, USP 10 mg

ANTIHISTAMINE

Indoor & Outdoor Allergies

For Adults and Children six years and older!

For 24 Hour Relief of:

- Sneezing
- Runny Nose
- Itchy, Watery Eyes
- Itching Throat or Nose


 $\textbf{COMPARE TO CLARITIN}^{\circledR}\textbf{REDITABS}^{\circledR}\textbf{active ingredient}^{\dag}$

10 ORALLY DISINTEGRATING TABLETS

*When taken as directed. See Drug Facts Panel.

DISTRIBUTED BY TOPCO ASSOCIATES LLC

5095192/R812



■ TAMPER EVIDENT: DO NOT USE IF BLISTER UNITS ARE TORN, BROKEN OR SHOW ANY SIGNS OF ■ Phenylketonurics: Contains Phenylalanine 0.6 mg Per Tablet.

Other information

5.41	
sak a doctor	consumers with liver or kidney disease
sak a doctor	children under 6 years of age
1 tablet daily; not more than 1 tablet in 24 hours	adults and children 6 years and over

■ place 1 tablet on tongue; tablet disintegrates, with or without water

Directions


away (1-800-222-1222). Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right If pregnant or breast-feeding, ask a health professional before use.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

When using this product do not take more than directed. Taking more than directed may cause

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a Do not use if you have ever had an allergic reaction to this product or any of its ingredients. Warnings

Drug Facts (continued)



NDC 36800-527-69

This product is not manufactured or distributed by Schering-Plough Healthcare Products, Inc. CLARITIN® and REDITABS® are registered trademarks of Schering Corporation.

Non Varnish Area

MELT-IN-YOUR-MOUTH TABLETS • NO WATER NEEDED

24 HOUR NON-DROWSY*

Original Prescription Strength

Allergy Relief

LORATADINE ORALLY DISINTEGRATING TABLETS, USP 10 mg

Indoor & Outdoor Allergies

For Adults and Children six years and older!

For 24 Hour Relief of:

Sneezing

Runny Nose

· Itchy Throat or Nose Itchy, Watery Eyes

actual size

10 ORALLY DISINTEGRATING TABLETS

When taken as directed. See Drug Facts Panel.

Keep the carton. It contains important information. See end panel for expiration date.

Questions? call 1-800-406-7984

COMPARE TO CLARITIN® REDITABS® active ingredient[†]

sodium stearyl fumarate, strawberry cream flavor, tutti-frutti flavor Inactive ingredients aspartame, croscarmellose sodium, magnesium stearate, mannitol, mint flavor,

Drug Facts (continued)



095192

OHMR812



LORATADINE

loratadine tablet, orally disintegrating

Product Information

Product TypeHUMAN OTC DRUGItem Code (Source)NDC:36800-527

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	10 mg

Inactive Ingredients			
Ingredient Name	Strength		
ASPARTAME (UNII: Z0H242BBR1)			
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MANNITOL (UNII: 3OWL53L36A)			
SODIUM STEARYL FUMARATE (UNII: 7CV7WJK4UI)			

Product Characteristics			
Color	white (White to Off White)	Score	no score
Shape	ROUND (flat faced beveled edge)	Size	10 mm
Flavor	STRAWBERRY, TUTTI FRUTTI, MINT	Imprint Code	RC17
Contains			

F	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:36800-527-69	10 in 1 BLISTER PACK		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA077153	08/31/2007	

Labeler - Topcare Associates LLC (006935977)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Ohm Laboratories Inc.		184769029	manufacture(36800-527)

Revised: 8/2012 Topcare Associates LLC