

**TYLENOL COLD PLUS FLU SEVERE DAY AND TYLENOL COLD PLUS FLU PLUS COUGH NIGHT 2-PACK- acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and guaifenesin, phenylephrine hydrochloride  
Kenvue Brands LLC**

-----  
**Tylenol® Cold Plus Flu Severe Day and Tylenol Cold Plus Flu Plus Cough Night 2-pack**

**Tylenol® Cold plus Flu plus Cough Night**

**Drug Facts**

**Active ingredients (in each 15 mL) Purpose**

Acetaminophen 325 mg Pain reliever/ fever reducer  
Dextromethorphan HBr 10 mg Cough suppressant  
Doxylamine Succinate 6.25 mg Antihistamine  
Phenylephrine HCl 5 mg Nasal decongestant

<b>Active ingredients (in each 15 mL)</b>	<b>Purpose</b>
Acetaminophen 325 mg	Pain reliever/ fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - runny nose and sneezing
  - cough
  - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks

- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see overdose warning)**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> <li>▪ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li> <li>▪ do not take more than 150 mL in 24 hours, unless directed by a doctor</li> </ul>
children under 12 years	ask a doctor

**Other information**

- each 15 mL contains: **sodium 5 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with TYLENOL COLD+FLU+COUGH” or foil inner seal imprinted with “SAFETY SEAL® is broken or missing.**

**Inactive ingredients**

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors,

glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

## Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

## Tylenol® Cold plus Flu Severe

### Drug Facts

#### **Active ingredients (in each 30 mL) Purpose**

Acetaminophen 650 mg Pain reliever/ fever reducer

Dextromethorphan HBr 20 mg Cough suppressant

Guaifenesin 400 mg Expectorant

Phenylephrine HCl 10 mg Nasal decongestant

<b>Active ingredients (in each 30 mL)</b>	<b>Purpose</b>
Acetaminophen 650 mg	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

### Uses

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - sore throat
  - nasal congestion
  - cough
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
- temporarily reduces fever

### Warnings

#### Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Sore throat warning**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

### **When using this product do not exceed recommended dose**

### **Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

## Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## Directions

- **do not take more than directed (see overdose warning)**
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"><li>▪ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li><li>▪ do not take more than 150 mL in 24 hours, unless directed by a doctor</li></ul>
children under 12 years	ask a doctor

## Other information

- each 30 mL contains: **sodium 10 mg**
- store between 20-25°C (68-77°F).
- **do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing.**

## Inactive ingredients

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

## Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

## PRINCIPAL DISPLAY PANEL

NDC 50580-526-01

DAYTIME

NIGHTTIME



## PRINCIPAL DISPLAY PANEL

NDC 50580-369-01

TYLENOL<sup>®</sup>  
FOR ADULTS  
COLD + FLU + COUGH

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,  
Pain Reliever-Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE

NIGHT  
WILD BERRY  
BURST

Alcohol 0.7%  
8 fl oz (240 mL)

30040371

NDC 50580-369-01

**TYLENOL**<sup>®</sup>

FOR ADULTS

**COLD + FLU + COUGH****Acetaminophen**, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,  
Pain Reliever—Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



NIGHT

WILD BERRY  
BURST

Alcohol 0.7%

8 fl oz (240 mL)

OPEN TO READ DRUG FACTS  
(Warnings, Directions...)

PEEL

30040372 Do not use if neck band imprinted with  
"TYLENOL COLD+FLU+ COUGH" or foil  
inner seal imprinted with "SAFETY  
SEAL<sup>®</sup>" is broken or missing.

**DO NOT USE WITH OTHER MEDICINES  
CONTAINING ACETAMINOPHEN**



3 0045-0562-081

**Drug Facts****Active ingredients Purpose  
(in each 15 mL)**

Acetaminophen 325 mg ..... Pain reliever/  
fever reducer  
Dextromethorphan HBr 10 mg... Cough suppressant  
Doxylamine Succinate 6.25 mg..... Antihistamine  
Phenylephrine HCl 5 mg..... Nasal decongestant

**Uses**

- temporarily relieves these common cold/flu symptoms:
  - minor aches and pains
  - headache
  - nasal congestion
  - runny nose and sneezing
  - cough
  - sinus congestion and pressure
- helps clear nasal passages

**Drug Facts (continued)**

- relieves cough to help you sleep
- temporarily reduces fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

**Drug Facts (continued)**

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional

**Drug Facts (continued)**

conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- If you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as

**Drug Facts (continued)**

- emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- do not exceed recommended dose
- excitability may occur, especially in children
- marked drowsiness may occur

**Drug Facts (continued)**

- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
  - pain, nasal congestion or cough gets worse or lasts more than 7 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts
- These could be signs of a serious condition.

Drug Facts (continued)

Drug Facts (continued)

Drug Facts (continued)

**TYLENOL**<sup>®</sup>

If pregnant or breast-feeding, ask a health professional before use.  
**Keep out of reach of children.**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

#### Directions

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over

- take 30 mL in the dosing cup provided every 4 hours while symptoms last
- do not take more than 150 mL in 24 hours, unless directed by a doctor

children under 12 years

ask a doctor

#### Other information

- each 15 mL contains: sodium 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.
- do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®"

is broken or missing

#### Inactive ingredients

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

#### Questions or comments?

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

## PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

## COLD + FLU + COUGH NIGHT

How can we help?

1-877-895-3665

SNAP THIS CODE OR TEXT TYCOLD7 TO 87715

MSG & DATA RATES MAY APPLY. TEXT HELP FOR HELP. TEXT STOP FOR STOP.



Acetaminophen (325 mg per 15 mL)  
Dextromethorphan HBr (10 mg per 15 mL)  
Doxylamine succinate (6.25 mg per 15 mL)  
Phenylephrine HCl (5 mg per 15 mL)

8 fl oz (240 mL)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
J&JCI 2017 30040372

## PRINCIPAL DISPLAY PANEL

NDC 50580-403-08

30049843

Warming Honey Lemon Flavor

TYLENOL®

FOR ADULTS

COLD + FLU SEVERE

Acetaminophen, Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin

Pain Reliever-Fever Reducer, Cough Suppressant, Nasal Decongestant, Expectorant

Alcohol Free

DAY

NON-DROWSY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION

8 fl oz (240 mL)



# TYLENOL

## COLD + FLU SEVERE

**Acetaminophen**

Dextromethorphan HBr, Phenylephrine HCl, Guaifenesin  
Pain Reliever–Fever Reducer, Cough Suppressant,  
Nasal Decongestant, Expectorant

Alcohol Free

 **DAY**  
NON-DROWSY

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- MUCUS + CHEST CONGESTION



8 fl oz (240 mL)

OPEN TO READ DRUG FACTS  
(Warnings, Directions...)


PEEL

EXP./LOT:

30049845

Do not use if neck band imprinted with  
"WARMING SENSATION" or foil  
inner seal imprinted with "TYLENOL"  
is broken or missing.

**DO NOT USE WITH OTHER MEDICINES  
CONTAINING ACETAMINOPHEN**



### Drug Facts

**Active ingredients Purpose**  
(in each 30 mL)

Acetaminophen 650 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 20 mg.....	Cough suppressant
Guaifenesin 400 mg.....	Expectorant
Phenylephrine HCl 10 mg.....	Nasal decongestant

### Drug Facts (continued)

- temporarily reduces fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing

PAGE 2

PAGE 3

## Uses

- temporarily relieves the following cold/flu symptoms:
  - minor aches and pains
  - headache
  - nasal congestion
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

## acetaminophen

- 3 or more alcoholic drinks every day while using this product
- Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
  - blisters
  - rash

## Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression,

## Drug Facts (continued)

psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

## Drug Facts (continued)

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dose**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days

## Drug Facts (continued)

- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts** (continued)

**Directions**

- do not take more than directed (see overdose warning)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> <li>■ take 30 mL in the dosing cup provided every 4 hours while symptoms last</li> <li>■ do not take more than 150 mL in 24 hours, unless directed by a doctor</li> </ul>
children under 12 years	ask a doctor

**Drug Facts** (continued)

**Other information**

- each 30 mL contains: sodium 10 mg
- store between 20-25°C (68-77°F).
- do not use if neck band imprinted with "WARMING SENSATION" or foil inner seal imprinted with "TYLENOL" is broken or missing

**Inactive ingredients**

anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, FD&C yellow no. 6, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

**Drug Facts** (continued)

**Questions or comments?**

call 1-877-895-3665 (toll-free) or 215-273-8755 (collect)

**PARENTS:**

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

**TYLENOL**  
COLD + FLU SEVERE

How can we help?  
☎ 1-877-895-3665

- Acetaminophen (650 mg per 30 mL)
- Dextromethorphan HBr (20 mg per 30 mL)
- Guaifenesin (400 mg per 30 mL)
- Phenylephrine HCl (10 mg per 30 mL)

8 fl oz (240 mL)

Distributed by:  
**JOHNSON & JOHNSON CONSUMER INC.**  
McNeil Consumer Healthcare Division  
Fort Washington, PA 19034 USA  
© J&JCI 2021

30049845

**TYLENOL COLD PLUS FLU SEVERE DAY AND TYLENOL COLD PLUS FLU PLUS COUGH NIGHT 2-PACK**

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and guaifenesin, phenylephrine hydrochloride kit

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:50580-526
---------------------	----------------	---------------------------	---------------

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-526-01	1 in 1 PACKAGE	10/28/2019	

### Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	240 mL
Part 2	1 BOTTLE, PLASTIC	240 mL

### Part 1 of 2

#### TYLENOL COLD PLUS FLU PLUS COUGH NIGHT

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, and phenylephrine hydrochloride solution

### Product Information

<b>Item Code (Source)</b>	NDC:50580-369
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ACETIC ACID</b> (UNII: Q40Q9N063P)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	

<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

### Product Characteristics

<b>Color</b>	purple	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY (Berry Mint)	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-369-01	240 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/17/2019	

## Part 2 of 2

### TYLENOL COLD PLUS FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride solution

### Product Information

<b>Item Code (Source)</b>	NDC:50580-403
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	

## Product Characteristics

<b>Color</b>	brown, yellow (Amber colored)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	HONEY, LEMON (honey lemon flavored)	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-403-08	240 mL in 1 BOTTLE, PLASTIC; Type 1: Convenience Kit of Co-Package		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/05/2010	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	10/28/2019	

**Labeler** - Kenvue Brands LLC (118772437)