THERAFLU FLU RELIEF MAX STRENGTH NIGHTTIME- acetaminophen, chlorpheniramine maleate, dextromethorphan hbr syrup Haleon US Holdings LLC

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 1000 mg Chlorpheniramine maleate 4 mg Dextromethorphan HBr 30 mg

Purposes

Pain reliever/Fever reducer

Antihistamine

Cough suppressant

Uses

- temporarily relieves these symptoms due to a common cold:
 - headache
 - minor aches and pains
 - cough due to minor throat and bronchial irritation
 - minor sore throat pain
 - runny nose
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed
- measure the dose correctly using the enclosed dosing cup
- adults and children 12 years of age and over: take every 6 hours in dosing cup provided, while symptoms persist
- do not take more than 3 doses (90 mL) in 24 hours unless directed by a doctor
- children under 12 years of age: do not use

Age	Dose
adults and children 12 years of age and	30 mL
over	
children under 12 years of age	do not use

Other information

- each 30 mL contains: potassium 5 mg, sodium 34 mg
- store at controlled room temperature 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, carboxymethylcellulose sodium, glycerin, nautral and artificial flavors, natural grade A honey, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium gluconate, sucralose, xanthan gum, zinc gluconate

Questions or comments?

1-855-328-5259

Additional Information

DO NOT TAKE MORE THAN 3 DOSES IN TOTAL IN ANY 24 HOUR PERIOD. USE AS DIRECTED

PEEL BACK HERE

DO NOT USE IF NECKBAND PRINTED WITH "SEALED FOR SAFETY" IS TORN OR MISSING.

*Maximum Strength per 6 hour dose. PARENTS: Learn about teen medicine abuse www.StopMedicineAbuse.org

Principal Display Panel THERAFLU FLU RELIEF **MAX STRENGTH*** NIGHTTIME Acetaminophen Pain Reliever/Fever Reducer **Chlorpheniramine Maleate** Antihistamine **Dextromethorphan HBr Cough Suppressant HELPS YOU REST**** Powerful fever fightingformula that relieves: / Body ache / Headache / Sore throat pain / Cough

/ Runny nose

Honey & Elderberry Flavor

8.3 FL OZ (245.5 mL)

HALEON

FLU RELIEF MAX STRENGTH*

HERAFLU

SELP.

NIGHTTIME

Acetaminophen Pain Reliever/Fever Reducer

Chlorpheniramine Maleate Antihistamine Dextromethorphan HBr Cough Suppressant

Powerful *fever fighting* formula that relieves:

Body ache / Headache Sore throat pain Cough / Runny nose

Honey & Elderberry Flavor

8.3 FL 0Z (245.5 mL)

THERAFLU FLU RELIEF MAX STRENGTH NIGHTTIME

acetaminophen, chlorpheniramine maleate, dextromethorphan hbr syrup

Product Information					
Product Type	HUMAN OTC DRUG	ltem Code (Source)	NDC:006	57-8205
Route of Administration	ORAL				
Active Ingredient/Active Moiety					
Ingred	lient Name		Basis of Stre	ngth	Strength
		יחט ודוטטנאניו			1000 mg

	SOZOALITAD) (ACETAMINOLUEIA - OIAII'SOZOALITA		in 30 mL
CHLORPHENIRAMINE MA - UNII:3U6IO1965U)	ALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMIN	E CHLORPHENIRAMINE MALEATE	4 mg in 30 mL
DEXTROMETHORPHAN I (DEXTROMETHORPHAN - U	HYDROBROMIDE (UNII: 9D2RTI9KYH) NII:7355X3ROTS)	DEXTROMETHORPHA HYDROBROMIDE	N 30 mg in 30 mL
Inactive Ingredien			
	Ingredient Name	211)	Streng
	JLOSE SODIUM, UNSPECIFIED (UNII: K679OBS	311)	
GLYCERIN (UNII: PDC6A3)			
HONEY (UNII: Y9H1V576F	n) ERYL ETHER (UNII: B2138XJ83G)		
PROPYLENE GLYCOL (UI	-		
WATER (UNII: 059QF0K00			
SODIUM BENZOATE (UN			
	PECIFIED FORM (UNII: 1Q73Q2JULR)		
SODIUM GLUCONATE (L	-		
SUCRALOSE (UNII: 96K6L			
XANTHAN GUM (UNII: TT			
ZINC GLUCONATE (UNII:	U6WSN5SQ1Z)		
Other Ingredients			
-	Ingredient Nan		Oursetit
Ingredient Kind	_		Quantit
May contain	ANHYDROUS CITRIC ACID (UNII: XF417D3	PSL)	
Product Character	ristics		
Color	S	core	
Shape	S	ize	
Flavor	BERRY (Elderberry)	nprint Code	
Contains			
Packaging			
		Marketing Start	

# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:0067- 8205-01	245.5 mL in 1 BOTTLE; Type 1: Convenience Kit of Co-Package	06/27/2022	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/27/2022	
OTC Monograph Drug	MOTZ	00/27/2022	

Revised: 1/2025