

OPTION ONE- hydrogen peroxide solution IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

OPTION ONE SANITIZING TEAT DIP A HIGHLY EFFECTIVE PRE-MILKING AND POST-MILKING TEAT DIP WITH GLYCERIN

ACTIVE INGREDIENT

Hydrogen Peroxide.....1.00%

INERT INGREDIENTS

Emollients(skin conditioners).....10.00%

Other Inerts.....89.00%

(Includes Buffering Agents, Wetting Agents and Surfactants)

Total.....100%

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes dirty and/or noticeably loses color, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment
4. Wash Teat Dip cups after each milking.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. REMOVE VISIBLE SOILS FROM THE TEATS.
2. OBSERVE FOREMILK BY STRIPPING TWO OR THREE STREAMS OF MILK INTO A STRIP CUP.
3. DIP OR SPRAY THE COW'S TEATS FULL LENGTH WITH OPTION ONE TEAT DIP.
- 4 WAIT APPROXIMATELY 30 SECONDS.
5. REMOVE ALL TEAT DIP BY THOROUGHLY DRYING THE TEATS AND TEAT ORIFICES AND UDDER (IF SPRAYING), WITH A CLEAN, SINGLE SERVICE PAPER TOWEL(S).
6. ATTACH MILKING UNIT

B. POST-MILKING (POST DIPPING)

Immediately after milking dip or spray all teats full length with OPTION ONE Teat Dip Ensure good coverage of all sides of the teats and teat orifices.

WARNING

DO NOT DILUTE THIS PRODUCT

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER

CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.
PROTECT THIS PRODUCT FROM FREEZING.

CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.

FIRST AID PROCEDURES

EYE CONTACT: AVOID CONTACT WITH EYES. IF CONTACT OCCURS, FLUSH EYE WITH LARGE QUANTITIES OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUE. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE. IF IRRITATION DEVELOPS, CONSULT A PHYSICIAN.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DRINK TWO GLASSES OF WATER AND CONSULT A PHYSICIAN.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION

FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

OPTION ONE

NDC 29983-0055-4



OPTION ONE

SANITIZING TEAT DIP

A Highly Effective Pre-Milking and Post-Milking Teat Dip With Glycerin

ACTIVE INGREDIENT:

Hydrogen Peroxide 1.00%

INERT INGREDIENTS:

Emollients (Skin Conditioners) 10.00%

Other Inerts: 89.00%

(Includes Buffering Agents, Surfactants and Wetting Agents)

TOTAL: 100.00%

GENERAL RECOMMENDATIONS

1. Use teat dip applicators made of plastic or other non-corrosive materials. Do not use cups made of aluminum.
2. When solution in the teat dip cup becomes visibly dirty, discard remaining contents and refill cup with fresh teat dip.
3. Do not use this product for cleaning or sanitizing equipment.
4. Wash teat dip cups after each milking.

CAUTION

KEEP OUT OF REACH OF CHILDREN. AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY. PROTECT EYES AND MUCOUS MEMBRANES FROM CONTACT WITH THIS PRODUCT.


FIRST AID PROCEDURES

EYE CONTACT: AVOID CONTACT WITH EYES. IF CONTACT OCCURS, FLUSH EYE WITH LARGE QUANTITIES OF FRESH, TEPID WATER FOR 15 MINUTES. HOLD EYELID OPEN TO ENSURE COMPLETE IRRIGATION OF EYE AND LID TISSUE. TILT HEAD TO THE SIDE AND IRRIGATE THE EYE FROM THE BRIDGE OF THE NOSE TO THE OUTSIDE OF THE FACE. IF IRRITATION DEVELOPS, CONSULT A PHYSICIAN.

SKIN CONTACT: WASH ANY SENSITIVE AREAS.

INGESTION: DRINK TWO GLASSES OF WATER AND CONSULT A PHYSICIAN.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300



RECOMMENDED PROCEDURES FOR TEAT SANITATION

A. PRE-MILKING (PRE-DIPPING)

1. Remove visible soils from the teats.
2. Observe foremilk by stripping two or three streams of milk into a strip cup.
3. Dip or spray the cow's teats full length with OPTION ONE Teat Dip.
4. Wait approximately 30 seconds.
5. Remove all teat dip by thoroughly drying the teats, teat orifices and udder (if spraying), with a clean, single service paper towel(s) to avoid contamination of milk.
6. Attach milking unit.

B. POST-MILKING (POST-DIPPING)

1. Immediately after milking, dip or spray all teats full length with OPTION ONE Teat Dip. Ensure good coverage of all sides of the teats and teat orifices.

WARNING

DO NOT MIX OPTION ONE TEAT DIP WITH ANY OTHER PRODUCT.

DO NOT DILUTE THIS PRODUCT.

DO NOT RETURN TEAT DIP LEFT IN THE TEAT DIPPER TO STORAGE CONTAINER.

ENSURE THAT DIPPED TEATS ARE DRY BEFORE EXPOSING THE ANIMALS TO WEATHER CONDITIONS WHICH MAY CAUSE DAMAGE TO THE TEATS.

PROTECT THIS PRODUCT FROM FREEZING.

A PART OF THE COMPLETE IBA SANITATION PROGRAM

EXP. DATE:

LOT NO.:

NET CONTENTS: 5 GALLONS

IBA STOCK # 160225

308E

MANUFACTURED EXCLUSIVELY FOR IBA INC., SANITATION DIVISION, 27 PROVIDENCE ROAD, MILLBURY, MA 01527

OPTION ONE

hydrogen peroxide solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:29983-0055
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	10.0 g in 1 L

Product Characteristics

Color	orange	Score	
-------	--------	-------	--

Shape		Size		
Flavor		Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:29983-0055-4	18.9 L in 1 DRUM		
2	NDC:29983-0055-6	56.7 L in 1 DRUM		
3	NDC:29983-0055-7	113.4 L in 1 DRUM		
4	NDC:29983-0055-8	207.9 L in 1 DRUM		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
unapproved drug other			05/14/2010	

Labeler - IBA (019494160)