

ASPIRIN- aspirin tablet, film coated
Amerisource Bergen

Good Neighbor Pharmacy 44-157

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

temporarily relieves:

- headache
- pain and fever of colds
- toothache
- menstrual pain
- muscle pain
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction, which may include:

- shock
- facial swelling
- hives
- asthma (wheezing)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have asthma
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
- children under 12 years: do not use unless directed by a doctor

Other information

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments?

1-800-426-9391 8:30 AM - 4:00 PM ET Monday - Friday

Principal Display Panel

**Good
Neighbor
Pharmacy®**

NDC 46122-691-78

**Compare to
Bayer® Genuine
Aspirin active ingredient****

REGULAR STRENGTH

Aspirin
325 mg (NSAID)

Pain Reliever / Fever Reducer

Actual
Size

100
Tablets
325 mg Each

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

Good Neighbor Pharmacy

**Satisfaction
Guaranteed**
Brand Products

Distributed By
AmerisourceBergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

**This product is not manufactured or distributed by Bayer AG, owner of the registered trademark Bayer® Genuine Aspirin.

50844 ORG012215712



Compare to Bayer® Genuine Aspirin active ingredient**
NDC 46122-691-78



Distributed By
AmersourceBergen
1 West First Avenue
Conshohocken, PA 19428
Questions or Concerns?
www.mygnp.com

REGULAR STRENGTH

Aspirin

325 mg (NSAID)

Pain Reliever / Fever Reducer

Actual Size



100 Tablets
325 mg Each

ABC#: 10282784 Rev: 10/23



3 46122 69178 9

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Drug Facts	Active ingredient (in each tablet) Aspirin 325 mg (NSAID)* Pain reliever/fever reducer *nonsteroidal anti-inflammatory drug	Purpose	No Print/No Varnish Lot & EXP
Uses	temporarily relieves: ■ headache ■ muscle pain ■ toothache ■ menstrual pain ■ pain and fever of colds ■ minor pain of arthritis	Warnings	Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness. Allergy alert: Aspirin may cause a severe allergic reaction, which may include: ■ facial swelling ■ shock ■ hives ■ asthma (wheezing)
Do not use	■ if you are allergic to aspirin or any other pain reliever/fever reducer ■ if you have ever had an allergic reaction to this product or any of its ingredients	Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ take more or for a longer time than directed ■ have 3 or more alcoholic drinks every day while using this product ■ have had stomach ulcers or bleeding problems ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ are age 60 or older	

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

■ take a blood thinning (anticoagulant) or steroid drug

Do not use ■ if you are allergic to aspirin or any other pain reliever/fever reducer
■ if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if ■ you are taking a diuretic
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease ■ you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ vomit blood ■ have bloody or black stools
■ have stomach pain that does not get better ■ feel faint
■ ringing in the ears or a loss of hearing occurs
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
■ drink a full glass of water with each dose
■ adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours not to exceed 12 tablets in 24 hours unless directed by a doctor
■ children under 12 years: do not use unless directed by a doctor

Other information
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) ■ use by expiration date on package

Inactive ingredients corn starch, hypromellose, polyethylene glycol, propylene glycol

Questions or comments? 1-800-426-9391
8:30 AM - 4:00 PM ET Monday - Friday

STOP PEELING

Adhesive Only

GNP 44-157

ASPIRIN			
aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-691
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	

ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
---	---------	--------

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

Product Characteristics

Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-691-78	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/10/2021	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	03/10/2021	

Labeler - Amerisource Bergen (007914906)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-691)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(46122-691)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(46122-691)

Establishment

Name	Address	ID/FEI	Business Operations

LNK International, Inc.		868734088	manufacture(46122-691)
-------------------------	--	-----------	------------------------

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(46122-691)

Revised: 9/2025

Amerisource Bergen