

**ANTI-DIARRHEAL- loperamide hcl tablet
DOLGENCORP, LLC**

Dollar General 44-375

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are

taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- **each caplet contains:** calcium 15 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal Display Panel

DG® | health

Compare to active

**ingredient of
Imodium® A-D***

Anti-Diarrheal

Loperamide HCl Tablets, 2 mg
Anti-Diarrheal

- Controls the symptoms
of diarrhea

2
mg each

24 Caplets

Actual Caplet Size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**

1. To open, tear
at perforations and
remove a section.

2. Use ✂ to cut
through plastic
for caplet.

*This product is not manufactured or distributed by Kenvue Inc.,
owner of the registered trademark Imodium® A-D.
50844 REV0325A37508

DISTRIBUTED BY
OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

100%
Satisfaction
Guaranteed!
(888)309-9030

Drug Facts (continued)

Heart alert: Taking more than directed can cause serious heart problems or death.

Do not use if you have bloody or black stool.

Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease ■ a history of abnormal heart rhythm

Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.

When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bloating. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea. ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information ■ each caplet contains calcium 15 mg

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BUSTERS IS TORN OR BROKEN ■ score between 20'-25' C (68'-77' F) ■ see end flap for expiration date and lot number

Inactive ingredients corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions or comments? Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Drug Facts **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Active ingredient (in each caplet) Loperamide HCl 2 mg

Purpose Anti-diarrheal

Use controls symptoms of diarrhea, including Traveler's Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

B-0315-375-08
REV0325A37508



Compare to active ingredient of Imodium® A-D*

Anti-Diarrheal

Loperamide HCl Tablets, 2 mg
Anti-Diarrheal

- Controls the symptoms of diarrhea

1. To open, tear at perforations and remove a section.
2. Use 1/2 to cut through plastic for caplet.




DO NOT USE IF PACKAGE IS TORN, BROKEN OR SIGNS OF TAMPERING

Dispense in original container. Do not use if package is torn, broken or shows signs of tampering.

*This product is not manufactured or distributed by Kenvue Inc., owner of the registered trademark Imodium® A-D.
50844 REV0325A37508

DISTRIBUTED BY
OLD EAST MAIN CO.
100 MISSION RIDGE
GOODLETTSVILLE, TN 37072

10
Satisfy
Guara
(888) 3-



ANTI-DIARRHEAL

loperamide hcl tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55910-375
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X90C3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
D&C YELLOW NO. 10 ALUMINUM LAKE (UNII: CQ3XH3DET6)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	

Product Characteristics

Color	green	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	44;375
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-375-02	2 in 1 CARTON	01/12/2024	
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:55910-375-08	4 in 1 CARTON	01/12/2024	
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	01/12/2024	

Labeler - DOLGENCORP, LLC (068331990)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55910-375) , pack(55910-375)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55910-375)

Revised: 8/2025

DOLGENCORP, LLC