

**EQUATE DAYTIME NIGHTTIME COLD AND FLU- acetaminophen,
dextromethorphan hydrobromide, doxylamine succinate, guaifenesin,
phenylephrine hydrochloride
WALMART INC.**

Wal-Mart Daytime Nighttime Cold and Flu Drug Facts

Active ingredients (in each 15 mL) {Nighttime}

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine Succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- minor aches and pains
- headache
- fever
- sore throat
- runny nose and sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours

- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacists before taking this product.
- to make a child sleepy
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- glaucoma
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur

- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-888-287-1915

Active ingredients (in each 15 mL) {Daytime}

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Guaifenesin 200 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Expectorant

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- minor aches and pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotion conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present

- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments?

1-888-287-1915

Package/Label Principal Display Panel

VALUE PACK

equate™

Compare to Vicks® DayQuil® Severe Honey Flavor active ingredients

DAYTIME

SEVERE

HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

Daytime Relief

- Acetaminophen – Minor aches & pains, fever
- Phenylephrine HCl – Nasal congestion & sinus pressure
- Dextromethorphan HBr – Cough
- Guaifenesin – Chest congestion

Compare to Vicks® NyQuil® Severe Honey Flavor active ingredients

NIGHTTIME

SEVERE

HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

Nighttime Relief

- Acetaminophen – Minor aches & pains, fever
- Phenylephrine HCl – Nasal congestion & sinus pressure
- Dextromethorphan HBr – Cough
- Doxylamine Succinate – Runny nose & sneezing

2 – 12 FL OZ (355mL) BOTTLES/24 FL OZ (1 PT 8 FL OZ) 710mL TOTAL

equate™



NON-DROWSY

SEVERE HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant

equate™



SEVERE

HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

VALUE PACK

equate™

NDC 79903-348-24



equate™



SEVERE

HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Satisfaction Guaranteed

For more information call 1-888-287-1915 or visit Walmart.com/equate

DISTRIBUTED BY: Walmart Inc., Bentonville, AR 72716

*These products are not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil® and DayQuil®.

2- 12 FL OZ (355mL) BOTTLES / 24 FL OZ (1 PT 8 FL OZ) 710mL TOTAL



BOX BOTTLE CUP



44X02 2E C1



Nighttime Severe Honey Flavor Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

- Uses** temporarily relieves common cold/flu symptoms:
- nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - headache
 - fever
 - sore throat
 - cough due to minor throat and bronchial irritation
 - runny nose and sneezing
 - reduces swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage
 - cough to help you sleep

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if
- adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- to make a child sleepy
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- glaucoma
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
 - pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-888-287-1915

Daytime Severe Honey Flavor Cold & Flu

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

- Uses** temporarily relieves common cold/flu symptoms:
- nasal congestion
 - sinus congestion and pressure
 - cough due to minor throat and bronchial irritation
 - minor aches and pains
 - headache
 - fever
 - sore throat
 - reduces swelling of nasal passages
 - temporarily restores freer breathing through the nose
 - promotes nasal and/or sinus drainage
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if
- adult takes more than 4,000 mg of acetaminophen in 24 hours
 - child takes more than 5 doses in 24 hours
 - taken with other drugs containing acetaminophen
 - adult has 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
- If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
 - pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present
 - new symptoms occur
 - cough comes back or occurs with rash or headache that lasts.
- These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed - see Overdose warning
 - only use the dose cup provided
 - do not exceed 4 doses per 24 hrs
- | | |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs | 15 mL every 4 hrs |
| children 4 to under 6 yrs | ask a doctor |
| children under 4 yrs | do not use |

Other information

- each 15 mL contains: sodium 15 mg
- store at 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions or comments? 1-888-287-1915

equate™

DAYTIME

NON-DROWSY

SEVERE

HONEY FLAVOR

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Expectorant, Nasal Decongestant



EQUATE DAYTIME NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, guaifenesen, phenylephrine hydrochloride kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-348
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-348-24	1 in 1 CARTON; Type 0: Not a Combination Product	07/07/2025	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	355 mL
Part 2	1 BOTTLE	355 mL

Part 1 of 2

EQUATE NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:79903-122
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-122-12	1 in 1 CARTON		
1		355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2023	

Part 2 of 2

EQUATE COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, guaifenesin, phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:79903-123
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0K00R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	YELLOW	Score	
Shape		Size	
Flavor	HONEY, MENTHOL	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-123-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing	Application Number or Monograph	Marketing Start	Marketing End
-----------	---------------------------------	-----------------	---------------

Category	Citation	Date	Date
OTC Monograph Drug	M012	06/30/2023	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/07/2025	

Labeler - WALMART INC. (051957769)

Revised: 4/2026

WALMART INC.