

CHILDRENS TYLENOL- acetaminophen powder
Kenvue Brands LLC

Children's TYLENOL

Drug Facts

Active ingredient (in each powder)

Acetaminophen 160 mg

Purpose

Pain reliever/fever reducer

Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
 - the common cold
 - flu
 - headache
 - sore throat
 - toothache

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **this product does not contain directions or complete warnings for adult use.**
- **do not give more than directed (see overdose warning)**
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- tear packet and pour powder on your child's tongue
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (powder) *
under 48	under 6 years	do not use
48-59 lbs	6-8 years	2 powders
60-71 lbs	9-10 years	2 powders
72-95 lbs	11 years	3 powders

* or as directed by a doctor

Other information

- store between 20-25°C (68-77°F)

- **do not use if packet is torn or damaged**

Inactive ingredients

citric acid, ethylcellulose, flavor, magnesium stearate, maltodextrin, sodium bicarbonate, sucralose, xylitol

Questions or comments?

call **1-800-458-1635** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-555-04

Children's
TYLENOL®

Acetaminophen
Pain Reliever-Fever Reducer

Ibuprofen Free
Aspirin Free

Pain+Fever
Ages 6-11

DISSOLVE PACKS

No Water Needed
Dissolves in Seconds
Dye - Free

30 Packets*
160 mg
each packet

Wild Berry
Flavor

*Packets of Powder



NDC 50580-555-04

Children's TYLENOL®

Acetaminophen
Pain Reliever-Fever Reducer
Pain + Fever

Ibuprofen Free
Aspirin Free
Ages 6-11

DISSOLVE PACKS

- ☑ No Water Needed
- ☑ Dissolves in Seconds
- ✓ Dye - Free

30 Packets*
160 mg
each packet



**Wild Berry
Flavor**
*Packets of Powder

Children's
TYLENOL®

DISSOLVE PACKS

- ☑ No Water Needed
- ☑ Dissolves In Seconds
- ✓ Dye - Free



How can we help?
1-800-458-1635



Important: Read all product information before using. Keep this box for Important Information.

Drug Facts

Active ingredient (in each powder) **Purpose**
Acetaminophen 160 mg Pain reliever/fever reducer

Uses temporarily: ■ reduces fever
■ relieves minor aches and pains due to: ■ the common cold
■ flu ■ headache ■ sore throat ■ toothache

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
■ more than 5 doses in 24 hours, which is the maximum daily amount
■ with other drugs containing acetaminophen

Drug Facts (continued)

■ fever gets worse or lasts more than 3 days
■ new symptoms occur ■ redness or swelling is present
These could be signs of a serious condition

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ this product does not contain directions or complete warnings for adult use.
■ do not give more than directed (see overdose warning)

Children's
TYLENOL®

DISSOLVE PACKS

Important: See Drug Facts for Complete Product Information

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days

- do not give more than directed (see overdose warning)
- find right dose on chart below. If possible, use weight to dose; otherwise, use age.
- tear packet and pour powder on your child's tongue
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (powder)*
under 48	under 6 years	do not use
48-59 lbs	6-8 years	2 powders
60-71 lbs	9-10 years	2 powders
72-95 lbs	11 years	3 powders

*or as directed by a doctor

Other information ■ store between 20-25°C (68-77°F)

- do not use if packet is torn or damaged

Inactive ingredients citric acid, ethylcellulose, flavor, magnesium stearate, maltodextrin, sodium bicarbonate, sucralose, xylitol

Questions or comments?
call 1-800-458-1635 (toll-free) or 215-273-8755 (collect)

Directions

STEP 1
Fold on line and tear



STEP 2
Pour powder on your child's tongue





CDJ-11606-004-A



3 00450 41630 8

Distributed by:
**JOHNSON & JOHNSON
CONSUMER INC.**
McNeil Consumer
Healthcare Division
Fort Washington, PA
19034 USA
US D894,008
©J&JCI 2021



30049667/JNJ0BOI000002-V3

CHILDRENS TYLENOL acetaminophen powder			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50580-555
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)		ACETAMINOPHEN	160 mg
Inactive Ingredients			
Ingredient Name			Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)			
ETHYLCELLULOSE, UNSPECIFIED (UNII: 7Z8S9VYZ4B)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MALTODEXTRIN (UNII: 7CVR7L4A2D)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
SUCRALOSE (UNII: 96K6UQ3ZD4)			
XYLITOL (UNII: VCQ006KQ1E)			
Product Characteristics			

Color	white (off white)	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50580-555-04	30 in 1 CARTON	07/15/2019	
1		1 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:50580-555-05	18 in 1 CARTON	07/15/2019	
2		1 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/15/2019	

Labeler - Kenvue Brands LLC (118772437)