

**COLD AND FLU NIGHTTIME- acetaminophen, dextromethorphan hbr,
doxylamine succinate solution**

Bi-Mart

Bi-Mart 44-014

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation
 - minor aches and pains
 - sore throat
 - headache
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- liver disease
- glaucoma

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**

- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- **each 30 mL contains:** sodium 22 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients

anhydrous citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?

1-800-935-6737

Principal Display Panel

BI-MART

NDC 37835-524-02

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu*

NIGHTTIME COLD & FLU

Acetaminophen

(Pain reliever/fever reducer)

Dextromethorphan HBr

(Cough suppressant)

Doxylamine succinate

(Antihistamine)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Sneezing, Runny Nose,
Cough

Ages 12 Years
and Over

Eucalyptus Mint Flavor

12 FL OZ (355 mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

PARENTS:

Learn About Teen Medicine Abuse

www.StopMedicineAbuse.org

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE

If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® Cold & Flu.
50844 ORG032301402

Distributed by: BI-MART
Eugene, OR 97402
B-014 ORG
G9524-012-11-3

BI-MART

NDC 37835-524-02

Compare to the active ingredients in Vicks® NyQuil® Cold & Flu*

NIGHTTIME

COLD & FLU

Acetaminophen
(Pain reliever/fever reducer)
Dextromethorphan HBr
(Cough suppressant)
Doxylamine succinate
(Antihistamine)

Headache, Fever, Sore Throat,
Minor Aches & Pains,
Sneezing, Runny Nose,
Cough

Ages 12 Years
and Over

Eucalyptus Mint Flavor

12 FL OZ (355 mL)

F-014
ORG



PEEL BACK FOR COMPLETE DRUG FACTS

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED NECK WRAP IS BROKEN OR MISSING

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 30 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine

Uses ■ temporarily relieves common cold and flu symptoms:
■ runny nose and sneezing
■ cough due to minor throat and bronchial irritation
■ minor aches and pains ■ sore throat ■ headache
■ temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the registered trademark Vicks® NyQuil® Cold & Flu. 50844 ORG032301402

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by: BI-MART
Eugene, OR 97402
B-014 ORG
G9524-012-11-3



no print / no varnish area
lot no. & exp. date

Drug Facts (continued)

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- liver disease ■ glaucoma

Ask a doctor or pharmacist before use if you are

Drug Facts (continued)

- fever gets worse or lasts more than 3 days
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- do not exceed 4 doses per 24 hours
- mL = milliliter
- only use the dose cup provided
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years: do not use

Other information

- each 30 mL contains: sodium 22 mg

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- excitability may occur, especially in children
- avoid alcoholic beverages
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- new symptoms occur
- redness or swelling is present
- pain or cough gets worse or lasts more than 7 days

Each of the contains 30 mg

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- use by expiration date on package

Inactive ingredients anhydrous citric acid, D&C yellow #10, FD&C green #3, FD&C yellow #6, flavors, glycerin, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate dihydrate, sodium saccharin, sucralose

Questions or comments?
 ☎ 1-800-935-6737

BI-MART UNCONDITIONAL MONEY BACK GUARANTEE
 If you are not completely satisfied with your BI-MART QUALITY PRODUCT, regardless of the reason, return the unused portion and your entire purchase price will be cheerfully refunded.

BiMart 44-014

COLD AND FLU NIGHTTIME

acetaminophen, dextromethorphan hbr, doxylamine succinate solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37835-524
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	

SUCRALOSE (UNII: 96K6UQ3ZD4)

Product Characteristics

Color	green	Score	
Shape		Size	
Flavor	MINT (Eucalyptus)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37835-524-02	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	01/19/2023	
2	NDC:37835-524-30	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/18/2018	01/19/2023

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/18/2018	

Labeler - Bi-Mart (027630078)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(37835-524) , pack(37835-524)

Revised: 7/2025

Bi-Mart