

DEXAMETHASONE SODIUM PHOSPHATE- dexamethasone sodium phosphate injection
Somerset Therapeutics, LLC


Dexamethasone Sodium Phosphate Injection, USP
For Intravenous or Intramuscular Use Only
Rx Only

DESCRIPTION

Dexamethasone Sodium Phosphate Injection, USP, is a water-soluble inorganic ester of dexamethasone which produces a rapid response even when injected intramuscularly.

Dexamethasone Sodium Phosphate, USP chemically is Pregna-1,4-diene-3,20-dione, 9-fluoro- 11,17-dihydroxy-16-methyl-21-(phosphonoxy)-, disodium salt, (11 β , 16 α).

It occurs as a white to practically white powder, is hygroscopic, is soluble in water and its solutions have a pH between 7.5 and 9.5. It has the following structural formula:



$C_{22}H_{28}FNa_2O_8P$

M.W. 516.41

Each mL of Dexamethasone Sodium Phosphate Injection, USP (**Preservative Free**) contains dexamethasone sodium phosphate, USP equivalent to 10 mg dexamethasone phosphate; 24.75 mg sodium citrate, dihydrate; and Water for Injection, q.s. pH adjusted with citric acid monohydrate or sodium hydroxide, if necessary. pH: 7.0 to 8.5.

ACTIONS:

Naturally occurring glucocorticoids (hydrocortisone), which also have salt-retaining properties, are used as replacement therapy in adrenocortical deficiency states. Their synthetic analogs are primarily used for their potent anti-inflammatory effects in disorders of many organ systems.

Glucocorticoids cause profound and varied metabolic effects. In addition, they modify the body's immune responses to diverse stimuli.

INDICATIONS AND USAGE

A. Intravenous or intramuscular administration

When oral therapy is not feasible and the strength, dosage form, and route of administration of the drug reasonably lend the preparation to the treatment of the condition, those products labeled for intravenous or intramuscular use are indicated as follows:

1. Endocrine Disorders

Primary or secondary adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; synthetic analogs may be used in conjunction with mineralocorticoids where applicable; in infancy, mineralocorticoid supplementation is of particular importance).

Acute adrenocortical insufficiency (hydrocortisone or cortisone is the drug of choice; mineralocorticoid supplementation may be necessary, particularly when synthetic analogs are used).

Preoperatively, and in the event of serious trauma or illness, in patients with known adrenal insufficiency or when adrenocortical reserve is doubtful.

Shock unresponsive to conventional therapy if adrenocortical insufficiency exists or is suspected.

Congenital adrenal hyperplasia

Nonsuppurative thyroiditis

Hypercalcemia associated with cancer

2. Rheumatic Disorders

As adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

Post-traumatic osteoarthritis

Synovitis of osteoarthritis

Rheumatoid arthritis, including juvenile rheumatoid arthritis (selected cases may require low-dose maintenance therapy).

Acute and subacute bursitis

Epicondylitis

Acute nonspecific tenosynovitis

Acute gouty arthritis

Psoriatic arthritis

Ankylosing spondylitis

3. Collagen Diseases

During an exacerbation or as maintenance therapy in selected cases of:

Systemic lupus erythematosus

Acute rheumatic carditis

4. *Dermatologic Diseases*

Pemphigus

Severe erythema multiforme (Stevens-Johnson syndrome)

Exfoliative dermatitis

Bullous dermatitis herpetiformis

Severe seborrheic dermatitis

Severe psoriasis

Mycosis fungoides

5. *Allergic States*

Control of severe or incapacitating allergic conditions intractable to adequate trials of conventional treatment in:

Bronchial asthma

Contact dermatitis

Atopic dermatitis

Serum sickness

Seasonal or perennial allergic rhinitis

Drug hypersensitivity reactions

Urticarial transfusion reactions

Acute noninfectious laryngeal edema (epinephrine is the drug of first choice).

6. *Ophthalmic Diseases*

Severe acute and chronic allergic and inflammatory processes involving the eye, such as:

Herpes zoster ophthalmicus

Iritis, iridocyclitis

Chorioretinitis

Diffuse posterior uveitis and choroiditis

Optic neuritis

Sympathetic ophthalmia

Anterior segment inflammation

Allergic conjunctivitis

Allergic corneal marginal ulcers

Keratitis

7. *Gastrointestinal Diseases*

To tide the patient over a critical period of the disease in:

Ulcerative colitis (systemic therapy)

Regional enteritis (systemic therapy)

8. *Respiratory Diseases*

Symptomatic sarcoidosis

Berylliosis

Fulminating or disseminated pulmonary tuberculosis when used concurrently with appropriate anti-tuberculous chemotherapy.

Loeffler's syndrome not manageable by other means.

Aspiration pneumonitis

9. *Hematologic Disorders*

Acquired (autoimmune) hemolytic anemia.

Idiopathic thrombocytopenic purpura in adults (IV only; IM administration is contraindicated).

Secondary thrombocytopenia in adults

Erythroblastopenia (RBC anemia)

Congenital (erythroid) hypoplastic anemia

10. *Neoplastic Diseases*

For palliative management of:

Leukemias and lymphomas in adults

Acute leukemia of childhood

11. *Edematous States*

To induce diuresis or remission of proteinuria in the nephrotic syndrome, without uremia, of the idiopathic type or that due to lupus erythematosus.

12. *Nervous System*

Acute exacerbations of multiple sclerosis

13. *Miscellaneous*

Tuberculous meningitis with subarachnoid block or impending block when used concurrently with appropriate anti-tuberculosis chemotherapy.

Trichinosis with neurologic or myocardial involvement.

Diagnostic testing of adrenocortical hyperfunction.

Cerebral edema of diverse etiologies in conjunction with adequate neurological evaluation and management.

B. Intra-articular or soft tissue administration

When the strength and dosage form of the drug lend the preparation to the treatment of the condition, those products labeled for intra-articular or soft tissue administration

are indicated as adjunctive therapy for short-term administration (to tide the patient over an acute episode or exacerbation) in:

Synovitis of osteoarthritis.

Rheumatoid arthritis.

Acute and subacute bursitis.

Acute gouty arthritis.

Epicondylitis.

Acute nonspecific tenosynovitis.

Post-traumatic osteoarthritis.

C. Intralesional administration

When the strength and dosage form of the drug lend the preparation to the treatment of the condition, those products labeled for intralesional administration are indicated for:

Keloids.

Localized hypertrophic, infiltrated, inflammatory lesions of: lichen planus, psoriatic plaques,

granuloma annulare, and lichen simplex chronicus (neurodermatitis).

Discoid lupus erythematosus.

Necrobiosis lipoidica diabetorum.

Alopecia areata.

They also may be useful in cystic tumors of an aponeurosis tendon (ganglia).

CONTRAINDICATIONS

Systemic fungal infections.

WARNINGS

Serious Neurologic Adverse Reactions with Epidural Administration

Serious neurologic events, some resulting in death, have been reported with epidural injection of corticosteroids. Specific events reported include, but are not limited to, spinal cord infarction, paraplegia, quadriplegia, cortical blindness, and stroke. These serious neurologic events have been reported with and without use of fluoroscopy. The safety and effectiveness of epidural administration of corticosteroids have not been established, and corticosteroids are not approved for this use.

In patients on corticosteroid therapy subjected to any unusual stress, increased dosage of rapidly acting corticosteroids before, during, and after the stressful situation is indicated.

Immunosuppression and Increased Risk of Infection

Corticosteroids, including dexamethasone sodium phosphate injection, suppress the

immune system and increase the risk of infection with any pathogen, including viral, bacterial, fungal, protozoan, or helminthic pathogens. Corticosteroids can:

- Reduce resistance to new infections
- Exacerbate existing infections
- Increase the risk of disseminated infections
- Increase the risk of reactivation or exacerbation of latent infections
- Mask some signs of infection

Corticosteroid-associated infections can be mild but can be severe and at times fatal. The rate of infectious complications increases with increasing corticosteroid dosages

Monitor for the development of infection and consider dexamethasone sodium phosphate injection withdrawal or dosage reduction as needed.

Tuberculosis

If dexamethasone sodium phosphate injection is used to treat a condition in patients with latent tuberculosis or tuberculin reactivity, tuberculosis may occur. Closely monitor such patients for reactivation. During prolonged therapy, patients with latent tuberculosis or tuberculin reactivity should receive chemoprophylaxis.

Varicella Zoster and Measles Viral Infections

Varicella and measles can have a serious or even fatal course in non-immune patients taking corticosteroids, including dexamethasone sodium phosphate injection. In corticosteroid-treated patients who have not had these diseases or are nonimmune, particular care should be taken to avoid exposure to varicella and measles:

- If a dexamethasone sodium phosphate injection-treated patient is exposed to varicella, prophylaxis with varicella zoster immune globulin may be indicated. If varicella develops, treatment with antiviral agents may be considered.
- If a dexamethasone sodium phosphate injection-treated patient is exposed to measles, prophylaxis with immunoglobulin may be indicated.

Hepatitis B Virus Reactivation

Hepatitis B virus reactivation can occur in patients who are hepatitis B carriers treated with immunosuppressive dosages of corticosteroids, including dexamethasone sodium phosphate injection. Reactivation can also occur infrequently in corticosteroid-treated patients who appear to have resolved hepatitis B infection.

Screen patients for hepatitis B infection before initiating immunosuppressive (e.g., prolonged) treatment with dexamethasone sodium phosphate injection. For patients who show evidence of hepatitis B infection, recommend consultation with physicians with expertise in managing hepatitis B regarding monitoring and consideration for hepatitis B antiviral therapy.

Fungal Infections

Corticosteroids, including dexamethasone sodium phosphate injection, may exacerbate systemic fungal infections; therefore, avoid dexamethasone sodium phosphate injection use in the presence of such infections unless dexamethasone sodium phosphate injection is needed to control drug reactions. For patients on chronic dexamethasone sodium phosphate injection therapy who develop systemic fungal infections, dexamethasone sodium phosphate injection withdrawal or dosage reduction is

recommended.

Amebiasis

Corticosteroids, including dexamethasone sodium phosphate injection, may activate latent amebiasis. Therefore, it is recommended that latent amebiasis or active amebiasis be ruled out before initiating dexamethasone sodium phosphate injection in patients who have spent time in the tropics or patients with unexplained diarrhea.

Strongyloides Infestation

Corticosteroids, including dexamethasone sodium phosphate injection, should be used with great care in patients with known or suspected *Strongyloides* (threadworm) infestation. In such patients, corticosteroid-induced immunosuppression may lead to *Strongyloides* hyperinfection and dissemination with widespread larval migration, often accompanied by severe enterocolitis and potentially fatal gram-negative septicemia.

Cerebral Malaria

Avoid corticosteroids, including dexamethasone sodium phosphate injection, in patients with cerebral malaria.

Kaposi's Sarcoma

Kaposi's sarcoma has been reported to occur in patients receiving corticosteroid therapy, most often for chronic conditions. Discontinuation of corticosteroids may result in clinical improvement of Kaposi's sarcoma.

Prolonged use of corticosteroids may produce posterior subcapsular cataracts, glaucoma with possible damage to the optic nerves, and may enhance the establishment of secondary ocular infections due to fungi or viruses.

Usage in Pregnancy

Since adequate human reproduction studies have not been done with corticosteroids, use of these drugs in pregnancy, nursing mothers or women of childbearing potential requires that the possible benefits of the drug be weighed against the potential hazards to the mother and embryo or fetus. Infants born of mothers who have received substantial doses of corticosteroids during pregnancy should be carefully observed for signs of hypoadrenalism.

Average and large doses of cortisone or hydrocortisone can cause elevation of blood pressure, salt and water retention, and increased excretion of potassium. These effects are less likely to occur with the synthetic derivatives except when used in large doses. Patients with a stressed myocardium should be observed carefully and the drug administered slowly since premature ventricular contractions may occur with rapid administration. Dietary salt restriction and potassium supplementation may be necessary. All corticosteroids increase calcium excretion.

While on corticosteroid therapy patients should not be vaccinated against smallpox. Other immunization procedures should not be undertaken in patients who are on corticosteroids, especially in high doses, because of possible hazards of neurological complications and lack of antibody response.

Because rare instances of anaphylactoid reactions have occurred in patients receiving parenteral

corticosteroid therapy, appropriate precautionary measures should be taken prior to administration, especially when the patient has a history of allergy to any drug.

PRECAUTIONS

Drug-induced secondary adrenocortical insufficiency may be minimized by gradual reduction of dosage. This type of relative insufficiency may persist for months after discontinuation of therapy; therefore, in any situation of stress occurring during that period, hormone therapy should be reinstated. Since mineralocorticoid secretion may be impaired, salt and/or a mineralocorticoid should be administered concurrently.

There is an enhanced effect of corticosteroids in patients with hypothyroidism and in those with cirrhosis.

Corticosteroids should be used cautiously in patients with ocular herpes simplex for fear of corneal perforation.

The lowest possible dose of corticosteroid should be used to control the condition under treatment, and when reduction in dosage is possible, the reduction must be gradual.

Psychic derangements may appear when corticosteroids are used, ranging from euphoria, insomnia, mood swings, personality changes, and severe depression to frank psychotic manifestations. Also, existing emotional instability or psychotic tendencies may be aggravated by corticosteroids.

Aspirin should be used cautiously in conjunction with corticosteroids in hypoprothrombinemia.

Steroids should be used with caution in nonspecific ulcerative colitis, if there is a probability of impending perforation, abscess, or other pyogenic infection, also in diverticulitis, fresh intestinal anastomoses, active or latent peptic ulcer, renal insufficiency, hypertension, osteoporosis, and myasthenia gravis.

Growth and development of infants and children on prolonged corticosteroid therapy should be carefully followed.

Patients who are on immunosuppressant doses of corticosteroids should be warned to avoid exposure to chickenpox or measles and, if exposed, to obtain medical advice.

Intra-articular injection of a corticosteroid may produce systemic as well as local effects.

Appropriate examination of any joint fluid present is necessary to exclude a septic process.

A marked increase in pain accompanied by local swelling, further restriction of joint motion, fever, and malaise are suggestive of septic arthritis. If this complication occurs and the diagnosis of sepsis is confirmed, appropriate antimicrobial therapy should be instituted.

Local injection of a steroid into a previously infected joint is to be avoided. Corticosteroids should not be injected into unstable joints.

Although controlled clinical trials have shown corticosteroids to be effective in speeding the resolution of acute exacerbations of multiple sclerosis they do not show that they

affect the ultimate outcome or natural history of the disease. The studies do show that relatively high doses of corticosteroids are necessary to demonstrate a significant effect. (See Dosage and Administration Section).

Since complications of treatment with glucocorticoids are dependent on the size of the dose and the duration of treatment a risk/benefit decision must be made in each individual case as to dose and duration of treatment and as to whether daily or intermittent therapy should be used.

ADVERSE REACTIONS

To report SUSPECTED ADVERSE REACTIONS, contact Somerset Therapeutics, LLC at 1-800-417-9175 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Fluid and electrolyte disturbances:

Sodium retention

Fluid retention

Congestive heart failure in susceptible patients

Potassium loss

Hypokalemic alkalosis

Hypertension

Musculoskeletal:

Muscle weakness

Steroid myopathy

Loss of muscle mass

Osteoporosis

Vertebral compression fractures

Aseptic necrosis of femoral and humeral heads

Pathologic fracture of long bones

Gastrointestinal:

Peptic ulcer with possible subsequent perforation and hemorrhage

Pancreatitis

Abdominal distention

Ulcerative esophagitis

Dermatological:

Impaired wound healing

Thin fragile skin

Facial erythema

Increased sweating

May suppress reactions to skin tests

Petechiae and ecchymoses

Neurological:

Convulsions

Increased intracranial pressure with papilledema (pseudotumor cerebri) usually after treatment

Vertigo

Headache

Ophthalmic:

Posterior subcapsular cataracts

Increased intraocular pressure

Glaucoma

Endocrine:

Menstrual irregularities

Development of cushingoid state

Suppression of growth in children

Secondary adrenocortical and pituitary unresponsiveness, particularly in times of stress, as in trauma, surgery, or illness

Decreased carbohydrate tolerance

Manifestations of latent diabetes mellitus

Increased requirements for insulin or oral hypoglycemic agents in diabetics

Metabolic:

Negative nitrogen balance due to protein catabolism

Miscellaneous:

Hyperpigmentation or hypopigmentation

Subcutaneous and cutaneous atrophy

Sterile abscess

Post-injection flare, following intra-articular use

Charcot-like arthropathy

Itching, burning, tingling in the ano-genital region

DOSAGE AND ADMINISTRATION

A. Intravenous or Intramuscular Administration

The initial dosage of dexamethasone sodium phosphate injection may vary from 0.50 mg/day to 9.0 mg/day depending on the specific disease entity being treated. In situations of less severity, lower doses will generally suffice, while in selected patients higher initial doses may be required. Usually the parenteral dosage ranges are one-third to one-half the oral dose given every 12 hours. However, in certain overwhelming, acute, life-threatening situations, administration of dosages exceeding the usual dosages may be justified and may be in multiples of the oral dosages.

For the treatment of unresponsive shock high pharmacologic doses of this product are currently

recommended. Reported regimens range from 1 to 6 mg/kg of body weight as a single intravenous injection to 40 mg initially followed by repeat intravenous injection every 2 to 6 hours while shock persists.

For the treatment of cerebral edema in adults an initial intravenous dose of 10 mg is recommended followed by 4 mg intramuscularly every six hours until maximum response has been noted. This regimen may be continued for several days postoperatively in patients requiring brain surgery. Oral dexamethasone, 1 to 3 mg t.i.d., should be given as soon as possible and dosage tapered off over a period of five to seven days. Nonoperative cases may require continuous therapy to remain free of symptoms of increased intracranial pressure. The smallest effective dose should be used in children, preferably orally. This may approximate 0.2 mg/kg/24 hours in divided doses.

In treatment of acute exacerbations of multiple sclerosis daily doses of 200 mg of prednisolone for a week followed by 80 mg every other day or 4–8 mg dexamethasone every other day for 1 month have been shown to be effective.

The initial dosage should be maintained or adjusted until a satisfactory response is noted. If after a reasonable period of time there is a lack of satisfactory clinical response, dexamethasone sodium phosphate injection should be discontinued and the patient transferred to other appropriate therapy. It should be emphasized that dosage requirements are variable and must be individualized on the basis of the disease under treatment and the response of the patient.

After a favorable response is noted, the proper maintenance dosage should be determined by decreasing the initial drug dosage in small decrements at appropriate time intervals until the lowest dosage which will maintain an adequate clinical response is reached. It should be kept in mind that constant monitoring is needed in regard to drug dosage. Included in the situations which may make dosage adjustments necessary are changes in clinical status secondary to remissions or exacerbations in the disease process, the patient's individual drug responsiveness and the effect of patient exposure to stressful situations not directly related to the disease entity under treatment. In this later situation it may be necessary to increase the dosage of dexamethasone sodium phosphate injection for a period of time consistent with the patient's condition. If after a long-term therapy the drug is to be stopped, it is recommended that it be withdrawn gradually rather than abruptly.

B. Intra-articular, soft tissue or intralesional administration.

The dose for intrasynovial administration is usually 2 to 4 mg for large joints and 0.8 to 1

mg for small joints. For soft tissue and bursal injections a dose of 2 to 4 mg is recommended. Ganglia require a dose of 1 to 2 mg. A dose of 0.4 to 1 mg is used for injection into tendon sheaths. Injection into intervertebral joints should not be attempted at any time and hip joint injection cannot be recommended as an office procedure.

Intrasynovial and soft tissue injections should be employed only when affected areas are limited to 1 or 2 sites. It should be remembered that corticoids provide palliation only and that other conventional or curative methods of therapy should be employed when indicated.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever the solution and container permit.

Frequency of injection usually ranges from once every 3 to 5 days to once every 2 to 3 weeks. Frequent intra-articular injection may cause damage to joint tissue.

HOW SUPPLIED

Dexamethasone Sodium Phosphate Injection, USP (**Preservative Free**) equivalent to 10 mg dexamethasone phosphate, is supplied in a single dose vial as follows:

NDC No	Strength	Vial Size
70069- 021 -25	10 mg/mL	1 mL Vial Packaged in twenty-fives.

The vial stopper closure is not made with natural rubber latex.

Storage

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Sensitive to heat.

Do not autoclave.

Protect from freezing.

Protect from light.

Single dose vials -Store in container until time of use. Discard unused portion.

SPL UNCLASSIFIED

Manufactured for:

Somerset Therapeutics, LLC.

Somerset, NJ 08873

Made in India

Code No.: KR/DRUGS/KTK/28/289/97

1200828

ST-DEX21/P/06

Revised: 02/2024

Made in India

Neutral Code No: TN/DRUGS/616/1996

1200832

ST-DEX-MPPL/P/01

Revised: 09/2024

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL

Container label - 1 mL

<p>NDC 70069-021-01 Rx only Dexamethasone Sodium Phosphate Injection, USP 10 mg/mL* For IM or IV Use Only 1 mL Single Dose Vial Preservative Free, Discard unused portion Sterile</p>	<p>*Each mL contains: Dexamethasone Sodium Phosphate, USP equivalent to 10 mg Dexamethasone Phosphate. PROTECT FROM LIGHT Manufactured for: Somerset Therapeutics, LLC. Somerset, NJ 08873 Made in India Code No.:KR/DRUGS/KTK/28/289/97</p>	<p>1200826 ST-DEX21/L/03 3 7 0 0 6 9 1 0 2 1 0 1 9</p>	<p>Keep area blank and varnish free for overprinting LOT and EXP 4.5 x 15mm</p>
---	--	--	---

<p>NDC 70069-021-01 Rx only Dexamethasone Sodium Phosphate Injection, USP 10 mg/mL* For IM or IV Use Only 1 mL Single Dose Vial Preservative Free, Discard unused portion Sterile</p>	<p>*Each mL contains: Dexamethasone Sodium Phosphate, USP equivalent to 10 mg Dexamethasone Phosphate. PROTECT FROM LIGHT Manufactured for: Somerset Therapeutics, LLC Somerset, NJ 08873 Made in India Neutral Code No: TN/DRUGS/616/1996</p>	<p>1200833 ST-DEX-MPPL/L/01 3 7 0 0 6 9 1 0 2 1 0 1 9</p>	<p>Unvarnished Area 13.5 x 10 mm Lot : Exp :</p>
---	--	---	--

Carton label - 1 mL

<p>*Each mL contains: Dexamethasone Sodium Phosphate, USP equivalent to 10 mg Dexamethasone Phosphate, 24.75 mg Sodium Citrate Dihydrate, Water for Injection, q.s. pH adjusted with Citric Acid or Sodium Hydroxide. The vial stopper closure is not made with natural rubber latex. Usual dosage: See Package Insert. Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Store container in carton until time of use. Discard unused portion. PROTECT FROM LIGHT.</p>	<p>Manufactured for: Somerset Therapeutics, LLC Somerset, NJ 08873 Customer Care # 1-800-417-9175 Made in India Code No: KR/DRUGS/KTK/28/289/97 1200827</p>	<p>Rx only NDC 70069-021-01 25 x 1 mL Single Dose Vials Dexamethasone Sodium Phosphate Injection, USP 10 mg/mL* For Intramuscular or Intravenous Use Only Preservative Free Sterile Somerset Therapeutics, LLC</p>	<p>3 7 0 0 6 9 1 0 2 1 0 1 9</p>
<p>*Each mL contains: Dexamethasone Sodium Phosphate, USP equivalent to 10 mg Dexamethasone Phosphate, 24.75 mg Sodium Citrate Dihydrate, Water for Injection, q.s. pH adjusted with Citric Acid or Sodium Hydroxide. The vial stopper closure is not made with natural rubber latex. Usual dosage: See Package Insert. Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Store container in carton until time of use. Discard unused portion. PROTECT FROM LIGHT.</p>	<p>Manufactured for: Somerset Therapeutics, LLC Somerset, NJ 08873 Customer Care # 1-800-417-9175 Made in India Neutral Code No: TN/DRUGS/616/1996 1200831</p>	<p>Rx only NDC 70069-021-01 25 x 1 mL Single Dose Vials Dexamethasone Sodium Phosphate Injection, USP 10 mg/mL* For Intramuscular or Intravenous Use Only Preservative Free Sterile Somerset Therapeutics, LLC</p>	<p>3 7 0 0 6 9 1 0 2 1 0 1 9 Unvarnished area for lot/exp and serialization barcode data GTIN S/N Lot Exp 50 x 30 mm</p>

DEXAMETHASONE SODIUM PHOSPHATE

dexamethasone sodium phosphate injection

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:70069-021
Route of Administration	INTRAMUSCULAR, INTRAVENOUS		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXAMETHASONE SODIUM PHOSPHATE (UNII: AI9376Y64P) (DEXAMETHASONE - UNII: 7S5I7G3JQL)	DEXAMETHASONE PHOSPHATE	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	24.75 mg in 1 mL
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (Clear, colorless solution)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70069-021-25	25 in 1 CARTON	04/19/2018	
1	NDC:70069-021-01	1 mL in 1 VIAL; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA207442	04/19/2018	

Labeler - Somerset Therapeutics, LLC (079947873)

Registrant - Somerset Therapeutics, LLC (079947873)

Establishment

Name	Address	ID/FEI	Business Operations
Somerset Therapeutics Private Limited		677236695	ANALYSIS(70069-021) , LABEL(70069-021) , PACK(70069-021) , MANUFACTURE(70069-021)

Establishment

Name	Address	ID/FEI	Business Operations
Maiva Pharma Private Limited		725656438	ANALYSIS(70069-021) , LABEL(70069-021) , MANUFACTURE(70069-021) , PACK(70069-021)

Revised: 12/2024

Somerset Therapeutics, LLC