

**BUMETANIDE- bumetanide tablet**  
**First Nation Group, LLC**

-----  
**Bumetanide Tablets USP**

**SPL UNCLASSIFIED**

Bumetanide Tablets USP, 0.5 mg, 1 mg, and 2 mg

**For oral use**

**Rx Only**


**BOXED WARNING**

Bumetanide tablets are potent diuretics which, if given in excessive amounts, can lead to a profound diuresis with water and electrolyte depletion. Therefore, careful medical supervision is required, and dose and dosage schedule have to be adjusted to the individual patient's needs (*see DOSAGE AND ADMINISTRATION*).

**DESCRIPTION**

Bumetanide tablets USP are loop diuretics available as 0.5 mg, 1 mg, and 2 mg (white to off white) tablets for oral administration; each tablet also contains corn starch, lactose monohydrate, magnesium oxide light, magnesium stearate, microcrystalline cellulose, pregelatinized starch, sodium lauryl sulfate, and talc.

Chemically, bumetanide is 3-(butylamino)-4-phenoxy-5-sulfamoylbenzoic acid. It is a practically white powder having a calculated molecular weight of 364.42, and the following structural formula:



FDA approved dissolution test specifications differ from USP.

**CLINICAL PHARMACOLOGY**

Bumetanide tablets are loop diuretics with a rapid onset and short duration of action.

Pharmacological and clinical studies have shown that 1 mg bumetanide tablets have a diuretic potency equivalent to approximately 40 mg furosemide. The major site of bumetanide tablets action is the ascending limb of the loop of Henle.

The mode of action has been determined through various clearance studies in both humans and experimental animals. Bumetanide tablets inhibit sodium reabsorption in the ascending limb of the loop of Henle, as shown by marked reduction of free-water clearance ( $CH_2O$ ) during hydration and tubular free-water reabsorption ( $T^C H_2O$ ) during hydropenia. Reabsorption of chloride in the ascending limb is also blocked by bumetanide tablets, and bumetanide tablets are somewhat more chloruretic than natriuretic.

Potassium excretion is also increased by bumetanide tablets, in a dose-related fashion.

Bumetanide tablets may have an additional action in the proximal tubule. Since phosphate reabsorption takes place largely in the proximal tubule, phosphaturia during bumetanide tablets induced diuresis is indicative of this additional action. This is further supported by the reduction in the renal clearance of bumetanide tablets by probenecid, associated with diminution in the natriuretic response. This proximal tubular activity does not seem to be related to an inhibition of carbonic anhydrase. Bumetanide tablets do not appear to have a noticeable action on the distal tubule.

Bumetanide tablets decrease uric acid excretion and increase serum uric acid. Following oral administration of bumetanide tablets the onset of diuresis occurs in 30 to 60 minutes. Peak activity is reached between 1 and 2 hours. At usual doses (1 mg to 2 mg) diuresis is largely complete within 4 hours; with higher doses, the diuretic action lasts for 4 to 6 hours. Diuresis starts within minutes following an intravenous injection and reaches maximum levels within 15 to 30 minutes.

Several pharmacokinetic studies have shown that bumetanide, administered orally or parenterally, is eliminated rapidly in humans, with a half-life of between 1 and 1½ hours. Plasma protein-binding is in the range of 94% to 96%.

Oral administration of carbon-14 labeled bumetanide tablets to human volunteers revealed that 81% of the administered radioactivity was excreted in the urine, 45% of it as unchanged drug. Urinary and biliary metabolites identified in this study were formed by oxidation of the N-butyl side chain. Biliary excretion of bumetanide tablets amounted to only 2% of the administered dose.

## **Pediatric Pharmacology**

Elimination of bumetanide appears to be considerably slower in neonatal patients compared with adults, possibly because of immature renal and hepatobiliary function in this population. Small pharmacokinetic studies of intravenous bumetanide in preterm and full-term neonates with respiratory disorders have reported an apparent half-life of approximately 6 hours, with a range up to 15 hours and a serum clearance ranging from 0.2 mL/min/kg to 1.1 mL/min/kg. In a population of neonates receiving bumetanide for volume overload, mean serum clearance rates were 2.2 mL/min/kg in patients less than 2 months of age and 3.8 mL/min/kg in patients aged 2 to 6 months. Mean serum half-life of bumetanide was 2.5 hours and 1.5 hours in patients aged less than 2 months and those aged 2 to 6 months, respectively. Elimination half-life decreased considerably during the first month of life, from a mean of approximately 6 hours at birth to approximately 2.4 hours at 1 month of age.

In preterm neonates, mean serum concentrations following a single 0.05 mg/kg dose ranged from 126 µg/L at 1 hour to 57 µg/L at 8 hours. In another study, mean serum concentrations following a single 0.05 mg/kg dose were 338 ng/mL at 30 minutes and 176 ng/mL after 4 hours. A single dose of 0.1 mg/kg produced mean serum levels of 314 ng/mL at 1 hour, and 195 ng/mL at 6 hours. Mean volume of distribution in neonates and infants has been reported to range from 0.26 L/kg to 0.39 L/kg.

The degree of protein binding of bumetanide in cord sera from healthy neonates was approximately 97%, suggesting the potential for bilirubin displacement. A study using pooled sera from critically ill neonates found that bumetanide at concentrations of 0.5 µg/mL to 50 µg/mL, but not 0.25 µg/mL, caused a linear increase in unbound bilirubin concentrations.

In 56 infants aged 4 days to 6 months, bumetanide doses ranging from 0.005 mg/kg to 0.1 mg/kg were studied for pharmacodynamic effect. Peak bumetanide excretion rates increased linearly with increasing doses of drug. Maximal diuretic effect was observed at a bumetanide excretion rate of about 7 µg/kg/h, corresponding to doses of 0.035 mg/kg to 0.040 mg/kg. Higher doses produced a higher bumetanide excretion rate but no increase in diuretic effect. Urine flow rate peaked during the first hour after drug administration in 80% of patients and by 3 hours in all patients.

### **Geriatric Pharmacology**

In a group of ten geriatric subjects between the ages of 65 and 73 years, total bumetanide clearance was significantly lower ( $1.8 \pm 0.3$  mL/min/kg) compared with younger subjects ( $2.9 \pm 0.2$  mL/min/kg) after a single oral bumetanide 0.5 mg dose. Maximum plasma concentrations were higher in geriatric subjects ( $16.9 \pm 1.8$  ng/mL) compared with younger subjects ( $10.3 \pm 1.5$  ng/mL). Urine flow rate and total excretion of sodium and potassium were increased less in the geriatric subjects compared with younger subjects, although potassium excretion and fractional sodium excretion were similar between the two age groups. Nonrenal clearance, bioavailability, and volume of distribution were not significantly different between the two groups.

### **INDICATIONS AND USAGE**

Bumetanide tablets are indicated for the treatment of edema associated with congestive heart failure, hepatic and renal disease, including the nephrotic syndrome.

Almost equal diuretic response occurs after oral and parenteral administration of bumetanide. Therefore, if impaired gastrointestinal absorption is suspected or oral administration is not practical, bumetanide should be given by the intramuscular or intravenous route.

Successful treatment with bumetanide tablets following instances of allergic reactions to furosemide suggests a lack of cross-sensitivity.

### **CONTRAINDICATIONS**

Bumetanide tablets are contraindicated in anuria. Although bumetanide tablets can be used to induce diuresis in renal insufficiency, any marked increase in blood urea nitrogen or creatinine, or the development of oliguria during therapy of patients with progressive renal disease, is an indication for discontinuation of treatment with bumetanide tablets.

Bumetanide tablets are also contraindicated in patients in hepatic coma or in states of severe electrolyte depletion until the condition is improved or corrected. Bumetanide tablets are contraindicated in patients hypersensitive to this drug.

## **WARNINGS**

### **Volume and Electrolyte Depletion**

The dose of bumetanide tablets should be adjusted to the patient's need. Excessive doses or too frequent administration can lead to profound water loss, electrolyte depletion, dehydration, reduction in blood volume and circulatory collapse with the possibility of vascular thrombosis and embolism, particularly in elderly patients.

### **Hypokalemia**

Hypokalemia can occur as a consequence of bumetanide tablets administration. Prevention of hypokalemia requires particular attention in the following conditions: patients receiving digitalis and diuretics for congestive heart failure, hepatic cirrhosis and ascites, states of aldosterone excess with normal renal function, potassium-losing nephropathy, certain diarrheal states, or other states where hypokalemia is thought to represent particular added risks to the patient, i.e., history of ventricular arrhythmias.

In patients with hepatic cirrhosis and ascites, sudden alterations of electrolyte balance may precipitate hepatic encephalopathy and coma. Treatment in such patients is best initiated in the hospital with small doses and careful monitoring of the patient's clinical status and electrolyte balance. Supplemental potassium and/or spironolactone may prevent hypokalemia and metabolic alkalosis in these patients.

### **Ototoxicity**

In cats, dogs and guinea pigs, bumetanide has been shown to produce ototoxicity. In these test animals' bumetanide was 5 to 6 times more potent than furosemide and, since the diuretic potency of bumetanide is about 40 to 60 times furosemide, it is anticipated that blood levels necessary to produce ototoxicity will rarely be achieved. The potential exists, however, and must be considered a risk of intravenous therapy, especially at high doses, repeated frequently in the face of renal excretory function impairment. Potentiation of aminoglycoside ototoxicity has not been tested for bumetanide. Like other members of this class of diuretics, bumetanide probably shares this risk.

### **Allergy to Sulfonamides**

Patients allergic to sulfonamides may show hypersensitivity to bumetanide tablets.

### **Thrombocytopenia**

Since there have been rare spontaneous reports of thrombocytopenia from postmarketing experience, patients should be observed regularly for possible occurrence of thrombocytopenia.

## **PRECAUTIONS**

## **General**

Serum potassium should be measured periodically and potassium supplements or potassium sparing diuretics added if necessary. Periodic determinations of other electrolytes are advised in patients treated with high doses or for prolonged periods, particularly in those on low-salt diets.

Hyperuricemia may occur; it has been asymptomatic in cases reported to date. Reversible elevations of the BUN and creatinine may also occur, especially in association with dehydration and particularly in patients with renal insufficiency. Bumetanide tablets may increase urinary calcium excretion with resultant hypocalcemia.

Diuretics have been shown to increase the urinary excretion of magnesium; this may result in hypomagnesemia.

## **Laboratory Tests**

Studies in normal subjects receiving bumetanide tablets revealed no adverse effects on glucose tolerance, plasma insulin, glucagon and growth hormone levels, but the possibility of an effect on glucose metabolism exists. Periodic determinations of blood sugar should be done, particularly in patients with diabetes or suspected latent diabetes.

Patients under treatment should be observed regularly for possible occurrence of blood dyscrasias, liver damage or idiosyncratic reactions, which have been reported occasionally in foreign marketing experience. The relationship of these occurrences to bumetanide tablets use is not certain.

## **Drug Interactions**

Drugs with Ototoxic Potential (*see WARNINGS*)

Especially in the presence of impaired renal function, the use of parenterally administered bumetanide in patients to whom aminoglycoside antibiotics are also being given should be avoided, except in life-threatening conditions.

Drugs with Nephrotoxic Potential

There has been no experience with the concurrent use of bumetanide tablets with drugs known to have a nephrotoxic potential. Therefore, the simultaneous administration of these drugs should be avoided.

Lithium

Lithium should generally not be given with diuretics (such as bumetanide tablets) because they reduce its renal clearance and add a high risk of lithium toxicity.

Probenecid

Pretreatment with probenecid reduces both the natriuresis and hyperreninemia produced by bumetanide tablets. This antagonistic effect of probenecid on bumetanide tablets natriuresis is not due to a direct action on sodium excretion but is probably secondary to its inhibitory effect on renal tubular secretion of bumetanide. Thus, probenecid should not be administered concurrently with bumetanide tablets.

Indomethacin

Indomethacin blunts the increases in urine volume and sodium excretion seen during

bumetanide tablets treatment and inhibits the bumetanide-induced increase in plasma renin activity. Concurrent therapy with bumetanide tablets are thus not recommended.

### Antihypertensives

Bumetanide tablets may potentiate the effect of various antihypertensive drugs, necessitating a reduction in the dosage of these drugs.

### Digoxin

Interaction studies in humans have shown no effect on digoxin blood levels.

### Anticoagulants

Interaction studies in humans have shown bumetanide tablets to have no effect on warfarin metabolism or on plasma prothrombin activity.

## **Carcinogenesis, Mutagenesis and Impairment of Fertility**

Bumetanide tablets were devoid of mutagenic activity in various strains of *Salmonella typhimurium* when tested in the presence or absence of an *in vitro* metabolic activation system. An 18-month study showed an increase in mammary adenomas of questionable significance in female rats receiving oral doses of 60 mg/kg/day (2000 times a 2-mg human dose). A repeat study at the same doses failed to duplicate this finding.

Reproduction studies were performed to evaluate general reproductive performance and fertility in rats at oral dose levels of 10, 30, 60 or 100 mg/kg/day. The pregnancy rate was slightly decreased in the treated animals; however, the differences were small and not statistically significant.

## **Pregnancy**

### **Teratogenic Effects**

Bumetanide tablets are neither teratogenic nor embryocidal in mice when given in doses up to 3400 times the maximum human therapeutic dose.

Bumetanide tablets have been shown to be nonteratogenic, but it has a slight embryocidal effect in rats when given in doses of 3400 times the maximum human therapeutic dose and in rabbits at doses of 3.4 times the maximum human therapeutic dose. In one study, moderate growth retardation and increased incidence of delayed ossification of sternebrae were observed in rats at oral doses of 100 mg/kg/day, 3400 times the maximum human therapeutic dose. These effects were associated with maternal weight reductions noted during dosing. No such adverse effects were observed at 30 mg/kg/day (1000 times the maximum human therapeutic dose). No fetotoxicity was observed at 1000 to 2000 times the human therapeutic dose.

In rabbits, a dose-related decrease in litter size and an increase in resorption rate were noted at oral doses of 0.1 mg/kg/day and 0.3 mg/kg/day (3.4 and 10 times the maximum human therapeutic dose). A slightly increased incidence of delayed ossification of sternebrae occurred at 0.3 mg/kg/day; however, no such adverse effects were observed at the dose of 0.03 mg/kg/day. The sensitivity of the rabbit to bumetanide tablets parallels the marked pharmacologic and toxicologic effects of the drug in this species.

Bumetanide tablets were not teratogenic in the hamster at an oral dose of 0.5

mg/kg/day (17 times the maximum human therapeutic dose). Bumetanide was not teratogenic when given intravenously to mice and rats at doses up to 140 times the maximum human therapeutic dose.

There are no adequate and well-controlled studies in pregnant women. A small investigational experience in the United States and marketing experience in other countries to date have not indicated any evidence of adverse effects on the fetus, but these data do not rule out the possibility of harmful effects. Bumetanide tablets should be given to a pregnant woman only if the potential benefit justifies the potential risk to the fetus.

### **Nursing Mothers**

It is not known whether this drug is excreted in human milk. As a general rule, nursing should not be undertaken while the patient is on bumetanide tablets since it may be excreted in human milk.

### **Pediatric Use**

Safety and effectiveness in pediatric patients below the age of 18 have not been established.

*In vitro* studies using pooled sera from critically ill neonates have shown bumetanide to be a potent displacer of bilirubin (see *CLINICAL PHARMACOLOGY: Pediatric Pharmacology*). The administration of bumetanide could present a particular concern if given to critically ill or jaundiced neonates at risk for kernicterus.

### **Geriatric Use**

Clinical studies of bumetanide tablets did not include sufficient numbers of subjects aged 65 and over to determine whether they responded differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

This drug is known to be substantially excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

## **ADVERSE REACTIONS**

The most frequent clinical adverse reactions considered probably or possibly related to bumetanide tablets are muscle cramps (seen in 1.1% of treated patients), dizziness (1.1%), hypotension (0.8%), headache (0.6%), nausea (0.6%) and encephalopathy (in patients with pre-existing liver disease) (0.6%). One or more of these adverse reactions have been reported in approximately 4.1% of patients treated with bumetanide tablets.

Serious skin reactions (i.e., Stevens-Johnson syndrome, toxic epidermal necrolysis) have been reported in association with bumetanide use.

Less frequent clinical adverse reactions to bumetanide tablets are impaired hearing (0.5%), pruritus (0.4%), electrocardiogram changes (0.4%), weakness (0.2%), hives (0.2%), abdominal pain (0.2%), arthritic pain (0.2%), musculoskeletal pain (0.2%), rash (0.2%) and vomiting (0.2%). One or more of these adverse reactions have been reported in approximately 2.9% of patients treated with bumetanide tablets.

Other clinical adverse reactions, which have each occurred in approximately 0.1% of patients, are vertigo, chest pain, ear discomfort, fatigue, dehydration, sweating, hyperventilation, dry mouth, upset stomach, renal failure, asterixis, itching, nipple tenderness, diarrhea, premature ejaculation and difficulty maintaining an erection.

Laboratory abnormalities reported have included hyperuricemia (in 18.4% of patients tested), hypochloremia (14.9%), hypokalemia (14.7%), azotemia (10.6%), hyponatremia (9.2%), increased serum creatinine (7.4%), hyperglycemia (6.6%), and variations in phosphorus (4.5%), CO content (4.3%), bicarbonate (3.1%) and calcium (2.4%). Although manifestations of the pharmacologic action of bumetanide tablets, these conditions may become more pronounced by intensive therapy.

Also reported have been thrombocytopenia (0.2%) and deviations in hemoglobin (0.8%), prothrombin time (0.8%), hematocrit (0.6%), WBC (0.3%) and differential counts (0.1%). There have been rare spontaneous reports of thrombocytopenia from postmarketing experience.

Diuresis induced by bumetanide tablets may also rarely be accompanied by changes in LDH (1.0%), total serum bilirubin (0.8%), serum proteins (0.7%), SGOT (0.6%), SGPT (0.5%), alkaline phosphatase (0.4%), cholesterol (0.4%) and creatinine clearance (0.3%). Increases in urinary glucose (0.7%) and urinary protein (0.3%) have also been seen.

**To report SUSPECTED ADVERSE REACTIONS, contact First Nation Group, LLC at 1-855-221-5332 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

## **OVERDOSAGE**

Overdosage can lead to acute profound water loss, volume and electrolyte depletion, dehydration, reduction of blood volume and circulatory collapse with a possibility of vascular thrombosis and embolism. Electrolyte depletion may be manifested by weakness, dizziness, mental confusion, anorexia, lethargy, vomiting and cramps. Treatment consists of replacement of fluid and electrolyte losses by careful monitoring of the urine and electrolyte output and serum electrolyte levels.

## **DOSAGE AND ADMINISTRATION**

Individualize dosage with careful monitoring of patient response.

### **Oral Administration**

The usual total daily dosage of bumetanide tablets USP is 0.5 mg to 2 mg and in most patients is given as a single dose.

If the diuretic response to an initial dose of bumetanide tablets USP is not adequate, in view of its rapid onset and short duration of action, a second or third dose may be given at 4- to 5-hour intervals up to a maximum daily dose of 10 mg. An intermittent dose schedule, whereby bumetanide tablets USP are given on alternate days or for 3 to 4

days with rest periods of 1 to 2 days in between, is recommended as the safest and most effective method for the continued control of edema. In patients with hepatic failure, keep the dosage to a minimum.

Because cross-sensitivity with furosemide has rarely been observed, bumetanide can be substituted at approximately a 1:40 ratio of bumetanide in proportion to furosemide in patients allergic to furosemide.

### **Parenteral Administration**

Bumetanide injection may be administered parenterally (intravenously and intramuscularly) to patients in whom gastrointestinal absorption may be impaired or in whom oral administration is not practical.

Terminate parenteral treatment and institute oral treatment as soon as possible.

### **HOW SUPPLIED**

Bumetanide tablets USP are supplied as 0.5 mg, 1 mg, and 2 mg tablets for oral administration.

Bumetanide tablets USP 0.5 mg are available as white to off white, elliptical, flat-faced, beveled edged tablets, engraved 'LH' on one side and plain on reverse side.

NDC 81469-221-01

in bottles of 100 tablets with child-resistant closure.

Bumetanide tablets USP 1 mg are available as white to off white, elliptical, flat-faced, beveled edged tablets, engraved 'LI' on one side and plain on reverse side.

NDC 81469-222-01

in bottles of 100 tablets with child-resistant closure

NDC 81469-222-05

in bottles of 500 tablets

Bumetanide tablets USP 2 mg are available as white to off white, elliptical, flat-faced, beveled edged tablets, engraved 'LJ' on one side and plain on reverse side.

NDC 81469-223-01

in bottles of 100 tablets with child-resistant closure

NDC 81469-223-05

in bottles of 500 tablets

Store at 20°C to 25°C (68°F to 77°F); excursions permitted to 15°C to 30°C (59°F to 86°F) [See USP Controlled Room Temperature].

Dispense contents in a tight, light-resistant container as defined in the USP with a child-resistant closure, as required.

### **Manufactured for:**

First Nation Group, LLC

4566 E Highway 20,  
Suite #208  
Niceville, FL 32578-8839

**Product of Italy**

**Revised:** September 2025, V-01

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 81469-221-01

Bumetanide Tablets USP

0.5 mg

100 Tablets

Rx Only

**NDC 81469-221-01**

**Bumetanide  
Tablets USP**

**0.5 mg**

Rx ONLY                      100 Tablets

**FIRST NATION  
GROUP**

Each tablet contains:  
Bumetanide, USP.....0.5 mg

USUAL DOSAGE:  
See prescribing information.  
Dispense in tight, light-resistant  
containers as defined in USP.

Store at 20° to 25°C  
(68° to 77°F); excursions  
permitted between  
15° to 30°C (59° to 86°F)  
[See USP Controlled  
Room Temperature].

KEEP THIS AND ALL  
DRUGS OUT OF THE  
REACH OF CHILDREN.

Rev: 09/25,V-01  
ECFEOTL303US329500

Manufactured for:  
**First Nation Group, LLC**  
**4566 E Highway 20, Suite #208**  
**Niceville, FL 32578-8839**  
Product of Italy  
Code No.: HP/Drugs/N-MNB/17/194

11090

12008445

8146912210115

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 81469-222-01

Bumetanide Tablets USP

1 mg

100 Tablets

Rx Only

**NDC 81469-222-01**

# Bumetanide Tablets USP

**1 mg**

Rx ONLY      100 Tablets



Each tablet contains:  
Bumetanide, USP.....1 mg


USUAL DOSAGE:  
See prescribing information.  
Dispense in tight, light-resistant containers as defined in USP.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Rev: 09/25,V-01  
ECFEITL303US329600

Manufactured for:  
**First Nation Group, LLC**  
4566 E Highway 20, Suite #208  
Niceville, FL 32578-8839  
Product of Italy  
Code No.: HP/Drugs/N-MNB/17/194

11091  
12008446  
N 8 1 4 6 9 1 2 2 0 1 2

NDC 81469-222-05

Bumetanide Tablets USP

1 mg

500 Tablets

Rx Only

**NDC 81469-222-05**

# Bumetanide Tablets USP

**1 mg**

Rx ONLY      500 Tablets



Each tablet contains:  
Bumetanide, USP.....1 mg


USUAL DOSAGE:  
See prescribing information.  
Dispense in tight, light-resistant containers as defined in USP.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.  
This container is not intended for household use.

Code No.: HP/Drugs/N-MNB/17/194  
ECFEITL303US329700      Rev: 09/25,V-01

Manufactured for:  
**First Nation Group, LLC**  
4566 E Highway 20, Suite #208  
Niceville, FL 32578-8839  
Product of Italy

11092  
12008447  
N 8 1 4 6 9 1 2 2 0 5 0

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

NDC 81469-223-01

Bumetanide Tablets USP

2 mg

100 Tablets

Rx Only

**NDC 81469-223-01**

# Bumetanide Tablets USP

**2 mg**

Rx ONLY      100 Tablets



Each tablet contains:  
Bumetanide, USP.....2 mg


USUAL DOSAGE:  
See prescribing information.  
Dispense in tight, light-resistant containers as defined in USP.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.

Rev: 09/25,V-01  
ECFE3TL303US329800

Manufactured for:  
**First Nation Group, LLC**  
4566 E Highway 20, Suite #208  
Niceville, FL 32578-8839  
Product of Italy  
Code No.: HP/Drugs/N-MNB/17/194

11093  
12008448  
N 8146912301 9

NDC 81469-223-05

Bumetanide Tablets USP

2 mg

500 Tablets

Rx Only

**NDC 81469-223-05**

# Bumetanide Tablets USP

**2 mg**

Rx ONLY      500 Tablets



Each tablet contains:  
Bumetanide, USP.....2 mg

USUAL DOSAGE:  
See prescribing information.  
Dispense in tight, light-resistant containers as defined in USP.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° to 30°C (59° to 86°F) [See USP Controlled Room Temperature].

KEEP THIS AND ALL DRUGS OUT OF THE REACH OF CHILDREN.  
This container is not intended for household use.

Rev: 09/25,V-01  
ECFE4TL303US329900

Manufactured for:  
**First Nation Group, LLC**  
4566 E Highway 20, Suite #208  
Niceville, FL 32578-8839  
Product of Italy  
Code No.: HP/Drugs/N-MNB/17/194




11094  
12008449  
N 8146912305 7

## BUMETANIDE

bumetanide tablet

### Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:81469-221
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BUMETANIDE</b> (UNII: 0Y2S3XUQ5H) (BUMETANIDE - UNII:0Y2S3XUQ5H)	BUMETANIDE	0.5 mg

Inactive Ingredients	
Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>MAGNESIUM OXIDE</b> (UNII: 3A3U0GI71G)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

Product Characteristics			
<b>Color</b>	WHITE (white to off white)	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	8mm
<b>Flavor</b>		<b>Imprint Code</b>	LH
<b>Contains</b>			

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81469-221-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2026	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA219291	02/19/2026	

<b>BUMETANIDE</b>			
bumetanide tablet			
Product Information			
<b>Product Type</b>	HUMAN PRESCRIPTION DRUG	<b>Item Code (Source)</b>	NDC:81469-222
<b>Route of Administration</b>	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>BUMETANIDE</b> (UNII: 0Y2S3XUQ5H) (BUMETANIDE - UNII:0Y2S3XUQ5H)	BUMETANIDE	1 mg	

## Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	

## Product Characteristics

Color	WHITE (white to off white)	Score	no score
Shape	OVAL	Size	11mm
Flavor		Imprint Code	LI
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81469-222-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2026	
2	NDC:81469-222-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA219291	02/19/2026	

## BUMETANIDE

bumetanide tablet

## Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:81469-223
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BUMETANIDE (UNII: 0Y2S3XUQ5H) (BUMETANIDE - UNII:0Y2S3XUQ5H)	BUMETANIDE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM OXIDE (UNII: 3A3U0GI71G)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	

## Product Characteristics

Color	WHITE (white to off white)	Score	no score
Shape	OVAL	Size	13mm
Flavor		Imprint Code	LJ
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81469-223-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2026	
2	NDC:81469-223-05	500 in 1 BOTTLE; Type 0: Not a Combination Product	02/19/2026	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA219291	02/19/2026	

**Labeler** - First Nation Group, LLC (078875731)

**Registrant** - Mankind Pharma Limited (915834068)

## Establishment

Name	Address	ID/FEI	Business Operations
Mankind Pharma Limited		916512493	MANUFACTURE(81469-221, 81469-222, 81469-223) , ANALYSIS(81469-221, 81469-222, 81469-223) , PACK(81469-221, 81469-222, 81469-223)

Revised: 2/2026

First Nation Group, LLC