

**CONRX SINUS- acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet**  
**Eagle Distributors, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ConRx™ Sinus**

**Drug Facts**

**Active ingredient (in each tablet)**

Acetaminophen 325mg

Guaifenesin 200mg

Phenylephrine HCl 5mg

**Purpose**

Pain reliever/ fever reducer

Expectorant

Nasal decongestant

**Uses**

Uses For the temporary relief of: sinus congestion and pressure, headache, nasal congestion, minor aches and pains, helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive, temporarily relieves nasal congestion due to the common cold, and hay fever and other upper respiratory allergies.

**Warnings**

- **Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: more than 8 tablets in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- If you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease

- diabetes
- trouble urinating due to an enlarged prostate gland
- a persistent chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**

- nervousness, dizziness, or sleeplessness occur
  - pain, nasal congestion or cough gets worse or lasts for more than 7 days
  - fever gets more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - cough comes back or occurs with rash or headache that lasts.
- These could be signs of serious conditions.

**If pregnant or breast-feeding, ask a health professional before use.**

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

Adults and Children 12 years and over	<ul style="list-style-type: none"> <li>• Take 1-2 tablets every 4 hours as needed</li> <li>• Do not taking more than 8 tablets in 24 hour</li> </ul>
Children under 12 years of age	Consult a doctor

**Other Information**

- Do not use if pouch is torn or damaged.
- Store between 15-30°C (59-86°F). Avoid excessive heat and humidity.
- See side panel for lot number and expiration date.

**Inactive ingredients**

Colloidal silicon dioxide, Sodium Lauryl Sulphate, Edetaet disodium, dioctyl sodium sulphosuccinate, polyvinyl pyrrollidone, gelatin, corn starch', magnesium Stearate, microcrystalline cellulose, polyethylene glycol 6000,Sodium Benzoate ,Sodium Starch Glycolate, Stearic acid, Purified Talc, Titanium Dioxide.

**PRINCIPAL DISPLAY PANEL - 2 Caplets × 50 Pouch Box**

**See New Warnings Information & Directions**

**Compare to the Active Ingredients in**

***Tylenol® Sinus\****

***Congestion & Pain\****

***ConRx™***

***SEVERE***

***SINUS***

***Headache & Nasal Congestion***

- Acetaminophen  
Headache & Pain
- Guaifenesin  
Chest Congestion
- Phenylephrine HCl  
Nasal Congestion

**TO OPEN**

**PUSH IN TAB AND PULL OUT**

**Compare to the Active Ingredients in**

***Tylenol® Sinus\****

***Congestion & Pain\****

***50 Pouches of 2 Caplets Each***

See New Warnings Information & Directions  
Compare to the Active Ingredients in

**Tylenol<sup>®</sup> Sinus\***  
Congestion & Pain\*

**ConRx<sup>™</sup> SEVERE SINUS**  
Headache & Nasal Congestion

- Acetaminophen Headache & Pain
- Guaifenesin Chest Congestion
- Phenylephrine HCl Nasal Congestion

See New Warnings Information & Directions  
Compare to the Active Ingredients in

**Tylenol<sup>®</sup> Sinus\***  
Congestion & Pain\*

**ConRx<sup>™</sup> SEVERE SINUS**  
Headache & Nasal Congestion

- Acetaminophen Headache & Pain
- Guaifenesin Chest Congestion
- Phenylephrine HCl Nasal Congestion

▲  
TO OPEN  
PUSH IN TAB AND PULL OUT

Compare to the Active Ingredients in

**Tylenol<sup>®</sup> Sinus\***  
Congestion & Pain\*

50 Pouches of 2 Caplets Each



50 Pouches of 2 Caplets Each

- Acetaminophen Headache & Pain
- Guaifenesin Chest Congestion
- Phenylephrine HCl Nasal Congestion

**ConRx<sup>™</sup> SEVERE SINUS**  
Headache & Nasal Congestion

**Tylenol<sup>®</sup> Sinus\***  
Congestion & Pain\*

See New Warnings Information & Directions  
Compare to the Active Ingredients in

**Drug Facts**

Active Ingredients (in each caplets)	Purposes
Acetaminophen 325 mg	Pain reliever
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

- Uses** ■ for the temporary relief of:
- sinus congestion and pressure
  - headache
  - nasal congestion
  - minor aches and pains
  - helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive
  - temporarily relieves nasal congestion due to the common cold, and hay fever and other upper respiratory allergies

See New Warnings Information & Directions  
Compare to the Active Ingredients in

**Tylenol<sup>®</sup> Sinus\***  
Congestion & Pain\*

**Warnings**  
**Liver warnings:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 8 tablets in 24 hours, which is the maximum daily amount  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product

**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

**Ask a doctor before use if you have**  
 ■ liver disease ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ trouble urinating due to an enlarged prostate gland  
 ■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema ■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin**

**When using this product do not exceed recommended dosage**

**Stop use and ask a doctor if**  
 ■ nervousness, dizziness, or sleeplessness occur  
 ■ pain, nasal congestion or cough gets worse or lasts for more than 7 days  
 ■ fever gets worse or lasts for more than 3 days  
 ■ redness or swelling is present  
 ■ new symptoms occur  
 ■ cough comes back or occurs with rash or headache that lasts  
 These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use.**  
**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

adults and children 12 years and over	■ take 1-2 tablets every 4 hours as needed ■ do not take more than 8 tablets in 24 hours
children under 12 years	■ consult a doctor

**Other information**  
 ■ avoid excessive heat and humidity  
 ■ store between 15-30°C (59-86°F)  
 ■ do not use if pouch is torn or open  
 ■ see side panel for lot number and expiration date

**Inactive ingredients** colloidal silicon dioxide, sodium lauryl sulphate, edetate disodium, diethyl sodium sulphosuccinate, polyvinyl pyrrolidone, gelatin, corn starch, magnesium stearate, microcrystalline cellulose, polyethylene glycol 6000, sodium benzoate, sodium starch glycolate, stearic acid, purified talc, titanium dioxide

**Questions or comments?** call 1-800-570-8650 (M-F 9 am to 5 pm PST)



**ConRx™ SEVERE SINUS**  
**Headache & Nasal Congestion**

- Acetaminophen
- Guaifenesin
- Phenylephrine HCl

Headache & Pain  
 Chest Congestion  
 Nasal Congestion

**DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN**

\*This product is not manufactured or distributed by: McNeil Consumer Speciality Pharmaceuticals Division of McNeil PPC, Inc. Eagle Distributors, Inc.,/ConRx™ does not own the Tylenol® Sinus Congestion & Pain trademark.



Product manufactured for:  
**Eagle Distributors, Inc.**  
 Los Angeles, CA 90011

Part No: 88885\_Ver:1\_2013

# CONRX SINUS

acetaminophen, guaifenesin, and phenylephrine hydrochloride tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68737-236
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>Acetaminophen</b> (UNII: 362O9ITL9D) (Acetaminophen - UNII:362O9ITL9D)	Acetaminophen	325 mg
<b>Guaifenesin</b> (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	200 mg
<b>Phenylephrine Hydrochloride</b> (UNII: 04JA59TNSJ) (Phenylephrine - UNII:1WS297W6MV)	Phenylephrine Hydrochloride	5 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>BIS(1-METHYLAMYL) SODIUM SULFO SUCCINATE</b> (UNII: 772Y8KZU65)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>POVIDONE K90</b> (UNII: RDH86HJV5Z)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL 6000 (UNII: 30IQX730WE)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

### Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	18mm
Flavor		Imprint Code	CRX
Contains			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68737-236-20	50 in 1 BOX		
1		2 in 1 POUCH		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	02/15/2013	

**Labeler** - Eagle Distributors, Inc. (929837425)

Revised: 2/2013

Eagle Distributors, Inc.