DRAMAMINE CHEWABLE DIMENHYDRINATE- dimenhydrinate tablet, chewable JC World Bell Wholesale Co., Inc.

Dramamine Chewable Dimenhydrinate

Drug Facts

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Use

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not givetochildren under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use

if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding,

ask a doctor before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- to prevent motion sickness, the first dose should be taken $\frac{1}{2}$ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

Adults and children 12 years and over	 take 1 to 2 chewable tablets every 4-6 hours do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor 			
Children 6 to under 12 years	 give ½ to 1 chewable tablet every 6-8 hours do not give more than 3 chewable tablets in 24 hours, or as directed by a doctor 			
Children 2 to under 6 years	 give ½ chewable tablet every 6-8 hours do not give more than 1 ½ chewable tablets in 24 hours, or as directed by a doctor 			

Other information

- Phenylketonurics: contains phenylalanine 0.84 mg per tablet
- store at room temperature 20-25°C (68-77°F)
- see side panel for lot number and expiration date

Inactive ingredients

anhydrous-citric-acid, aspartame, FD&C yellow no. 6 aluminum lake, flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol

Questions?

1-800-382-7219 Dramamine.com

Package Labeling:



FRONT

BACK



0.080

Drug Facts (continued) Ask a doctor before use if you have a breathing problem such as emphysema or chronic bronchits glaucoma trouble urinating due to an enlarged prostate gland. Ask a doctor or pharmacist before use if you are taking sedatives or tranquillers. When using this product marked drowsiness may occur avoid alcoholic drinks alcohol, sedatives, and tranquilzers may increase drowsiness be careful when driving a motor vehicle or operating machinery. If pregnant or breast-feeding, ask a doctor before use. Keep out of reach of children. In In pregnant or breast-rectung, ask a outcome borlo uses. Reep out or reach or chimters. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) immediately. *Directions*: to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity. To prevent or treat motion sickness: Adults and children 12 years and over: Take 1 to 2 tablets every 4 – 6 hours = do not take more than 8 chewable tablets in 24 hours, or as directed by a doctor. Children 6 to under 12 years: = give ½ to 1 tablet every 6 - 8 hours = do not give more than 3 chewable tablets 12 years. ** give 9: 10 I tablet every - 8 noturs* 40 not give more than 3 cnewand tablets in 24 hours, or as directed by a doctor. Children 2 to under \$\frac{1}{2}\$ years ** give \$\frac{1}{2}\$ tablet every \$\frac{1}{2}\$ doctor. Other information ** Phenylketonurics: contains phenylalanine 0.84 mg per tablet ** store at room temperature 20* - 25 ** 0 (88* - 77**) ** do not use if pouch is opened. Inactive ingredients anhydrous citric acid, aspartame, FD&C yellow no. 8 aluminum lake, flagues can explain a tablet ** store at room temperature 20* - 25 ** 0 (88* - 77**) ** do not use if pouch is opened. Inactive ingredients anhydrous citric acid, aspartame, FD&C yellow no. 8 aluminum lake, pouch so the property in detection productions and the production of the flavors, magnesium stearate, maltodextrin, methacrylic acid copolymer, modified starch, sorbitol Questions? 1-800-382-7219 Dramamine.com Dist. By: Medtech Products Inc

DRAMAMINE CHEWABLE DIMENHYDRINATE

ENLIVERINATE (LINII), IROZZEGEC) (O CUI ODOTUTORUVI LINI

dimenhydrinate tablet, chewable

Product Information HUMAN OTC DRUG NDC:50269-525(NDC:63029-902) **Product Type** Item Code (Source) **Route of Administration** ORAL **Active Ingredient/Active Moiety Basis of Ingredient Name** Strength Strength

DIMENHIDKINAIE (UNII: JB93/PEKOC) (8-CHLUKUI HEUPHILLINE -	DIMENHYDRINATE	50 ma
UNII:GE2UA340FM)	DIMENHIDRINATE	30 mg

Inactive Ingredients		
Ingredient Name	Strength	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)		
ASPARTAME (UNII: Z0H242BBR1)		
FD&C YELLOW NO. 6 ALUMINUM LAKE (UNII: GYP6Z2JR6Q)		
MAGNESIUM STEARATE (UNII: 70097M6I30)		
MALTODEXTRIN (UNII: 7CVR7L4A2D)		
SORBITOL (UNII: 506T60A25R)		

Product Characteristics				
Color	orange	Score	2 pieces	
Shape	ROUND	Size	12mm	
Flavor		Imprint Code		
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:50269-525- 00	2 in 1 BOX	01/15/2012		
1		20 in 1 PACKET; Type 0: Not a Combination Product			

Marketing Information				
Marketing Application Number or Monograph Category Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M009	01/15/2012		

Labeler - JC World Bell Wholesale Co., Inc. (805257581)

Registrant - JC World Bell Wholesale Co., Inc. (805257581)

Establishment				
Name	Address	ID/FEI	Business Operations	
JC World Bell Wholesale Co., Inc.		805257581	repack(50269-525)	

Revised: 9/2025 JC World Bell Wholesale Co., Inc.