

# **G-DOLOGEN- acetaminophen, dextbrompheniramine maleate tablet**

## **Kramer Novis**

-----

### **G-DOLOGEN® for adults**

#### ***Drug Facts***

##### ***Active ingredients (in each tablet)***

Acetaminophen 500 mg  
Dextbrompheniramine maleate 2mg

##### ***Purpose***

Pain reliever/fever reducer  
Antihistamine

##### ***Uses***

- for the temporary relief of minor aches and pains associated with
  - headache • the common cold • toothache • muscular aches • backache •
- premenstrual and menstrual cramps • the minor pain from arthritis and reduces fever.
- for the temporary relief of • runny nose • sneezing • itching of the nose or throat • itchy, watery eyes due to hay fever.

##### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 tablets in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are allergic to acetaminophen or any of the inactive ingredients in the product • this product unless directed by a doctor, if you have a breathing problem such as emphysema or chronic bronchitis, or if you have glaucoma or difficulty in urination due to enlargement of the prostate gland • if you are taking sedatives or tranquilizers, without first consulting your doctor.

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product** • may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect.

Avoid alcoholic beverages while taking this product.

• use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions:***

adults and children 12 years of age and older	take 1 tablet every 4 to 6 hours as needed, do not exceed 6 tablets in 24 hours, or as directed by a doctor.
children under 12 years of age	consult a doctor

***Other information:***

• tamper evident feature: do not use if security seal is torn, broken, or missing • store at controlled room temperature: 15°- 30°C (59° - 86°F) • avoid excessive heat or humidity

***Inactive ingredients:***

magnesium stearate, microcrystalline cellulose, sodium starch glycolate, stearic acid

***Questions or comments?***

**1-787-767-2072**

Call weekdays from 8:00 am to 4.00 pm AST.

**You may also report serious side effects to this phone number.**

Compare to Dologen®

Sugar & Dye Free

**Manufactured for:** Kramer Novis  
San Juan, PR 00917

Made in USA with imported ingredients

www.kramernovis.com

**Peel the label from the bottom right corner for Drug Facts (continued).**

## Packaging

**Drug Facts**

**Active Ingredients (in each tablet)**      **Purpose**

Acetaminophen 500 mg.....Pain reliever/fever reducer  
Dexbrompheniramine maleate 2mg.....Antihistamine

**Uses**

- for the temporary relief of minor aches and pains associated with
- headache • the common cold • toothache • muscular aches • backache • premenstrual and menstrual cramps
- the minor pain from arthritis and reduces fever.
- for the temporary relief of • runny nose • sneezing
- itching of the nose or throat • itchy, watery eyes due to hay fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 tablets in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

Peel the label from the bottom right corner for Drug Facts (continued).

**Manufactured for:** Kramer Novis  
San Juan, PR 00917  
Made in USA with imported ingredients  
www.kramernovis.com  
Rev. 08/25

NDC 52083-683-90

**G-DOLOGEN<sup>®</sup>**  
for adults  
Compare to Dologen<sup>®</sup>

**Acetaminophen,  
Dexbrompheniramine  
Maleate**

**Pain Reliever-  
Fever Reducer  
Antihistamine**

Sugar & Dye Free  
**Net Contents 90 Tablets**

**KramerNovis**  
Pharmaceuticals within reach of patients

## Drug Facts Label

**Drug Facts**

**Active ingredients (in each tablet)**      **Purpose**

Acetaminophen 500 mg.....Pain reliever/fever reducer  
Dexbrompheniramine maleate 2mg.....Antihistamine

**Uses**

- for the temporary relief of minor aches and pains associated with
- headache • the common cold • toothache • muscular aches • backache • premenstrual and menstrual cramps
- the minor pain from arthritis and reduces fever.
- for the temporary relief of • runny nose • sneezing
- itching of the nose or throat • itchy, watery eyes due to hay fever.

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take • more than 6 tablets in 24 hours, which is the maximum daily amount

- with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

**Drug Facts (continued)**

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist • if you are allergic to acetaminophen or any of the inactive ingredients in the product • this product unless directed by a doctor, if you have a breathing problem such as emphysema or chronic bronchitis, or if you have glaucoma or difficulty in urination due to enlargement of the prostate gland • if you are taking sedatives or tranquilizers, without first consulting your doctor.

**Ask a doctor before use if you have** liver disease

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin

**When using this product** • may cause drowsiness; alcohol, sedatives, and tranquilizers may increase the drowsiness effect.  
Avoid alcoholic beverages while taking this product.  
• use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if** • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • any new symptoms appear

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions:**

adults and children 12 years of age and older	take 1 tablet every 4 to 6 hours as needed, do not exceed 6 tablets in 24 hours, or as directed by a doctor.
children under 12 years of age	consult a doctor

**Other information:** • tamper evident feature: do not use if security seal is torn, broken, or missing • store at controlled room temperature: 15° - 30° C (59° - 86° F) • avoid excessive heat or humidity

**Inactive ingredients:** magnesium stearate, microcrystalline cellulose, sodium starch glycolate, stearic acid

**Questions or comments? 1-787-767-2072**  
Call weekdays from 8:00 am to 4:00 pm AST.  
**You may also report serious side effects to this phone number.**

# G-DOLOGEN

acetaminophen, dextbrompheniramine maleate tablet

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:52083-683
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg
DEXBROMPHENIRAMINE MALEATE (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg

## Inactive Ingredients

Ingredient Name	Strength
MAGNESIUM STEARATE (UNII: 70097M6I3O)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

## Product Characteristics

Color	white	Score	no score
Shape	CAPSULE	Size	19mm
Flavor		Imprint Code	D
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52083-683-90	90 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/15/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/15/2025	

**Labeler** - Kramer Novis (090158395)

**Registrant** - Kramer Novis (090158395)

