

GOOD SENSE IBUPROFEN CHILDRENS- ibuprofen suspension
Preferred Pharmaceuticals Inc.

Perrigo Children's Ibuprofen Oral Suspension Drug Facts

Active ingredient (in each 5 mL)

Ibuprofen 100 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Purposes

Pain reliever/fever reducer

Uses

temporarily:

- relieves minor aches and pains due to the common cold, flu, sore throat, headache and toothache
- reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more

than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease, or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- give with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint
 - vomits blood
 - has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain
 - trouble breathing
 - weakness in one part or side of body
 - slurred speech
 - leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area

- any new symptoms appear

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise use age.
- use only enclosed dosing cup. Do not use any other dosing device.
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**
- replace original bottle cap to maintain child resistance
- wash dosage cup after each use

Dosing Chart

Weight (lb)	Age (yr)	Dose (mL)**
under 24 lbs	under 2 years	ask a doctor
24-35 lbs	2-3 years	5 mL
36-47 lbs	4-5 years	7.5 mL
48-59 lbs	6-8 years	10 mL
60-71 lbs	9-10 years	12.5 mL
72-95 lbs	11 years	15 mL

**or as directed by a doctor

Other information

- **each 5 mL contains:** sodium 2 mg
- **do not use if printed neckband is broken or missing**
- store at 20-25°C (68-77°F)
- do not freeze

Inactive ingredients

anhydrous citric acid, FD&C red #40, glycerin, high fructose corn syrup, hypromellose, natural and artificial bubblegum flavors, polysorbate 80, purified water, sodium benzoate, sorbitol solution, xanthan gum

Questions or comments?
1-800-719-9260

Principal Display Panel

GOODSENSE®
68788-4036-1
For Ages 2 to 11 Years
Children’s Ibuprofen
Oral Suspension
100 mg per 5 mL
Pain Reliever
Fever Reducer
(NSAID)
Lasts Up To 8 Hours
Alcohol Free
Bubble Gum Flavor
Compare to active ingredient of Children’s Motrin®
8 FL OZ (240 mL)

Childrens
Ibuprofen Oral
Suspension

Generic for Childrens Motrin

Active ingredient (in each 5mL): Ibuprofen,
USP 100mg (NSAID)...pain reliever/fever
reducer

Pkg Size: Exp Date: ####/####
Lot#: Batch#:

Ins:
Mfg: L. Perrigo Company
Prod#:

Warning
Do not use if printed neckband is broken or missing.
Store at 20°- 25°C (68°- 77°F). Allergy alert: Ibuprofen
may cause a severe allergic reaction, especially in people
allergic to aspirin. Stomach bleeding warning: this
product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding.
Sore throat warning: severe or persistent sore throat or
sore throat accompanied by a high fever, headache,
nausea, and vomiting may be serious. Consult doctor
promptly. Do not use more than 2 days or administer to
children under 3 years of age unless directed by doctor.
Read box for additional warnings, drug facts, directions,
and other information. Bubble Gum Flavor. Alcohol
Free.

Directions English

Take _____teaspoonful(s)
) every _____ hours.

CAUTION: Federal law PROHIBITS transfer of
this drug to any person other than the patient for
whom it was prescribed.

GTIN

SN #####
EXP #####

Instrucciones Espanol:
Toma _____cuchardita(s)
) cada _____ horas

Childrens Ibuprofen Oral
Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Childrens Ibuprofen Oral
Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Childrens Ibuprofen Oral
Suspension
Qty: Ins:
Insurance NDC:
Lot: Bat:

Childrens Ibuprofen Oral
Suspension
Qty: Ins:
Lot: Bat:
Prod# (NDC):

Log

Chart

Billing

Patient

GOOD SENSE IBUPROFEN CHILDRENS ibuprofen suspension			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:68788-4036(NDC:0113-0166)

Route of Administration		ORAL		
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)		IBUPROFEN	100 mg in 5 mL	
Inactive Ingredients				
Ingredient Name		Strength		
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
POLYSORBATE 80 (UNII: 6OZP39ZG8H)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
SORBITOL SOLUTION (UNII: 8KW3E207O2)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color	PINK (light)	Score		
Shape		Size		
Flavor	BUBBLE GUM	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68788-4036-1	1 in 1 CARTON	12/23/2025	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA074937		12/23/2025	

Labeler - Preferred Pharmaceuticals Inc. (791119022)

Registrant - Preferred Pharmaceuticals Inc. (791119022)

Establishment			
Name	Address	ID/FEI	Business Operations
Preferred Pharmaceuticals Inc.		791119022	RELABEL(68788-4036)

Revised: 12/2025

Preferred Pharmaceuticals Inc.