

ANTI DIARRHEAL- loperamide hcl tablet, film coated
Lidl US LLC

Lidl US LLC Anti-Diarrheal Drug Facts

Active ingredient (in each caplet)

Loperamide HCl 2 mg

Purpose

Anti-diarrheal

Use

controls symptoms of diarrhea, including Travelers' Diarrhea

Warnings

Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl

Heart alert: Taking more than directed can cause serious heart problems or death

Do not use

if you have bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool
- a history of liver disease

Ask a doctor or pharmacist before use if you are

taking antibiotics

When using this product

tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery.

Stop use and ask a doctor if

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

Other information

- store at 20°-25°C (68°-77°F)
- see end panel for lot number and expiration date

Inactive ingredients

anhydrous lactose, carnauba wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to the active ingredient in Imodium[®] A-D

LOPERAMIDE HYDROCHLORIDE TABLETS, 2 mg

Anti-Diarrheal

Actual Size

Controls the Symptoms of Diarrhea

12 caplets*

*capsule-shaped tablets

anti-diarrheal

SEE NEW WARNING AND DIRECTIONS

DISTRIBUTED BY: LIDL US, LLC
3500 S. CLARK STREET, ARLINGTON, VA 22202

follow us on



GLUTEN FREE



NDC 71141-001-53

Compare to the
active ingredient in
Imodium® A-D**

LOPERAMIDE
HYDROCHLORIDE
TABLETS, 2 mg



Actual Size

Anti-Diarrheal

Controls the Symptoms
of Diarrhea

12
caplets*

*capsule-shaped tablets
anti-diarrheal

SEE NEW WARNING
AND DIRECTIONS



22453 F3 C2

**This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Imodium® A-D.

Drug Facts Active ingredient (in each caplet) Loperamide HCl 2 mg.....Anti-diarrheal Purpose controls symptoms of diarrhea, including Travelers' Diarrhea	
Warnings Allergy alert: Do not use if you have ever had a rash or other allergic reaction to loperamide HCl Heart alert: Taking more than directed can cause serious heart problems or death Do not use if you have bloody or black stool Ask a doctor before use if you have ■ fever ■ mucus in the stool ■ a history of liver disease Ask a doctor or pharmacist before use if you are taking antibiotics When using this product tiredness, drowsiness or dizziness may occur. Be careful when driving or operating machinery. Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days ■ you get abdominal swelling or bulging. These may be signs of a serious condition. If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). ▶	
Drug Facts (continued) Directions ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.	
adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34 to 47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use
Other information ■ store at 20°-25°C (68°-77°F) ■ see end panel for lot number and expiration date ▼	



Drug Facts (continued) Inactive ingredients anhydrous lactose, calcium wax, D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch ▶	Drug Facts (continued) Questions or comments? 1-800-719-9260 DO NOT USE IF BLISTER UNIT IS BROKEN OR TORN
---	---

ANTI DIARRHEAL			
loperamide hcl tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71141-001
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	LOPERAMIDE HYDROCHLORIDE (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4I)	LOPERAMIDE HYDROCHLORIDE	2 mg
Inactive Ingredients			

Ingredient Name	Strength
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	

Product Characteristics

Color	GREEN	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	L2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71141-001-53	12 in 1 CARTON	03/15/2017	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA075232	03/15/2017	

Labeler - Lidl US LLC (079389709)