

**MUCUS RELIEF CHILDRENS- dextromethorphan hbr, guaifenesin,
phenylephrine hcl solution
TOPCO ASSOCIATES LLC**

TopCare 44-057

Active ingredients (in each 5 mL)

Dextromethorphan HBr 5 mg
Guaifenesin 100 mg
Phenylephrine HCl 2.5 mg

Purpose

Cough suppressant
Expectorant
Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - nasal congestion due to a cold
 - the intensity of coughing
 - stuffy nose
 - the impulse to cough to help the child get to sleep

Warnings

Do not use

in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- diabetes
- heart disease
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- **each 5 mL contains:** sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose, xanthan gum

Questions or comments?

1-888-423-0139

Principal display panel

+TopCare®
health

NDC 76162-057-88

COMPARE TO MUCINEX®
CHILDREN'S MULTI-SYMPTOM COLD
ACTIVE INGREDIENTS*

children's

**Multi-Symptom
Cold
Mucus
Relief**

DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
GUAIFENESIN - EXPECTORANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

ORAL SOLUTION

- **Controls Cough**
- **Relieves Chest Congestion**
- **Breaks Up Mucus**
- **Relieves
Stuffy Nose**

**Ages 4
to
11 Years**

6.8 FL OZ (201 mL)

BERRY
FLAVOR

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ITASCA, IL 60143
©TOPCO LNKA0923
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com

*This product is not manufactured or distributed by
RB Health (US) LLC, owner of the registered trademark
Mucinex® Children's Multi-Symptom Cold.
50844 ORG022305788

**PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org**

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each 5 mL)	Purpose
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 2.5 mg	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - nasal congestion due to a cold
 - the intensity of coughing
 - stuffy nose
 - the impulse to cough to help the child get to sleep

Warnings

Do not use in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- glaucoma
- heart disease
- thyroid disease
- high blood pressure
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nausea, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- do not take more than 6 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- dose as follows or as directed by a doctor

Age	Dose
children 6 to under 12 years	10 mL every 4 hours
children 4 to under 6 years	5 mL every 4 hours
children under 4 years	do not use

Other information

- each 5 mL contains: sodium 3 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients anhydrous citric acid, FD&C red #40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sorbitol, sucralose, xanthan gum

Questions or comments? 1-888-423-0139

TopCare
health

NDC 76162-057-88

COMPARE TO MUCINEX®
CHILDREN'S MULTI-SYMPOM COLD
ACTIVE INGREDIENTS*

children's Multi-Symptom Cold Mucus Relief

TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

DISTRIBUTED BY TOPCO ASSOCIATES LLC
ITASCA, IL 60143
©TOPCO LNKAD923
QUESTIONS? 1-888-423-0139
topcare@topco.com
www.topcarebrand.com



Scan here for more
information or call 1-888-423-0139

QUALITY GUARANTEED

*This product is not manufactured or distributed by
RB Health (US) LLC, owner of the registered trademark
Mucinex® Children's Multi-Symptom Cold.
50844 ORG022305788

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

B-1910-057-88
ORG022305788

TopCare
health

children's Multi-Symptom Cold Mucus Relief

6.8 FL OZ (201 mL)

TopCare
health

NDC 76162-057-88

COMPARE TO MUCINEX®
CHILDREN'S MULTI-SYMPOM COLD
ACTIVE INGREDIENTS*

children's Multi-Symptom Cold Mucus Relief

DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
GUAIFENESIN - EXPECTORANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

ORAL SOLUTION

- Controls Cough
- Relieves Chest Congestion
- Breaks Up Mucus
- Relieves Stuffy Nose

Ages 4
to
11 Years



BERRY
FLAVOR

6.8 FL OZ (201 mL)

TopCare
health

NDC 76162-057-88

COMPARE TO MUCINEX®
CHILDREN'S MULTI-SYMPOM COLD
ACTIVE INGREDIENTS*

children's Multi-Symptom Cold Mucus Relief

DEXTROMETHORPHAN HBr - COUGH SUPPRESSANT
GUAIFENESIN - EXPECTORANT
PHENYLEPHRINE HCl - NASAL DECONGESTANT

ORAL SOLUTION

- Controls Cough
- Relieves Chest Congestion
- Breaks Up Mucus
- Relieves Stuffy Nose

Ages 4
to
11 Years



BERRY
FLAVOR

6.8 FL OZ (201 mL)



0 36800 00340 8

No Print/No Varnish
Lot & EXP

TopCare 44-057

MUCUS RELIEF CHILDRENS

dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:76162-057

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	BERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:76162-057-88	1 in 1 CARTON	03/07/2024	
1		201 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/07/2024	

Labeler - TOPCO ASSOCIATES LLC (006935977)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(76162-057) , pack(76162-057)

Revised: 3/2026

TOPCO ASSOCIATES LLC