ULTRA-GUARD ACTIVATOR- ultra-guard activator solution IBA

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

ULTRA-GUARD SANITIZING BARRIER TEAT DIP PROVIDES A PROTECTIVE BARRIER OVER THE TEAT END UNTIL THE NEXT MILKING CONTAINS EMOLLIENTS FOR EXCELLENT SKIN CONDITIONING

ACTIVE INGREDIENT

Organic Acids......2.85%

INERT INGREDIENTS

Contains A Special Emollient System

INDICATIONS AND USAGE

GENERAL RECOMMENDATIONS

To make a ready-to-use solution of Ultra-Guard Sanitizing Barrier Teat dip, add equal volumes of Ultra-Guard Activator and Ultra-Guard Base into a clean dip cup/container and mix until the color is uniform throughout. Do not dilute. Mix only the amount needed for one milking of the herd. Discard any unused teat dip. Do not mix with any other products.

RECOMMENDED PROCEDURES FOR TEAT SANITATION

POST-MILKING (POST-DIPPING)

Immediately after milking, dip all teats full length with READY-TO-USE ULTRA-GUARD SANITIZING BARRIER TEAT DIP. Ensure good coverage of all sides of the teats and teat orifices. Allow to dry. Do not wipe. Do not turn cows out in freezing weather until teat dip is completely dry.

NOTE: Before milking, use a compatible teat dip or udder wash to thoroughly clean the teats.

WARNING

Do not dilute this product.

Do not return teat dip left in the teat dipper to storage container.

Ensure that dipped teats are dry before exposing the animals to weather conditions which may cause damage to the teats.

Store at room temperature. Always store away from direct sunlight.

Protect this product from freezing. However, freezing will not affect ULTRA-GUARD ACTIVATOR provided that the contents are thawed completely and agitated thoroughly

before mixing base and activator together.


CAUTION: KEEP OUT OF REACH OF CHILDREN.
AVOID CONTACT WITH FOOD. DO NOT TAKE INTERNALLY.


FIRST AID PROCEDURES

EYE CONTACT: Rinse cautiously with fresh, tepid water for several minutes. Remove contact lenses if present and easy to do. Continue rinsing for 15 minutes. Get immediate medical advice/attention. SKIN CONTACT: For RTU products wash any sensitive areas. INGESTION: Do not induce vomiting. Never give anything by mouth to an unconscious person. Drink large amounts of water. If vomiting occurs, keep airways open. Keep head lower than the hips to prevent aspiration into the lungs. Get immediate medical advice/attention.

ALWAYS SEEK IMMEDIATE MEDICAL ATTENTION
FOR MEDICAL EMERGENCIES CALL 1-800-424-9300

ULTRA-GUARD ACTIVATOR





ULTRA-GUARD ACTIVATOR

ultra-guard activator solution

Product Information Product Type OTC ANIMAL DRUG Item Code (Source) NDC:29983-0087 Route of Administration TOPICAL

Active Ingredient/Active Moiety Ingredient Name Basis of Strength LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT) LACTIC ACID (28.5 g in 1 L

Inactive Ingredients			
Strength			

Product Characteristics				
Color	orange	Score		

Shape	Size	
Flavor	Imprint Code	
Contains		

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:29983-0087-4	18.9 L in 1 DRUM					
2	NDC:29983-0087-6	56.7 L in 1 DRUM					
3	NDC:29983-0087-8	207.9 L in 1 DRUM					
4	NDC:29983-0087-9	1039.5 L in 1 DRUM					

Marketing Information					
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
unapproved drug other		04/15/2022			
outei					

Labeler - IBA (019494160)

Revised: 12/2024 IBA