# PAIN RELIEVER AND FEVER REDUCER- acetaminophen tablet, film coated Spirit Pharmaceuticals LLC

-----

Pain Reliever and Fever Reducer

## **Drug Facts**

## Active Ingredient (in each tablet)

Acetaminophen 500 mg

## **Purpose**

Pain reliever/fever reducer

#### Uses

• temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis • toothache • muscular aches • premenstrual and menstrual cramps • temporarily reduces fever

## Warning

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

more than 4,000 mg of acetaminophen in 24 hours
 with other drugs containing acetaminophen
 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash. If a skin reaction occurs, stop use and seek medical help right away.

#### Do not use

• with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are allergic to acetaminophen or any of the inactive ingredients in this product

# Ask a doctor before use if you have

liver disease

# Ask a doctor or pharmacist before use if you are if you are

taking the blood thinning drug warfarin

## Stop use and ask a doctor if

• pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

## If pregnant or breast-feeding,

ask a health professional before use.

## Keep out of reach of children.

## Overdose warning:

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions**

do not take more than directed (see overdose warning)

adults & children 12 years & over	<ul> <li>take 2 tablets every 6 hours while symptoms last</li> <li>do not take more than 6 tablets in 24 hours, unless directed by a doctor</li> <li>do not use for more than 10 days unless directed by a doctor</li> </ul>
children under 12 years	ask a doctor

#### Other information

• store between 20-25°C (68-77°F)

# Inactive ingredients

Hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl pyrrolidone, pregelatinized starch, stearic acid, talc, titanium dioxide

# Inactive ingredients

1-888-333-9792

# Distributed by:

Cabinet Health P.B.C.

# **Principal Display Panel**





# CABINET:

# Pain Reliever & Fever Reducer Refill

COMPARE TO THE ACTIVE INGREDIENT IN:

Extra Strength Tylenol®\*

# Acetaminophen 500 mg

### **RELIEVES:**

HEADACHE · FEVER · MUSCLE ACHES AND PAINS MENSTRUAL CRAMPS



#### TO OPEN:

- 1. POSITION THUMBS INSIDE THE SHORT FLANGE OVER THE LOCKS.
- 2. GRIP WHILE PIVOTING THUMBS OUTWARD TO OPEN THE LOCKS.

Tamper evident: do not use if pouch is open

## **Drug Facts**

## Active Ingredient (in each tablet)

**Purpose** 

Acetaminophen 500 mg......Pain reliever/fever reducer

**Uses** • temporarily relieves minor aches and pains due to: • the common cold • headache • backache • minor pain of arthritis

- toothache muscular aches premenstrual and menstrual cramps
- temporarily reduces fever

#### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take . more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away

Do not use • with any other drug containing acetaminophen (prescription or non prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

• if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

## **Directions** • do not take more than directed (see overdose warning)

adu	lts -	&	ch	ildr	en
121	/ea	rs	&	ove	er

• take 2 tablets every 6 hours while symptoms last . do not take more than 6 tablets in 24 hours, unless directed by a doctor . do not use for more than 10 days unless directed by a doctor

children under 12 vears

ask a doctor

## **Drug Facts** (continued)

#### Other Information

• store between 20-25°C (68-77°F)

**Inactive ingredients** Hypermellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinylpyrrolidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments? 1-888-333-9792

# Single-use plastic is a headache of the past.

- \* This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Extra Strength Tylenol®
- † Less than the limit of detection and consistent with gluten-free diet labeling per FDA

Distributed by: Cabinet Health P.B.C. Question/Comments? 1-888-333-9792 www.cabinethealth.com

Lot / Expiration ▼









## PAIN RELIEVER AND FEVER REDUCER

acetaminophen tablet, film coated

B			
Prodi	CT	Inform	ation
I I Out			ation

Route of Administration ORAL

## **Active Ingredient/Active Moiety**

Ingredient Name	<b>Basis of Strength</b>	Strength
-----------------	--------------------------	----------

ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D) ACETAMINOPHEN 500 mg

## **Inactive Ingredients**

Ingredient Name	Strength
HYPROMELLOSES (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

## **Product Characteristics**

Color	white	Score	no score
Shape	CAPSULE	Size	18mm
Flavor		Imprint Code	S500
Contains			

## **Packaging**

# Item Code	Package Description	Marketing Start Date	Marketing End Date
<b>1</b> NDC:68210-4235-1	120 in 1 POUCH; Type 0: Not a Combination Product	11/17/2022	

# **Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	11/15/2022	
OTC Monograph Drug	M013	11/15/2022	

# Labeler - Spirit Pharmaceuticals LLC (179621011)

Revised: 12/2023 Spirit Pharmaceuticals LLC