

BLUEDOT ONE-STEP CORN REMOVER STRIPS- salicylic acid plaster
CREST MEDICAL LIMITED

BLUEDOT™ One-Step Corn Remover Strips

Drug Facts

Active ingredient (in each dosage unit)

Salicylic acid 40% in a plaster vehicle

Purposes

Corn and calluses removal

Uses

- For the removal of corns and calluses
- Relieves pain by removing corns and calluses
- Adhesive bandage holds the medicated pad in position

Warnings

For external use only

Do not use

- If you are a diabetic
- If you have poor blood circulation
- On irritated skin or any area that is infected or reddened

Stop use and ask a doctor if

Discomfort lasts

Keep out of reach of children

If swallowed, get medical help or contact a Poison Control Center right away

Directions

- Wash affected area and dry thoroughly
- Apply medicated bandage after 48 hours, remove medicated bandage
- Repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- May soak corn in warm water for 5 minutes to assist in removal

Other information

Store between 0° to 40°C (32° to 104°F)

Inactive ingredients

Acrylic adhesive, Titanium dioxide, Ferric hydroxide

Questions?

1-800-723-2465

- **Maximum Strength**
- **Medicated Pad**
- **Water Resistant**
- **Stays in place**

DISTRIBUTED BY:

Crest Medical Ltd.
3 Chesford Grange,
Warrington, Cheshire.
WA1 4RQ. UK.

www.crestmedical.co.uk

MADE IN BULGARIA

Packaging



BLUEDOT™

One-Step Corn Remover Strips

Salicylic Acid 40%

- Maximum Strength
- Medicated Pad
- Water Resistant
- Stays in place





Drug Facts

Active ingredient (in each dosage unit) Salicylic acid 40% in a plaster vehicle	Purposes Corn and calluses removal
---	--

Uses

- For the removal of corns and calluses
- Relieves pain by removing corns and calluses
- Adhesive bandage holds the medicated pad in position

Warnings

For external use only

Do not use

- If you are a diabetic
- If you have poor blood circulation
- On irritated skin or any area that is infected or reddened

Stop use and ask a doctor if
Discomfort lasts

Keep out of reach of children
If swallowed, get medical help or contact a Poison Control Center right away

Directions

- Wash affected area and dry thoroughly
- Apply medicated bandage after 48 hours, remove medicated bandage
- Repeat procedure every 48 hours as needed for up to 14 days (until corn is removed)
- May soak corn in warm water for 5 minutes to assist in removal

Other information
Store between 0° to 40°C (32° to 104°F)

Inactive ingredients
Acrylic adhesive, Titanium dioxide, Ferric hydroxide

Questions?
1-800-723-2465

Text prepared 8/24 ITEM CM997620

DISTRIBUTED BY:
Crest Medical Ltd.
3 Chesford Grange,
Warrington, Cheshire.
WA1 4RQ. UK.
www.crestmedical.co.uk
MADE IN BULGARIA



5 039139 012535 >

LOT



BLUEDOT ONE-STEP CORN REMOVER STRIPS

salicylic acid plaster

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84713-101
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	40 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
FERRIC HYDROXIDE (UNII: 2UA751211N)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84713-101-06	1 in 1 BOX	09/15/2024	
1		6 in 1 BAG		
1		0.0044 g in 1 PATCH; Type 4: Device Coated/Impregnated/Otherwise Combined with Drug		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M030	09/15/2024	

Labeler - CREST MEDICAL LIMITED (238858539)

Establishment

Name	Address	ID/FEI	Business Operations
KRE EOOD		565504983	manufacture(84713-101) , label(84713-101) , pack(84713-101)

Revised: 9/2024

CREST MEDICAL LIMITED