

**LAXATIVE MAXIMUM STRENGTH- sennosides tablet
MEIJER, INC.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient (in each tablet)

Sennosides USP, 25 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have

- noticed a sudden change in bowel habits that persists over a period of 2 weeks

Ask a doctor or pharmacist before use if you

- are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product

- do not use for a period longer than 1 week

Stop use and ask a doctor if

- rectal bleeding or failure to have a bowel movement occur after use of a laxative.

These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- swallow tablet(s) with a glass of water
- swallow tablet(s) whole, do not crush, break, or chew

adults and children 12 years of age and older	take 2 tablets once or twice daily
children 6 to under 12 years of age	take 1 tablet once or twice daily
children under 6 years of age	ask a doctor

Other information

- **each tablet contains:** calcium 40 mg
- very low sodium
- store between 20-25°C (68-77°F)

Inactive ingredients

acacia, calcium phosphate dihydrate*, calcium sulfate anhydrous*, carnauba wax, corn starch*, croscarmellose sodium, FD&C blue #1, iron oxide*, iron oxide black, iron oxide yellow*, kaolin*, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone*, shellac, silicon dioxide*, stearic acid*, sucrose, talc, titanium dioxide,

*contains one or more of these ingredients

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

†Compare to the active ingredient in ex-lax® Maximum Strength

MAXIMUM STRENGTH

laxative

Sennosides USP | 25 mg

Stimulant laxative

Gentle, Dependable Overnight Relief

Tablets

†This product is not manufactured or distributed by GlaxoSmithKline plc, distributor of ex-lax®.

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOW SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

DIST BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

www.meijer.com

Product Label



NDC 41250-815-24

†Compare to the active ingredient in ex-lax® Maximum Strength

MAXIMUM STRENGTH laxative

Sennosides USP | 25 mg Stimulant laxative

Gentle, Dependable Overnight Relief

24 Tablets



Actual Size

TAMPER EVIDENT: DO NOT USE IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING. KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:
Exp. Date:

Drug Facts (continued)

Active ingredient (in each tablet)
Sennosides USP, 25 mg.....Stimulant laxative

Purpose

Uses
 ■ relieves occasional constipation (irregularly)
 ■ generally produces bowel movement in 6 to 12 hours

Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have
 ■ noticed a sudden change in bowel habits that persists over a period of 2 weeks

Drug Facts (continued)

Ask a doctor or pharmacist before use if you
 ■ are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product
 ■ do not use for a period longer than 1 week

Stop use and ask a doctor if
 ■ rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole, do not crush, break, or chew

adults and children 12 years of age and older
 take 2 tablets once or twice daily

Drug Facts (continued)

Drug Facts (continued)

Ask a doctor or pharmacist before use if you
 ■ are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product
 ■ do not use for a period longer than 1 week

Stop use and ask a doctor if
 ■ rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole, do not crush, break, or chew

adults and children 12 years of age and older
 take 2 tablets once or twice daily

Questions or comments?
 Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Other information
 ■ each tablet contains: calcium 40 mg
 ■ very low sodium
 ■ store between 20-25°C (68-77°F)

Inactive ingredients acacia, calcium phosphate dihydrate, calcium sulfate anhydrous, carnauba wax, colloidal silicon dioxide, corn starch, croscarmellose sodium, FD&C blue #1 aluminum lake, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, povidone, shellac, stearic acid, sugar, talc, titanium dioxide

Drug Facts (continued)

children 6 to under 12 years of age	take 1 tablet once or twice daily
children under 6 years of age	ask a doctor

†This product is not manufactured or distributed by GlaxoSmithKline plc. distributor of ex-lax®.

DIST. BY MEIJER DISTRIBUTION, INC.
 GRAND RAPIDS, MI 49544
 www.meijer.com



PLD-C16J
 FC005586

MEIJER Maximum Strength Laxative

LAXATIVE MAXIMUM STRENGTH

sennosides tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:41250-815
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	25 mg

Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM PHOSPHATE, DIBASIC, DIHYDRATE (UNII: O7TSZ97GEP)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
KAOLIN (UNII: 24H4NWX5CO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	TCL083
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-815-48	48 in 1 CARTON	01/31/2019	01/31/2025
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:41250-815-24	24 in 1 CARTON	01/31/2019	01/31/2025
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	01/31/2019	01/31/2025

Labeler - MEIJER, INC. (006959555)

Revised: 1/2023

MEIJER, INC.