

BROAD SPECTRUM SPF 50 KIDS PREFERRED PLUS PHARMACY- avobenzone, homosalate, octisalate, octocrylene, oxybenzone lotion
Kinray Inc.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredients

Avobenzone 3.0%, Homosalate 13.0%, Octisalate 5.0%, Octocrylene 7.0%, Oxybenzone 4.0%

Uses

Helps prevent sunburn.

Warnings

For external use only. Do not use on damaged or broken skin. **When using this product** keep out of eyes. Rinse with water to remove. **Stop use and ask a doctor** if rash occurs.

Directions

Apply liberally 15 minutes before sun exposure. Reapply: after 80 minutes of swimming or sweating, immediately after towel drying, at least every 2 hours.

Sun Protection Measures: spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including: limit time in the sun, specially from 10am - 2pm, wear long-sleeve shirts, pants, hats, and sunglasses.

Children under 6 months: ask a doctor.

Inactive ingredients

Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Carbomer, Dimethicone, Disodium EDTA, Fragrance, Methylparaben, Polyglyceryl-3 Distearate, Propylparaben, Sorbitan Isostearate, Sorbitol, Stearic Acid, Tocopherol, Triethanolamine, VP/Eicosene Copolymer, Water.

Purpose

Sunscreen.

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away.

Preferred Plus Pharmacy

Broad Spectrum SPF 50

Kids Moisturizing Sunscreen Lotion

Water Resistant (80 Minutes)



Drug Facts

Active ingredients	Purpose
Avobenzone 3.0%, Homosalate 13.0%, Octisalate 5.0%, Octocrylene 7.0%, Oxybenzone 4.0%	Sunscreen

Uses • helps prevent sunburn

Warnings
 For external use only
 Do not use on damaged or broken skin
 When using this product keep out of eyes. Rinse with water to remove.
 Stop use and ask a doctor if rash occurs
 Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions • apply liberally 15 minutes before sun exposure
 • reapply: • after 80 minutes of swimming or sweating
 • immediately after towel drying • at least every 2 hours
 • Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 • limit time in the sun, especially from 10 a.m. – 2 p.m.
 • wear long-sleeve shirts, pants, hats, and sunglasses
 • children under 6 months: Ask a doctor

Inactive ingredients Aluminum Starch Octenylsuccinate, Benzyl Alcohol, Carbomer, Dimethicone, Disodium EDTA, Fragrance, Methylparaben, Polyglyceryl-3 Distearate, Propylparaben, Sorbitan Isostearate, Sorbitol, Stearic Acid, Tocopherol, Triethanolamine, VP/Eicosene Copolymer, Water.

Other information
 • protect this product from excessive heat and direct sun

ALL PREFERRED PLUS PHARMACY PRODUCTS ARE SATISFACTION GUARANTEED OR YOUR MONEY BACK FROM RETAILER WHERE PURCHASED.

Distributed by: KINRAY INC.
 152-35 Tenth Avenue
 Whitestone, NY 11357

© 2011 Kinray Inc., All Rights Reserved, PREFERRED PLUS, PREFERRED PLUS +++ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinray Inc. All other marks are property of their respective owners.



7 27510 00130 2

BROAD SPECTRUM SPF 50 KIDS PREFERRED PLUS PHARMACY			
avobenzone, homosalate, octisalate, octocrylene, oxybenzone lotion			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61715-028
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			

Ingredient Name	Basis of Strength	Strength
AVOBENZONE (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	3 g in 100 mL
HOMOSALATE (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	13 g in 100 mL
OCTISALATE (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	5 g in 100 mL
OCTOCRYLENE (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	7 g in 100 mL
OXYBENZONE (UNII: 95OOS7VE0Y) (OXYBENZONE - UNII:95OOS7VE0Y)	OXYBENZONE	4 g in 100 mL

Inactive Ingredients

Ingredient Name	Strength
ALUMINUM STARCH OCTENYLSUCCINATE (UNII: I9PJ0O6294)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
CARBOMER 940 (UNII: 4Q93RCW27E)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
METHYL PARABEN (UNII: A2I8C7HI9T)	
POLYGLYCERYL-3 DISTEARATE (UNII: ZH1LK470XV)	
PROPYL PARABEN (UNII: Z8IX2SC1OH)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
SORBITOL (UNII: 506T60A25R)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TOCOPHEROL (UNII: R0ZB2556P8)	
TRIETHANOLAMINE BENZOATE (UNII: M3EN4GC19W)	
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61715-028-08	237 mL in 1 BOTTLE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	03/15/2013	

Labeler - Kinray Inc. (012574513)

Revised: 5/2013

Kinray Inc.