TOPCARE FLU AND SEVERE COLD AND COUGH DAYTIME- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder, for solution

Topco Associates LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Topco Associates LLC. Flu & Severe Cold & Cough Drug Facts

Active ingredients (in each packet)

Acetaminophen 650 mg Dextromethorphan HBr 20 mg

Phenylephrine HCl 10 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold:
- minor aches and pains
- minor sore throat pain
- headache
- nasal and sinus congestion
- cough due to minor throat and bronchial irritation
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- in a child under 12 years of age
- if you have ever had an allergic reaction to this product or any of its ingredients
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occurs
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- pain, cough or nasal congestion gets worse or lasts more than 7 days
- cough comes back or occurs with rash or headache that lasts. These could be

signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not use more than directed (see overdose warning)
- take every 4 hours, while symptoms persist. Do not take more than 5 packets in 24 hours unless directed by a doctor.

Age	Dose		
adults and children 12 years of age and over	one packet		
children under 12 years of age	do not use		

- dissolve contents of one packet into 8 oz. hot water: sip while hot. Consume entire drink within 10-15 minutes.
- if using a microwave, add contents of one packet to 8 oz. of cool water: stir briskly before and after heating. Do not overheat.

Other information

- each packet contains: potassium 5 mg and sodium 43 mg
- **phenylketonurics:** contains phenylalanine 17 mg per packet
- store at 20-25°C (68-77°F). Protect product from heat and moisture.

Inactive ingredients

acesulfame potassium, anhydrous citric acid, aspartame, colloidal silicon dioxide, FD&C blue #1, FD&C red #40, flavor, maltodextrin, pregelatinized starch, sodium citrate, sucrose, tribasic calcium phosphate

Questions or comments?

1-888-423-0139

Package/Label Principal Display Panel

TopCare_® health

DAYTIME

COMPARE TO THERAFLU[®] DAYTIME SEVERE COLD & COUGH ACTIVE INGREDIENTS Flu & Severe Cold & Cough PAIN RELIEVER – FEVER REDUCER (ACETAMINOPHEN) COUGH SUPPRESSANT (DEXTROMETHORPHAN HBr) NASAL DECONGESTANT (PHENYLEPHRINE HCI) NON-DROWSY

FOR SOOTHING RELIEF OF:

- Nasal & Sinus Congestion
- Sore Throat Pain
- Headache
- Body Ache
- Fever
- Cough

6 PACKETS

BERRY INFUSED WITH MENTHOL FLAVOR



TOPCARE FLU AND SEVERE COLD AND COUGH DAYTIME

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride powder, for solution

Product Information						
Product Type	HUMAN OTC DRUG	Item Code (Source)		NDC:3680	NDC:36800-096	
Route of Administration	ORAL					
Active Ingredient/Active	Majaty					
Active ingredient/Active	molecy					
Ingredient Name		Basis of Str	ength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)			ACETAMINOPHEN		650 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)		DEXTROMETHORPHAN HYDROBROMIDE		20 mg		
PHENYLEPHRINE HYDROCHLORI UNII:1WS297W6MV)	DE (UNII: 04JA59TNSJ) (PHE	NYLEPHRINE -	PHENYLEPHRINE HYDROCHLORIDE		10 mg	

		Ingredient Name				Strength	
ACESULFAME P	OTAS	SIUM (UNII: 230V73Q5G9)					
		ACID (UNII: XF417D3PSL)					
ASPARTAME (UN							
SILICON DIOXID	DE (UN	II: ETJ7Z6XBU4)					
FD&C BLUE NO	. 1 (UI	NII: H3R47K3TBD)					
FD&C RED NO.	40 (UI	NII: WZB9127XOA)					
MALTODEXTRIN	I (UNII:	7CVR7L4A2D)					
SODIUM CITRAT	FE, UN	SPECIFIED FORM (UNII: 1Q73Q2JULR)					
SUCROSE (UNII:	C151H	l8M554)					
TRIBASIC CALC	IUM P	HOSPHATE (UNII: 91D9GV0Z28)					
Product Cha				_			
				Score			
Shape	e				Size		
	BERRY,	MENTHOL			Imprint Code		
Contains							
Packaging							
# Item Code	e	Package Description		Marketing Start Date	M	larketing End Date	
NDC-26900.00		in 1 CARTON; Type 0: Not a Combination oduct	01	/17/2012			
1 NDC:36800-09 91							
91	g In	formation					
	-	formation Application Number or Monograp Citation	h	Marketing Sta Date	rt	Marketing En Date	

Labeler - Topco Associates LLC (006935977)

Revised: 10/2022

Topco Associates LLC