ACYCLOVIR- acyclovir ointment OASIS TRADING

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Acyclovir

■ temporary relief of itching associated with minor skin irritations and rashes due to eczema, insect bites, poison ivy, poison oak, or poison sumac soaps detergents, cosmetics, jewelry, seborrheic dermatitis, psoriasis, external genital and anal itching

Keep out of reach of children

■ Adults and children 2 years of age and older: apply to the affected area no more than 3 to 4 times daily For external use only.

Do not use in the eyes by putting this product into the rectum by using fingers or any mechanical device or applicator.

Ask a doctor before use if you have a vaginal discharge, rectal bleeding, diaper rash.

When using this product consult a doctor before exceeding recommended dosage.

Stop use and ask a doctor if condition gets worse, condition persists for more than 7 days, condition clears up and occurs again with within a few days. Do not begin to use any other hydrocortisone product unless you have consulted a doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

cetostearyl alcohol, propylene glycol, purified water, sodium lauryl sulfate, white petrolatum For external use only

ACYCLOVIR

Drug Facts

Active Ingredients (in 1g)

Purpose

Acyclovir 50mg

Anti-itch

Uses

temporary relief of itching associated with minor skin irritations and rashes due to eczema, insect bites, poison ivy, poison oak, or poison sumac soaps detergents cosmetics, jewelry, seborrheic dermatitis, psoriasis

external genital and anal itching

Warnings/Precautions:

For external use only

Do not use in the eyes by putting this product into the rectum by using fingers or any mechanical device or applicator

Ask a doctor before use if you have a vaginal discharge rectal bleeding diaper rash

When using this product consult a doctor before exceeding recommended dosage

Stop use and ask a doctor if condition gets worse condition persists for more than 7 days

condition clears up and occurs again within a few days. Do not begin to use any other hydrocortisone product unless you have consulted a doctor.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Adults and children 2 years of age and older:

apply to affected area not more than 3 to 4 times daily.

Other Information

■ Store at room temperature

Inactive Ingredient

cetostearyl alcohol, propylene glycol, purified water, sodium lauryl sulfate, white petrolatum

Questions or comments?

Call weekdays from 9 a.m to 5 p.m EST at (201) 669-8405

Distributed By: P&K FRONTIER MARKETING CORP.

329 BROAD AVENUE # 2F, PALISADES PARK, NJ 07650, USA

Made in South Korea

ACYCLOVIR

acyclovir ointment

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72689-0018	
Route of Administration	TOPICAL			

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACYCLOVIR (UNII: X4HES1011F) (ACYCLOVIR - UNII:X4HES1011F)	ACYCLOVIR	50 mg in 1 g	

Inactive Ingredients		
Ingredient Name	Strength	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)		
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)		

Packaging				
ı	# Item Code	Package Description	Marketing Start Date	Marketing End Date
ı	1 NDC:72689-0018-1	5 g in 1 TUBE; Type 0: Not a Combination Product	11/16/2018	

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		11/15/2018		

Labeler - OASIS TRADING (689991468)

Registrant - OASIS TRADING (689991468)

Establishment			
Name	Address	ID/FEI	Business Operations
OASIS TRADING		689991468	manufacture(72689-0018)

Revised: 12/2018 OASIS TRADING