SUDOGEST- chlorpheniramine maleate, pseudoephedrine hcl tablet Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major 44-111

Active ingredients (in each tablet)

Chlorpheniramine maleate 4 mg Pseudoephedrine HCl 60 mg

Purpose

Antihistamine Nasal decongestant

Uses

- temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies:
 - runny nose
 - itchy, watery eyes
 - nasal congestion
 - sneezing
 - itching of the nose or throat
- temporarily relieves these symptoms due to the common cold:
 - runny nose
 - sneezing
 - sinus congestion and pressure

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- heart disease
- high blood pressure
- diabetes
- glaucoma
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- do not exceed recommended dose
- drowsiness may occur
- excitability may occur, especially in children
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.
- children under 12 years: do not use

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- store in a dry place and protect from light
- see end flap for expiration date and lot number

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, lactose anhydrous, magnesium stearate, microcrystalline cellulose, stearic acid

Questions or comments?

(800) 616-2471

Principal Display Panel

MAJOR®

NDC 0904-5351-24

Maximum Strength SudoGest Sinus & Allergy

Chlorpheniramine maleate 4 mg

Pseudoephedrine hydrochloride 60 mg

4 mg/60 mg

Antihis tamine/Nas al Deconges tant

Runny Nose & Sneezing, Itchy, Watery Eyes, Sinus Congestion & Pressure

24 Tablets

50844 REV0619K11108

Distributed by MAJOR® PHARMACEUTICALS 17177 N Laurel Park Drive, Suite 233 Livonia, MI 48152 Rev. 10/19 M-17 Re-order No. 235631

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Drug Facts (continued)

Inactive inaredients colloidal silicon dioxide, croscarmellose sodium, lactose anhydrous, magnesium stearate, microcrystalline cellulose, stearic acid

Questions or comments? (800) 616-2471

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS

Maximum Strength

SudoGest Sinus & Allergy

MAJOR°

NDC 0904-5351-24

Maximum Strength

SudoGest Sinus & Allergy

Chlorpheniramine maleate 4 mg Pseudoephedrine HCl 60 mg

4 mg/60 mg

Actual Size

Lot no. & Expiration Date

Antihistamine/Nasal Decongestant

Runny Nose & Sneezing, Itchy, Watery Eyes, Sinus Congestion & Pressure



24 Tablets

Pseudoephedrine HCl 60 mg.. Active ingredients

Nasal decongestant

(in each tablet

B-1212-111-08-R REV0619K11108

50844 REV0619K11108

Distributed by: MAJOR* PHARMACEUTICALS 17177 N Laurel Park Drive, Suite 233 Livonia, MI 48152 Rev. 10/19 M-17

Re-order No. 235631

Directions

ontrol Center (1-800-222-1222) right away

eep out of reach of children.

pregnant or breast-feeding, ask a health professional before use. **sep out of reach of children**. In case of overdose, get medical help or contact a Poison

■ children under 12 years: do not use

The r information

■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

■ Store in a dry place and protect from light ก light ก 15°-30°C (59°-86°F)

adults and children 12 years and over: take 1 tablet every 4 to 6 hours. Do not take more than 4 tablets in 24 hours.

■ avoid alcoholic beverages ■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

nervousness, dizziness, or sleeplessness occur symptoms do not improve within 7 days or occur with a fever

When using this product ■ do not exceed recommended dosage
■ drowsiness may occur ■ excitability may occur, especially in children
■ alcohol, sedatives and tranquilizers may increase drowsiness

lsk a doctor or pharmacist before use if you are taking sedatives or tranquilizers

■ diabetes ■ heart disease ■ high blood pressure ■ thyro
■ difficulty in unnation due to enlargement of the prostate gland
■ a breathing problem such as emphysema or chronic bronchitis

l temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper respiratory allergies: ■ runny nose ■ sneezing ■ fichy, watery eyes ■ nasal congestion ■ fiching of the nose or throat

temporarily relieves these symptoms due to the common cold:

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. sinus congestion and pressure

Major 44-111

SUDOGEST

chlorpheniramine maleate, pseudoephedrine hcl tablet

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-5351
Route of Administration	ORAL		

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	4 mg		
PSEUDO EPHEDRINE HYDRO CHLO RIDE (UNII: 6 V9 V2RYJ8 N) (PSEUDO EPHEDRINE - UNII:7CUC9 DDI9 F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg		

Inactive Ingredients				
Ingredient Name	Strength			
ANHYDRO US LACTO SE (UNII: 3S Y5LH9 PMK)				
CROSCARMELLOSE SODIUM (UNII: M28 OL1HH48)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
CELLULOSE, MICRO CRYSTALLINE (UNII: OP1R32D61U)				
SILICON DIO XIDE (UNII: ETJ7Z6 XBU4)				
STEARIC ACID (UNII: 4ELV7Z65AP)				

Product Characteristics			
Color	WHITE	Score	2 pieces
Shape	ROUND	Size	7mm
Flavor		Imprint Code	44;111
Contains			

]	Packaging				
1	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0904-5351-24	1 in 1 BOX	12/16/1992		
1	l	24 in 1 BLISTER PACK; Type 0: Not a Combination Product			
2	NDC:0904-5351-96	2 in 1 CARTON	12/16/1992		
2	2	24 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	12/16/1992	

Labeler - Major Pharmaceuticals (191427277)

Establishment			
Name	Address	ID/FEI	Business Operations

LNK International, Inc.	8328678	MANUFACTURE(0904-5351)	
-------------------------	---------	------------------------	--

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(0904-5351)

Revised: 12/2019 Major Pharmaceuticals