

**MAXIMUM STRENGTH ANTACID ANTIGAS- aluminum hydroxide, magnesium hydroxide, simethicone suspension
WALGREEN COMPANY**

WALgreens max antacid

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)

Magnesium hydroxide 400 mg

Simethicone 40mg

Purposes

Antacid

Antigas

Uses

relieves

- heartburn
- sour stomach
- acid indigestion
- the symptoms referred to as gas

Warnings

Do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are taking a prescription drug.

Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Directions

- shake well before use

- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals, at bedtime, or as directed by a doctor
- children under 12 years of age: ask a doctor

Other information

- **each 5 mL teaspoonful contains:** magnesium 165 mg, sodium 5 mg
- store at room temperature and avoid freezing
- keep tightly closed

Inactive ingredients

benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments?

1-800-540-3765

package Label

Walgreens



Compare to the active ingredients
in Maximum Strength Mylanta®[®]

Antacid + Anti-Gas Liquid

ALUMINA, MAGNESIA, AND SIMETHICONE
ORAL SUSPENSION / ANTACID & ANTI-GAS

Maximum Strength

- Fast relief of heartburn, pressure, bloating, acid indigestion & sour stomach
- ALCOHOL 0.5%



12 FL OZ (355 mL)

Classic
Flavor

222-06122-9 REV GC-0624

Drug Facts

TAMPER-EVIDENT: DO NOT USE IF THE BREAKAWAY BAND ON CAP IS BROKEN OR MISSING.

Active ingredients (in each 5 mL teaspoonful)

Aluminum hydroxide 400 mg (equivalent to dried gel, USP)	Antacid
Magnesium hydroxide 400 mg	Antacid
Simethicone 40 mg	Antigas

Purposes

Uses relieves • heartburn • sour stomach
• acid indigestion • symptoms referred to as gas

Warnings

Do not take more than 12 teaspoonfuls in 24 hours or use the maximum dosage for more than 2 weeks.

Ask a doctor before use if you have
• kidney disease • a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than 2 weeks

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.

Directions

- shake well before use
- adults and children 12 years and older: take 2 to 4 teaspoonfuls between meals as needed, at bedtime, or as directed by a doctor
- children under 12 years: ask a doctor

Other information

- each 5 mL teaspoonful contains: magnesium 165 mg, sodium 5 mg
- store at room temperature and avoid freezing • keep tightly closed

Inactive ingredients benzyl alcohol, butylparaben, carboxymethylcellulose sodium, flavor (contains alcohol), hypromellose, microcrystalline cellulose, propylparaben, purified water, saccharin sodium, sorbitol solution

Questions or comments? 1-800-540-3765

Your pharmacist recommends the Walgreens brand. We invite you to compare to national brands. This product is not manufactured or distributed by McNeil Consumer Pharmaceuticals Co., the owner of the registered trademark Mylanta®. DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD., DEERFIELD, IL 60015 3 100% SATISFACTION GUARANTEED walgreens.com ©2021 Walgreen Co. MADE IN U.S.A. W3ORFG1021-F ITEM 221700 W10071-1021-F 922-06122-9 REV GC-0624



MAXIMUM STRENGTH ANTACID ANTIGAS

aluminum hydroxide, magnesium hydroxide, simethicone suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-1691
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	400 mg in 5 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	400 mg in 5 mL
DIMETHICONE (UNII: 92RU3N3Y1O) (DIMETHICONE - UNII:92RU3N3Y1O)	DIMETHICONE	40 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
BUTYLPARABEN (UNII: 3QPI1U3FV8)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-1691-12	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/01/2014	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	09/01/2014	

Labeler - WALGREEN COMPANY (008965063)

Registrant - GCP Laboratories (965480861)

Establishment

Name	Address	ID/FEI	Business Operations
GCP Laboratories		965480861	manufacture(0363-1691) , pack(0363-1691)

Revised: 11/2025

WALGREEN COMPANY