

## **LOW DOSE ASPIRIN- aspirin tablet, coated**

### **Strategic Sourcing Services LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

## **Drug Facts**

### **Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

## **Purpose**

Pain reliever

## **Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor. **Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.**
- ask your doctor about other uses for safety coated 81 mg aspirin

## **Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives
- facial swelling
- shock
- asthma (wheezing)

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you are allergic to aspirin or any other pain reliever/fever reducer.
- if you have ever had an allergic reaction to his product or any of its ingredients

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are**

taking a prescription drug for

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stool
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever get worse or lasts more than 3 days
- redness or swelling is present
- ringing in the ears or loss of hearing occurs
- new symptoms occur

These could be signs of a serious condition

### **If pregnant or breast-feeding,**

ask a health professional before use. **It is especially important not to use aspirin at during the last 3 month of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor

- children under 12 years: consult a doctor

### **Other information**

- store between 15-30°C (59-86°F)

### **Inactive ingredients**

croscarmellose sodium, D&C yellow #10, lactose monohydrate, methacrylic acid, microcrystalline cellulose 102, polacrillin potassium, polyethylene glycol 6000, polyvinyl alcohol, purified talc, silicon dioxide, sodium lauryl sulfate, soybean, titanium dioxide, triethyl citrate

### **Questions or comments?**

Call **1-833-358-6431** Monday to Friday 9:00am to 7:00pm EST

### **Principal Display Panel**

**COMPARE TO BAYER® LOW DOSE ASPIRIN ACTIVE INGREDIENT†**

**LOW DOSE**

**Aspirin 81 mg**

**PAIN RELIEVER (NSAID)**

**Aspirin regimen±**

**Enteric coated±±**

**Safety coated±±**

**TABLETS**

†This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

≠Talk to your doctor or other healthcare provider before using this product for your heart. Aspirin is not right for everyone.

≠≠Coating helps protect against stomach upset

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**Distributed by: McKesson Corp.,**

via Strategic Sourcing Services LLC. Memphis, TN 38141

[www.fosterand thrive.com](http://www.fosterand thrive.com)

## Product Label

Exp. Date:

Lot No.:

Rev 03/23 PLD-A730A FC008594



Product of India

www.fosterandthrive.com

Money Back Guarantee

Sourcing Services LLC, Memphis, TN 38141

Distributed by: McKesson Corp., via Strategic

@2023 McKesson Corporation

### Drug Facts

#### Active ingredient (in each tablet)

Aspirin 81 mg (NSAID)\* .....Pain reliever  
\*nonsteroidal anti-inflammatory drug

#### Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor.
- Because of its delayed action, this product will not provide fast relief of headaches or other symptoms needing immediate relief.
- ask your doctor about other uses for safety coated 81 mg aspirin

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction, which may include:

- hives ■ facial swelling
  - shock ■ asthma (wheezing)
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: ■ are age 60 or older
- have had stomach ulcers or bleeding problems
  - take a blood thinner (anticoagulant) or steroid drug
  - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed

### Drug Facts (continued)

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic ■ you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for** ■ gout ■ diabetes ■ arthritis

#### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
  - you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - ringing in the ears or loss of hearing occurs
  - new symptoms occur
- These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Drug Facts (continued)

#### Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

#### Other information

- store between 15-30°C (59-86°F)

#### Inactive ingredients

croscarmellose sodium, D&C yellow #10, lactose monohydrate, methacrylic acid, microcrystalline cellulose 102, potassium hydroxide, polyethylene glycol 6000, polyvinyl alcohol, purified talc, silicon dioxide, sodium lauryl sulphate, soybean, titanium dioxide, triethyl citrate

#### Questions or comments?

Call 833-358-6431 Monday to Friday 9:00am to 7:00pm EST

\*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Bayer® Low Dose Aspirin.

†Talk to your doctor or other healthcare provider before using this product for your heart.

‡Aspirin is not right for everyone.

‡Coating helps protect against stomach upset

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

NDC 70677-1150-1

Foster & Thrive™

LOW DOSE

Aspirin 81 mg  
PAIN RELIEVER (NSAID)

Aspirin regimen†  
Enteric coated‡  
Safety coated‡‡

COMPARE TO BAYER® LOW DOSE ASPIRIN  
ACTIVE INGREDIENT

ACTUAL SIZE

120 TABLETS

## FOSTER & THRIVE Low Dose Aspirin

### LOW DOSE ASPIRIN

aspirin tablet, coated

#### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:70677-1150
<b>Route of Administration</b>	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ASPIRIN</b> (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

#### Inactive Ingredients

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>CROSCARMELOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>LACTOSE MONOHYDRATE</b> (UNII: EWQ57Q8I5X)	
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>METHACRYLIC ACID</b> (UNII: 1CS02G8656)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>POLACRILIN POTASSIUM</b> (UNII: 0BZ5A00FQU)	
<b>SODIUM LAURYL SULFATE</b> (UNII: 368GB5141J)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	
<b>POLYETHYLENE GLYCOL 6000</b> (UNII: 30IQX730WE)	
<b>SOYBEAN</b> (UNII: L7HT8F1ZOD)	
<b>POLYVINYL ALCOHOL, UNSPECIFIED</b> (UNII: 532B59J990)	

#### Product Characteristics

<b>Color</b>	yellow	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	P
<b>Contains</b>			

#### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1150-2	1 in 1 BOX	04/01/2023	
1		180 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
	NDC:70677			

2	NDC: 70677-1150-3	1 in 1 BOX	04/01/2023	
2		300 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
3	NDC: 70677-1150-1	1 in 1 BOX	04/01/2023	
3		120 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	04/01/2023	

**Labeler** - Strategic Sourcing Services LLC (116956644)

Revised: 5/2023

Strategic Sourcing Services LLC