

**ACID REDUCER- ranitidine tablet
Preferred Plus (Kinray)**

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride, USP 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do Not Use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR INDIVIDUAL BLISTER UNIT IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

Preferred plus Pharmacy[®]

NDC 61715-089-24

†Compare to the active ingredient of Zantac 150[®]

Maximum Strength

Ranitidine Tablets, USP 150 mg

Acid Reducer

24 TABLETS

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

Distributed by: Kinray Inc.

5104667/0813

Keep the carton. It contains important information.

Read the directions, consumer information leaflet and warnings before use.

<p>Questions? call 1-800-406-7984</p> <p>Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hydroxypropyl methylcellulose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide</p> <p>Other information ■ this product is sodium and sugar free ■ avoid excessive heat or humidity ■ store at 20° - 25° C (68° - 77° F) ■ OPEN OR TOURN. ■ CARTON OR INDIVIDUAL BUSTER UNITS ■ TAMPER EVIDENT: DO NOT USE IF THE</p> <p>Directions ■ adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water ■ to prevent symptoms, swallow 1 tablet with a glass of water ■ can be used up to twice daily (do not take more than 2 tablets in 24 hours) ■ children under 12 years: ask a doctor</p> <p>Drug Facts (continued)</p>	<p>Stop use and ask a doctor if ■ you need to take this product for more than 14 days ■ your heartburn continues or worsens</p> <p>Ask a doctor before use if you have ■ frequent chest pain, particularly with heartburn ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ had heartburn over 3 months. This may be a sign of a more serious condition ■ heartburn with lightheadedness, sweating or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness</p> <p>Warnings ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers ■ if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers ■ if you have kidney disease, except under the advice and supervision of a doctor</p> <p>Drug Facts (continued)</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

Drug Facts (continued)

Warnings
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ with other acid reducers
 ■ if you have kidney disease, except under the advice and supervision of a doctor

Do not use
 ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
 ■ with other acid reducers
 ■ if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have
 ■ frequent chest pain, particularly with heartburn
 ■ frequent wheezing, particularly with heartburn
 ■ unexplained weight loss
 ■ nausea or vomiting
 ■ stomach pain
 ■ had heartburn over 3 months. This may be a sign of a more serious condition
 ■ heartburn with **lightheadedness, sweating or dizziness**
 ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if
 ■ you need to take this product for more than 14 days
 ■ your heartburn continues or worsens

Questions? call 1-800-406-7984

24 TABLETS

Preferred Plus Pharmacy

Maximum Strength

Ranitidine Tablets, USP

150 mg

Acid Reducer

NDC 61715-80-24

Compare to the active ingredient of Zantac 150®

Preferred Plus Pharmacy

Maximum Strength

Ranitidine Tablets, USP

150 mg

Acid Reducer

24 TABLETS

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

SATISFACTION GUARANTEED

Expiry Date

NON VARNISH

Batch No.

KIN 263-368

1 (c) 2013 Kinney Inc. All Rights Reserved. PREFERRED PLUS, PREFERRED PLUS +++ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinney Inc. All other marks are property of their respective owners.

Distributed by: Kinney Inc., Whitestone, NY 11357
 All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased.

MADE IN INDIA

0813

ACID REDUCER

ranitidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61715-089
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL (Oval Shaped)	Size	12mm
Flavor		Imprint Code	9R
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:61715-089-24	3 in 1 CARTON		
1		8 in 1 BLISTER PACK		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	01/27/2014	

Labeler - Preferred Plus (Kinray) (012574513)

Registrant - Ranbaxy Pharmaceuticals Inc. (937890044)

Establishment

Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(61715-089)

Revised: 1/2014

Preferred Plus (Kinray)