

VALIANT PRE/POST- lactic acid and dodecylbenzene sulfonic acid solution
ABS Global Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, [click here](#).

KEEP OUT OF REACH OF CHILDREN
NOT FOR HUMAN USE

CAUTION: Not for internal use. May cause eye irritation. Avoid contact with eyes. Wash thoroughly after handling. For external use on dairy cows.

DIRECTIONS FOR USE

Mix Directions: Add 3.5 ounces of *Valiant Catalyst (US)* to one gallon of *Valiant Pre/Post* and mix (27.3 mL *Valiant Catalyst (US)* to 1 liter *Valiant Pre/Post (US)*). Inverting the container approximately 5-7 times will adequately mix the two solutions. This mixed solution will remain stable for 7 days. After 7 days, discard solution and mix a new solution.

IMPORTANT: Do not add mixed *Valiant Pre/Post (US)* ready-to-use to any other teat dip or other products. If transferred from this container to any other, make sure the other container is thoroughly precleaned and bears the proper container labeling for *Valiant Pre/Post (US)*.

Use of a complete cow-care program including both pre and post dipping may aid in reducing the spread of organisms which may cause mastitis.

DIRECTIONS FOR TEAT DIPPING

Pre-Milking: Fill teat dip cup with *Valiant Pre/Post (US)*. Before each cow is milked, dip the teats as far as possible into the teat dip cup containing *Valiant Pre/Post (US)*. Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Udder and teats which are heavily soiled should be washed before using a premilking teat dip. Teats should be dried with a single-service towel and dipped in *Valiant Pre/Post (US)*, wiping teats dry before milking.

Post-Milking: Immediately after milking, use *Valiant Pre/Post (US)* at full strength. Submerge teat 2/3 their length in *Valiant Pre/Post (US)* teat dip. Allow to air dry. Do

not wipe. Always use fresh full strength *Valiant Pre/Post (US)*. If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return unused product to the original container.

Do not turn cows into freezing weather until *Valiant Pre/Post (US)* is completely dry.

NOTE: *Valiant Pre/Post (US)* is not intended to cure or help in the healing of chapped or irritated teats. In case of irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

FIRST AID

Eye contact: Immediately flush eyes with plenty of water. Remove contact lenses and flush again. Get medical attention if irritation persists.

Skin contact: No known effect after skin contact. Rinse with water for a few minutes.

Inhalation: No special measures required. Treat symptomatically.

Ingestion: Get medical attention if symptoms occur.

FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, CALL: 1-800-328-0026.

FOR EMERGENCY MEDICAL INFORMATION WORLDWIDE, CALL: 1-651-222-5352 (IN THE USA).

Principal Display Panel and Representative Label

Valiant™

quality milk system

Pre/Post (US)

ABS

Sanitizing Pre/Post Teat Dip

Base - For use exclusively with Valiant Catalyst (US)

□An aid in reducing the spread of organisms which may cause mastitis.

ACTIVE INGREDIENTS:

Lactic Acid.....2.3%

Linear Dodecyl Benzene Sulfonic Acid.....1.6%

Contains an emollient system (lanolin and propylene glycol).....12.0%

ABS Global, Inc., P.O. Box 459
DeForest, Wisconsin 53532
www.absglobal.com

6300466

Patent Pending

ABS Global, Inc., P.O. Box 459
DeForest, Wisconsin 53532
www.absglobal.com

INSTRUCCIONES DE USO
Mazda: Agregue 105 ml (3.5 onzas) de Valiant Catalyst (US) a 3.8 litros (1 gal) de Valiant Pre/Post (US) y mezcle 27.3 ml de Valiant Catalyst (US) en 1.0 litro de Valiant Pre/Post (US). Para mezclar correctamente las dos soluciones, invierta el recipiente de 5 a 7 veces. Esta solución mezclada permanecerá estable por 7 días. Después de 7 días, tire la solución y mezcle otra nueva.

IMPORTANTE: No agregue Valiant Pre/Post (US) mezclado y listo para usar a ningún otro sellador de pezones o a otros productos. Si lo transfiere de este recipiente a otro, verifique que el otro esté minuciosamente limpio y que tenga la etiqueta de Valiant Pre/Post (US).

El uso de un programa completo para el cuidado de vacas, incluyendo el pre y post sellado, puede reducir la diseminación de organismos que pueden causar mastitis.

INSTRUCCIONES PARA EL SELLADO DE PEZONES
Pre-orden: Llene la copa de sellado con Valiant Pre/Post (US). Antes de ordenar cada vaca, sumerja los pezones hasta donde sea posible en la copa de sellado conteniendo Valiant Pre/Post (US). Luego, seque los pezones usando toallas desechables para evitar la contaminación de la leche.

Cuando las ubres y los pezones están demasiado sucios, se deben lavar antes de usar el sellador de pezones. Después de los pezones se deben lavar con una toalla desechable y sellarse con Valiant Pre/Post (US), y luego secarlos antes de la ordeña.

Post-orden: Inmediatamente después de la ordeña, use Valiant Pre/Post (US) sin diluir. Sumerja el pezón 2/3 partes de su tamaño en sellador Valiant Pre/Post (US). Deleto secal al aire. No lo limpie. Siempre use Valiant Pre/Post (US) fresco, sin diluir. Si el producto en la copa de sellado se seca, tire la solución y mezcle otra nueva a llenar con producto sin diluir. No vuelva a usar el producto residual ni lo devuelva al recipiente original.

No saque las vacas a la intermedia helada hasta que Valiant Pre/Post (US) se haya secado completamente.

NOTA: Valiant Pre/Post (US) no intenta curar o ayudar a cicatrizar pezones irritados o agrietados. En caso de irritación o agrietamiento, consulte a un veterinario.

MANTÉNGALO ALEJADO DEL ALCANCE DE LOS NIÑOS
NO ES PARA USO HUMANO
PRECAUCIÓN: No es para uso interno. Puede provocar irritación ocular. Evite el contacto con los ojos. Lávese completamente después del manejo. Para uso externo en vacas lecheras.

PRIMEROS AUXILIOS
Contacto con los ojos: Lávese de inmediato los ojos con abundante agua. Quite los lentes de contacto y lávese con abundante agua. Obtenga atención médica si persiste la irritación.

Contacto con la piel: No se conocen efectos sobre el contacto con la piel, enjuague con agua durante unos minutos. Inhalación: No se requieren medidas especiales. Tratar sintomáticamente. Ingestión: Obtenga atención médica si se presentan síntomas.

PARA INFORMACIÓN MEDICA DE EMERGENCIA EN LOS EE.UU. O CANADA, LLAME AL 1-800-328-9026.

PARA INFORMACIÓN MEDICA DE EMERGENCIA EN EL MUNDO ENTERO, LLAME AL 1-651-222-5352 (EN LOS EE.UU.).

Los ingredientes de la fórmula no contienen foltano.

Pre/post (us)



Valiant™

quality milk
system



6300466

Patent Pending

ABS Global, Inc., P.O. Box 459
DeForest, Wisconsin 53532
www.absglobal.com

DIRECTIONS FOR USE
Mix Directions: Add 3.5 ounces of Valiant Catalyst (US) to one gallon of Valiant Pre/Post and mix 27.3 ml of Valiant Catalyst (US) to 1 liter Valiant Pre/Post (US). Inverting the container approximately 5-7 times will adequately mix the two solutions. This mixed solution will remain stable for 7 days. After 7 days, discard solution and mix a new solution.

IMPORTANT: Do not add mixed Valiant Pre/Post (US) ready-to-use to any other test dip or other products. If transferred from this container to any other, make sure the other container is thoroughly pre-cleaned and bears the proper container labeling for Valiant Pre/Post (US).

Use of a complete cow-care program including both pre and post dipping may aid in reducing the spread of organisms which may cause mastitis.

Sanitizing Pre/Post Teat Dip

Base – For use exclusively with Valiant Catalyst (US)
An aid in reducing the spread of organisms which may cause mastitis.

ACTIVE INGREDIENTS:

Lactic Acid.....	2.3%
Linear Dodecyl Benzene Sulfonic Acid.....	1.6%
Contains an emollient system (lanolin and propylene glycol).....12.0%	

Base exclusiva para sellar pezones

Base – Para usar exclusivamente con Valiant Catalyst (US)
Una ayuda para reducir la diseminación de organismos que pueden causar mastitis.

INGREDIENTES ACTIVOS:

Ácido láctico.....	2.3%
Ácido lineal dodecilsulfónico.....	1.6%
Contiene un sistema emoliente (lanolina y glicol propileno).....12.0%	

**KEEP OUT OF REACH OF CHILDREN
NOT FOR HUMAN USE**

CAUTION: Not for internal use. May cause eye irritation. Avoid contact with eyes. Wash thoroughly after handling. For external use on dairy cows.

FIRST AID
Eye contact: Immediately flush eyes with plenty of water. Remove contact lenses and flush again. Get medical attention if irritation persists.
Skin contact: No known effect after skin contact. Rinse with water for a few minutes.
Inhalation: No special measures required. Treat symptomatically.
Ingestion: Get medical attention if symptoms occur.

FOR EMERGENCY MEDICAL INFORMATION IN USA OR CANADA, CALL: 1-800-328-9026

FOR EMERGENCY MEDICAL INFORMATION WORLDWIDE, CALL: 1-651-222-5352 (IN THE USA).

DIRECTIONS FOR TEST DIPPING
Pre-Milking: Fill test dip with Valiant Pre/Post (US). Before each cow is milked, dip the teats as far as possible into the test dip containing Valiant Pre/Post (US). Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Under and teats which are heavily soiled should be washed before using a pre-milking test dip. Teats should be dried with a single-service towel and dipped in Valiant Pre/Post (US), wiping teats dry before milking.

Post-Milking: Immediately after milking, use Valiant Pre/Post (US) at full strength. Submerge teat 2/3 their length in Valiant Pre/Post (US) test dip. Allow to air dry. Do not wipe. Always use fresh full strength Valiant Pre/Post (US). If product in dip becomes soiled or dirty, discard product and replenish with undiluted product. Do not reuse or return unused product to the original container.

Do not turn cows into freezing weather until Valiant Pre/Post (US) is completely dry.

NOTE: Valiant Pre/Post (US) is not intended to cure or help in the healing of chapped or irritated teats. In case of irritation or chapping, have the condition examined and, if necessary, treated by a veterinarian.

© 2012 Hecho en EEUU 703828/5300/0012
Todos los derechos reservados

© 2012 Made in USA 703828/5300/0012
All Rights Reserved

Formula ingredients contain no Propofol.

lactic acid and dodecylbenzene sulfonic acid solution

Product Information			
Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:59282-102
Route of Administration	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
LACTIC ACID (UNII: 33X04XA5AT) (LACTIC ACID - UNII:33X04XA5AT)		LACTIC ACID	23 mg in 1 mL
DODECYLBENZENESULFONIC ACID (UNII: 60NSK897G9) (DODECYLBENZENESULFONIC ACID - UNII:60NSK897G9)		DODECYLBENZENESULFONIC ACID	16 mg in 1 mL
Inactive Ingredients			
Ingredient Name			Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
PEG-75 LANOLIN (UNII: 09179OX7TB)			
WATER (UNII: 059QF0KO0R)			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:59282-102-16	18900 mL in 1 PAIL		
2	NDC:59282-102-17	56780 mL in 1 DRUM		
3	NDC:59282-102-21	208200 mL in 1 DRUM		
4	NDC:59282-102-44	1150000 mL in 1 PACKAGE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		09/09/1998	

Labeler - ABS Global Inc. (835053570)

Revised: 9/2018

ABS Global Inc.