

**TRIPLE ANTIBIOTIC PLUS- bacitracin zinc, neomycin sulfate, polymyxin b sulfate and pramoxine hydrochloride ointment  
Meijer, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Meijer Triple Antibiotic Plus Pain Relief Ointment 0.5oz (94028, 2018)**  
**Meijer Triple Antibiotic Plus Pain Relief Ointment 1.0oz (94037, 2018)**  
**Meijer Triple Antibiotic Plus Pain Relief Ointment 1.0oz Twin (94029, 2018)**

**Active ingredients (in each gram) Purposes**

Bacitracin zinc 500 units.....First Aid antibiotic  
Neomycin sulfate 3.5 mg.....First Aid antibiotic  
Polymyxin B sulfate 10,000 units.....First Aid antibiotic  
Pramoxine HCl 10 mg.....First Aid antibiotic

**Uses**

- first aid to help prevent infection and for the temporary relief of pain in minor:
- cuts
- scrapes
- burns

**Warnings**

For external use only

**Do not use**

- if you are allergic to any of the ingredients
- in the eyes or nose
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**Stop use and ask a doctor if**

- you need to use longer than 1 week
- rash or other allergic reaction develops
- symptoms persist for more than 1 week or clear up and occur again within a few days
- condition persists or gets worse
- redness, irritation, swelling, or pain persists or increases

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

**Directions**

- adults and children 2 years of age and older:
- clean the affected area
- apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the

area 1 to 3 times

- may be covered with a sterile bandage
- children under 2 years of age: ask a doctor

Other information

- store at room temperature 20-25°C (68-77°F)

Inactive ingredients alcohol, methylparaben, mineral oil, petrolatum, polyoxl 40 stearate, propylparaben

Questions or comments?

Call 1-800-814-8028

DIST. BY MEIJER DISTRIBUTION, INC.

GRAND RAPIDS, MI 49544

[www.meijer.com](http://www.meijer.com)

MADE IN SOUTH KOREA

# MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

**+ PAIN RELIEF**

**Dual Action**  
Soothes Cuts, Scrapes & Burns

**Powerful Protection & Pain Relief for  
Cuts | Scrapes | Burns**

**meijer**

## MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

**+ PAIN RELIEF**

NET WT 0.5 OZ (14 g)

First Aid Antibiotic/Pain Relieving Ointment

Compare to Neosporin®  
+ Pain Relief Ointment  
active ingredients\*

MAXIMUM STRENGTH  
triple antibiotic  
+ pain relief ointment  
Bacitracin Zinc | Neomycin Sulfate |  
Polymyxin B Sulfate | Pramoxine HCl



DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

MADE IN SOUTH KOREA

94028-TK-BX-R1  
OM-MD-TK106-R1

**Drug Facts** (continued)

**Active Ingredients (in each gram)** / **Purposes**

Bacitracin zinc 500 units.....First aid antibiotic  
Neomycin sulfate 3.5 mg.....First aid antibiotic  
Polymyxin B sulfate 10,000 units.....First aid antibiotic  
Pramoxine HCl 10 mg.....External analgesic

**Uses**

First aid to help prevent infection and for the temporary relief of pain in minor: ■ cuts ■ scrapes ■ burns

**Warnings**

For external use only

**Do not use**

■ if you are allergic to any of the ingredients  
■ in the eyes or nose ■ over large areas of the body

**Ask a doctor before use if you have**

■ deep or puncture wounds ■ animal bites ■ serious burns

**Stop use and ask a doctor if**

■ you need to use longer than 1 week  
■ rash or other allergic reaction develops

■ symptoms persist for more than 1 week, or clear up and occur again within a few days ■ condition persists or gets worse

■ redness, irritation, swelling, or pain persists or increases

**Directions**

■ adults and children 2 years of age and older:  
■ clean the affected area  
■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily ■ may be covered with a sterile bandage

■ children under 2 years of age: ask a doctor

**Other information**

■ store at room temperature 20-25°C (68-77°F)

**Inactive ingredients** alcohol, methylparaben, mineral oil, petrolatum, polyoxy 40 stearate, propylparaben

**Questions or comments?**  
Call 1-800-814-8028

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Neosporin®.

TAMPER EVIDENT. DO NOT USE IF CARTON IS OPEN OR FOIL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTSIDE CARTON FOR COMPLETION.

MAXIMUM STRENGTH  
triple antibiotic  
+ pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

Dual Action  
Soothes Cuts, Scrapes & Burns

Powerful Protection & Pain Relief for  
Cuts | Scrapes | Burns

meijer

MAXIMUM STRENGTH

triple antibiotic  
+ pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

Compare to Neosporin®  
+ Pain Relief Ointment  
active ingredients\*

+ PAIN RELIEF

First Aid Antibiotic/Pain Relieving Ointment

NET WT 1 OZ (28 g)

MAXIMUM STRENGTH  
triple antibiotic  
+ pain relief ointment  
Bacitracin Zinc | Neomycin Sulfate |  
Polymyxin B Sulfate | Pramoxine HCl



DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com

Made in South Korea

94029-TK-BX-R1  
OM-MD-TK106-R1

**Drug Facts (continued)**

**Active ingredients (in each gram)** Purposes: Bacitracin zinc 500 units. First aid antibiotic. Neomycin sulfate 3.5 mg. First aid antibiotic. Polymyxin B sulfate 10,000 units. First aid antibiotic. Pramoxine HCl 10 mg. External analgesic.

**Uses**  
First aid to help prevent infection and for the temporary relief of pain in minor: cuts, scrapes, burns.

**Warnings**  
For external use only.  
Do not use:  
If you are allergic to any of the ingredients or over large areas of the body.  
Ask a doctor before use if you have:  
deep or puncture wounds, animal bites, serious burns.  
Stop use and ask a doctor if:  
you need to use longer than 1 week.  
rash or other allergic reaction develops.  
symptoms persist for more than 1 week, or clear up and occur again within a few days.  
redness, irritation, swelling, or pain persists or increases.

**Directions**  
adults and children 2 years of age and older:  
clean the affected area.  
apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily.  
may be covered with a sterile bandage.  
children under 2 years of age: ask a doctor.  
store at room temperature 20-25°C (68-77°F).

**Other information**  
petroleum, polyoxy 40 stearate, methylparaben, mineral oil.

**Inactive ingredients**  
petroleum, polyoxy 40 stearate, methylparaben, mineral oil.

**Questions or comments?** Call 1-800-814-8028.  
\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Neosporin®.

TAMPEN EVIDENT; DO NOT USE IF CARTON IS OPEN OR POIL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLET.

94029-TK-BX-R1  
OM-MD-TK106-R1

DIST. BY MEIJER DISTRIBUTION, INC.  
GRAND RAPIDS, MI 49544  
www.meijer.com  
MADE IN SOUTH KOREA



MAXIMUM STRENGTH  
triple antibiotic  
+ pain relief ointment  
Bacitracin Zinc | Neomycin Sulfate |  
Polymyxin B Sulfate | Pramoxine HCl



PFD 2727846

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Neosporin®.

**Questions or comments?** Call 1-800-814-8028

**Inactive ingredients** alcohol, methylparaben, mineral oil, petrolatum, polyoxyl 40 stearate, propylparaben

**Other information** ■ store at room temperature 20-25°C (68-77°F)

■ children under 2 years of age: ask a doctor

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily ■ may be covered with a sterile bandage

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ clean the affected area

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

**Drug Facts**

**Active ingredients (in each gram)**

Bacitracin zinc 500 units

Neomycin sulfate 3.5 mg

Polymyxin B sulfate 10,000 units

Pramoxine HCl 10 mg

**Purposes**

First aid antibiotic

External analgesic

**Uses**

First aid to help prevent infection and for the temporary relief of

■ pain in minor: ■ cuts ■ scrapes ■ burns

**Warnings**

For external use only

Do not use

■ if you are allergic to any of the ingredients

■ in the eyes or nose

■ over large areas of the body

**Ask a doctor before use if you have**

■ deep or puncture wounds

■ animal bites

■ serious burns

**Stop use and ask a doctor if**

■ you need to use longer than 1 week

■ rash or other allergic reaction develops

■ symptoms persist for more than 1 week, or clear up and occur again within a few days

■ redness, irritation, swelling, or pain persists or increases

**Drug Facts (continued)**

**Directions**

■ adults and children 2 years of age and older:

■ apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times

■ daily

■ may be covered with a sterile bandage

■ children under 2 years of age: ask a doctor

**Other information**

■ store at room temperature 20-25°C (68-77°F)

**Inactive ingredients** alcohol, methylparaben, mineral oil, petrolatum, polyoxyl 40 stearate, propylparaben

**Questions or comments?** Call 1-800-814-8028

\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Neosporin®.

TAMPER EVIDENT: DO NOT USE IF CAP OR CARTON IS OPEN OR FOIL UNDER CAP IS BROKEN OR MISSING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

VALUE 2 PACK

MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

Dual Action Soothes Cuts, Scrapes & Burns

Powerful Protection & Pain Relief for Cuts | Scrapes | Burns

meijer

MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

First Aid Antibiotic/Pain Relieving Ointment

Compare to Neosporin® + Pain Relief Ointment active ingredients\*

meijer

MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

First Aid Antibiotic/Pain Relieving Ointment

Compare to Neosporin® + Pain Relief Ointment active ingredients\*

2 TUBES

NET WT 1 OZ (28 g) | TOTAL NET WT 2 OZ (56 g)

meijer

MAXIMUM STRENGTH triple antibiotic + pain relief ointment

Bacitracin Zinc | Neomycin Sulfate | Polymyxin B Sulfate | Pramoxine HCl

+ PAIN RELIEF

First Aid Antibiotic/Pain Relieving Ointment

Compare to Neosporin® + Pain Relief Ointment active ingredients\*

TRIPLE ANTIBIOTIC PLUS

M triple antibiotic + pain relief Bacitracin Polymyxin

bacitracin zinc, neomycin sulfate, polymyxin b sulfate and pramoxine hydrochloride ointment

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:41250-961
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
NEOMYCIN SULFATE (UNII: 057Y626693) (NEOMYCIN - UNII:I16QD7X297)	NEOMYCIN SULFATE	3.5 mg in 1 g
PRAMOXINE HYDROCHLORIDE (UNII: 88AYB867L5) (PRAMOXINE - UNII:068X84E056)	PRAMOXINE HYDROCHLORIDE	10 mg in 1 g
BACITRACIN ZINC (UNII: 89Y4M234ES) (BACITRACIN - UNII:58H6RWO52I)	BACITRACIN	500 [iU] in 1 g
POLYMYXIN B SULFATE (UNII: 19371312D4) (POLYMYXIN B - UNII:J2VZ07J96K)	POLYMYXIN B	10000 [iU] in 1 g

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
MINERAL OIL (UNII: T5L8T28FGP)	
ALCOHOL (UNII: 3K9958V90M)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
POLYOXYL 40 STEARATE (UNII: 13A4J4NH9I)	
PETROLATUM (UNII: 4T6H12BN9U)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:41250-961-14	1 in 1 CARTON	12/18/2018	
1		14 g in 1 TUBE; Type 0: Not a Combination Product		
2	NDC:41250-961-56	2 in 1 CARTON	12/18/2018	
2		28 g in 1 TUBE; Type 0: Not a Combination Product		
3	NDC:41250-961-28	1 in 1 CARTON	12/18/2018	
3		28 g in 1 TUBE; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part333B	12/18/2018	

**Labeler** - Meijer, Inc. (006959555)