

**STOMACH RELIEF REGULAR STRENGTH- bismuth subsalicylate suspension
CARDINAL HEALTH 110, LLC. DBA LEADER**

Regular Strength Stomach Relief Drug Facts

Active ingredient (in each 30 mL dose)

Bismuth subsalicylate 525 mg

Purposes

Upset stomach reliever and antidiarrheal

Uses

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

Warnings

Reye's syndrome:Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

Do not use

if you have

- an ulcer
- a bleeding problem
- bloody or black stool

Ask a doctor before use if you have

- fever
- mucus in the stool

Ask a doctor or pharmacist before use if you are

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

When using this product

a temporary, but harmless, darkening of the stool and/or tongue may occur

Stop use and ask a doctor if

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- shake well before use
- Only use dose cup provided
- **adults and children 12 years and over:** 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

Other information

- **each 30 mL contains:** potassium 14 mg, sodium 10 mg
- salicylate 260 mg
- low sodium
- sugar free
- store at room temperature
- protect from freezing
- avoid excessive heat (more than over 104°F or 40°C).

Inactive ingredients

carboxymethylcellulose sodium, D&C red #22, D&C red #28, flavor, gellan gum, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, Simethicone emulsion, sodium benzoate, sucralose, xanthan gum

Questions or comments?

1-866-467-2748

Safety Sealed: DO NOT USE IF IMPRINTED SHRINK BAND IS MISSING OR BROKEN

Principal Display Panel

NDC 70000-0698-1

COMPARE TO PEPTO-BISMOL® ORIGINAL LIQUID active ingredient*

LEADER™

Regular Strength

Stomach Relief

Bismuth Subsalicylate, 525 mg

Upset Stomach Reliever/Antidiarrheal

Original Flavor

Relieves:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

8 FL OZ (236 mL)

100% Money Back Guarantee

Dist. By CAH DUBLIN, OH 43017

www.myleader.com

1-800-200-6313

©2024 Cardinal Health.

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Original Liquid.

6.75" Width

3.25" Height

Drug Facts (continued)
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.
Directions ■ shake well before use ■ only use dose cup provided ■ adults and children 12 years and over: ■ 30 mL (1 dose) every ½ hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea ■ 30 mL (1 dose) every ½ hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea) ■ do not exceed 8 doses (240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea
Other information ■ each 30 mL contains: potassium 14 mg, sodium 10 mg ■ salicylate 260 mg ■ low sodium ■ sugar free ■ store at room temperature ■ protect from freezing ■ avoid excessive heat (over 104°F or 40°C)
Inactive ingredients carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, gellan gum, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum
Questions or comments? 1-866-467-2748

LOT: 70608LDRLR DIST. BY CAH DUBLIN, OH 43017
www.myleader.com 1-800-200-6313
©2024 Cardinal Health.

CIN 5920020 REV. 06/24



EXP:

NDC 70000-0698-1

Regular Strength

Stomach Relief

Bismuth Subsalicylate, 525 mg
Upset Stomach Reliever / Antidiarrheal

Original Flavor

Relieves:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

COMPARE TO
PEPTO-BISMOL®
ORIGINAL LIQUID
active ingredient*

100% Money
Back Guarantee

8 FL OZ (236 mL)

Do not use if imprinted shrink band is missing or broken

Drug Facts

Active ingredient (in each 30 mL dose)	Purpose
Bismuth subsalicylate 525 mg	Upset stomach reliever and antidiarrheal

Uses relieves ■ travelers' diarrhea ■ diarrhea ■ upset stomach due to overindulgence in food and drink, including: ■ heartburn ■ indigestion ■ nausea ■ gas ■ belching ■ fullness

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Contains salicylate. Do not take if you are allergic to salicylates (including aspirin) ■ taking other salicylate products

Do not use if you have ■ an ulcer ■ a bleeding problem ■ bloody or black stool

Ask a doctor before use if you have ■ fever ■ mucus in the stool

Ask a doctor or pharmacist before use if you are taking any drug for ■ anticoagulation (thinning the blood) ■ diabetes ■ gout ■ arthritis

When using this product a temporary, but harmless, darkening of the stool and/or tongue may occur.

Stop use and ask a doctor if ■ symptoms get worse or last more than 2 days ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days

*This product is not manufactured or distributed by Procter & Gamble, distributor of Pepto-Bismol® Original Liquid.

STOMACH RELIEF REGULAR STRENGTH

bismuth subsalicylate suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0698
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISMUTH SUBSALICYLATE (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	
D&C RED NO. 22 (UNII: 1678RKX8RT)	
D&C RED NO. 28 (UNII: 7671P0Y5NH)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POTASSIUM HYDROXIDE (UNII: WZH3C48M4T)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
WATER (UNII: 059QF0KO0R)	
SALICYLIC ACID (UNII: O414PZ4LPZ)	
DIMETHICONE, UNSPECIFIED (UNII: 92RU3N3Y1O)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3Z D4)	

XANTHAN GUM (UNII: TTV12P4NEE)

GELLAN GUM (HIGH ACYL) (UNII: W1L7G7ROMD)

Product Characteristics

Color	pink (viscous)	Score	
Shape		Size	
Flavor	WINTERGREEN (ORIGINAL)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0698-1	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	10/21/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	10/21/2024	

Labeler - CARDINAL HEALTH 110, LLC. DBA LEADER (063997360)

Revised: 10/2025

CARDINAL HEALTH 110, LLC. DBA LEADER