

**S.T.37- hexylresorcinol solution**  
**NUMARK BRANDS, INC**

-----  
**Drug Facts**

<b>Active Ingredient</b>	<b>Purposes</b>
Hexylresorcinol 0.1%	Oral Pain Reliever
	First aid antiseptic

**Uses** temporarily relieves:

- minor irritations of the mouth
- minor pain in the mouth
- sore mouth
- sore throat
- first aid to help prevent skin infection in minor cuts, scrapes, and burns

**Warnings**

**For external use only**

**Sore throat warning** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- In the eyes
- over large areas of the body

**Ask a doctor before use if you have**

- deep or puncture wounds
- animal bites
- serious burns

**When using this product**

- **do not exceed recommended dosage**

**Stop use and ask a doctor if**

- condition persists
- condition worsens

- you need to use product longer than 1 week

**Stop use and ask a doctor or dentist if**

- sore mouth symptoms do not improve in 7 days
- irritation, pain, and redness persist or worsen

**If pregnant or breast feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** (as a First Aid antiseptic)

- use full strength
- clean affected area
- apply a small amount of product on the area
- use 1-3 times daily
- may be covered with a sterile bandage
- if bandaged let dry first

**Directions** (as an Oral pain reliever)

- apply to affected area
- gargle, swish around, or allow to remain in place for at least one minute and then spit out

adults and children 2 years old and over	use up to 4 times daily or as directed by a dentist or doctor
children under 2 years old	ask a dentist or doctor

**Other Information**

- **store at controlled room temperature 59° - 86°F (15° - 30°C)**

**Inactive Ingredients**

citric acid, edetate disodium, glycerin, propylene glycol, sodium bisulfite, sodium citrate, purified water.

**Questions** call 1-800-214-2379

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL**

**8 oz SIZE**

**2.5 more ounces at the same low price**

**S.T.37®  
FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
SOLUTION**

**FOR BURNS, CUTS, ABRASIONS & HYGIENIC CARE OF THE MOUTH.**

**Do not use if foil seal under cap is broken or missing at time of purchase**

8 FL OZ (236.5 mL)

WON'T STING — WON'T STAIN

S.T.37 kills common pathogenic bacteria quickly — on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an order.

**Everyday Uses for S.T.37**

**Cuts & Abrasions:** Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

**Burns:** Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

**Hygienic Care of Mouth:** Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.  
27070 Miles Rd, Suite A, Solon Ohio 44139  
[www.numarkbrands.com](http://www.numarkbrands.com)

B10082  
Rev. 3/18

NOIT 70 S

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER

# S.T.37

WON'T STING—WON'T STAIN

## S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
SOLUTION

### Drug Facts

**Active ingredient** Hexyresorcinol 0.1%  
**Purposes** Oral pain reliever  
First aid antiseptic

**Uses** temporarily relieves:  
■ minor irritations of the mouth ■ sore mouth  
■ minor pain in the mouth ■ sore throat  
■ first aid to help prevent skin infection in minor cuts, scrapes, and burns

#### Warnings

**For external use only**

**Sore throat warning** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly.

#### Do not use

■ in the eyes ■ over large areas of the body

**Ask a doctor before use if you have**

■ deep or puncture wounds ■ animal bites  
■ serious burns

**When using this product** ■ do not exceed recommended dosage

**Stop use and ask a doctor if**

■ condition persists ■ condition worsens  
■ you need to use product longer than 1 week

**Stop use and ask a doctor or dentist if**

■ sore mouth symptoms do not improve in 7 days  
■ irritation, pain, and redness persist or worsen

**If pregnant or breast feeding, ask a health professional before use.**

WON'T STING—WON'T STAIN

## S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
SOLUTION

### Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** (as a First Aid antiseptic)  
■ use full strength ■ clean affected area  
■ apply a small amount of product on the area  
■ use 1-3 times daily  
■ may be covered with a sterile bandage  
■ if bandaged, let dry first

**Directions** (as an Oral pain reliever)  
■ apply to affected area  
■ gargle, swish around, or allow to remain in place for at least one minute and then spit out

adults and children  
2 years old and over

use up to 4 times  
daily or as directed  
by a dentist or doctor  
children under 2  
years old ask a dentist or  
doctor

**Other information** ■ store at controlled room temperature 59°-86°F (15°-30°C)

**Inactive ingredients** citric acid, edetate disodium, glycerin, propylene glycol, sodium bisulfite, sodium citrate, purified water.

**Questions** call 1-800-214-2379

Distributed by Numark Brands, Inc.  
27070 Miles Rd, Suite A, Solon Ohio 44139  
www.numarkbrands.com

**8 oz SIZE**  
2.5 more ounces  
at the same low price

# S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
SOLUTION

FOR BURNS,  
CUTS, ABRASIONS  
& HYGIENIC CARE  
OF THE MOUTH.

Do not use if foil seal under cap is  
broken or missing at time of purchase

8 FL OZ (236.5 mL)

WON'T STING—WON'T STAIN

## S.T.37

FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
SOLUTION

S.T.37 kills common pathogenic bacteria quickly —on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

#### Everyday Uses for S.T.37

**Cuts & Abrasions:** Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

**Burns:** Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

**Hygienic Care of Mouth:** Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.



B10082  
Rev. 3/18

**S.T.37®**  
**FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER**  
**SOLUTION**

## FOR BURNS, CUTS, ABRASIONS & HYGIENIC CARE OF THE MOUTH

8 FL OZ (236.5 mL)

**Do not use if foil seal under cap is broken or missing at time of purchase**

S.T.37 kills common pathogenic bacteria quickly - on contact!

Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

### Everyday Uses for S.T.37

**Cuts & Abrasions:** Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

**Burns:** Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

**Hygienic Care of Mouth:** Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.

27070 Miles Rd, Suite A, Solon Ohio 44139 1-800-214-2379

www.numarkbrands.com Rev. 3/18 J12008

**REFER TO OUTER CARTON FOR FULL INFORMATION**

**USES:** temporarily relieves: • minor irritations of the mouth • minor pain in the mouth • sore mouth • sore throat • first aid to help prevent skin infection in minor cuts, scrapes, and burns.

**DIRECTIONS:** (as a First Aid antiseptic) • use full strength • clean affected area • apply a small amount of product on the area • use 1-3 times daily • may be covered with a sterile bandage • if bandaged let dry first.

**DIRECTIONS:** (as an Oral pain reliever) • apply to affected area • gargle, swish around, or allow to remain in place for at least one minute and then spit out. For adults and children 2 years old and over: use up to 4 times daily or as directed by a dentist or doctor. For children under 2 years old: ask a dentist or doctor.

**WARNINGS:** For external use only. **Sore throat warning** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, swelling, nausea, or vomiting, consult a doctor promptly. **Do not use** • in the eyes • over large areas of the body. **Ask a doctor before use if you have** • deep or puncture wounds • animal bites • serious burns. **When using this product** • do not exceed recommended dosage. **Stop use and ask a doctor if** • condition persists • condition worsens • you need to use product longer than 1 week. **Stop use and ask a doctor or dentist if** • sore mouth symptoms do not improve in 7 days • irritation, pain, and redness persist or worsen. **If pregnant or breast feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**ACTIVE INGREDIENT:** Hexylresorcinol 0.1%.

**INACTIVE INGREDIENTS:** Citric acid, Edetate disodium, Glycerin, Propylene glycol, Sodium bisulfite, Sodium citrate, Purified water.



**ST.37**  
FIRST AID ANTISEPTIC/ORAL PAIN RELIEVER  
**SOLUTION**

**FOR BURNS, CUTS, ABRASIONS & HYGIENIC CARE OF THE MOUTH.**

8 FL OZ (236.5 mL)

**Do not use if foil seal under cap is broken or missing at time of purchase**

S.T.37 kills common pathogenic bacteria quickly - on contact! Laboratory tests have established S.T.37 antiseptic solution as a general antiseptic for household use.

S.T.37 is pleasant to use because it produces little or no stinging or burning. In fact, it soothes irritated areas.

S.T.37 does not stain the skin or leave an odor.

**Everyday Uses for S.T.37**

**Cuts & Abrasions:** Apply full strength S.T.37 into the cut or over the injured area. Since it is germicidal for the common pathogenic bacteria, S.T.37 may minimize the development of infection. Excellent for cleansing cuts and scrapes.

**Burns:** Apply full strength S.T.37 for treatment of minor burns. S.T.37 combats the cause of infections.

**Hygienic Care of Mouth:** Use undiluted and gargle or rinse to help keep mouth clean and fresh. S.T.37 temporarily relieves minor pain and protects irritated areas of a sore mouth or throat.

Distributed by Numark Brands, Inc.  
27070 Miles Rd, Suite A, Solon Ohio 44139 1-800-214-2379  
www.numarkbrands.com Rev. 3/18 J12008

## S.T.37

hexylresorcinol solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69846-860
<b>Route of Administration</b>	ORAL, TOPICAL		

### Active Ingredient/Active Moiety

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>Hexylresorcinol</b> (UNII: R9QTB5E82N) (Hexylresorcinol - UNII:R9QTB5E82N)	Hexylresorcinol	1.1 mg in 1 mL

### Inactive Ingredients

<b>Ingredient Name</b>	<b>Strength</b>
<b>Citric Acid Monohydrate</b> (UNII: 2968PHW8QP)	
<b>Edetate Disodium</b> (UNII: 7FLD91C86K)	
<b>Glycerin</b> (UNII: PDC6A3C0OX)	
<b>Propylene Glycol</b> (UNII: 6DC9Q167V3)	
<b>Sodium Bisulfite</b> (UNII: TZX5469Z6I)	
<b>Sodium Citrate</b> (UNII: 1Q73Q2JULR)	
<b>Water</b> (UNII: 059QF0KO0R)	

### Packaging

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:69846-860-00	1 in 1 AMPULE	11/20/1929	
1		236.5 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:69846-860-01	1 in 1 CARTON	11/20/1929	
2		473 mL in 1 BOTTLE; Type 0: Not a Combination Product		

### Marketing Information

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M022	11/20/1929	

**Labeler** - NUMARK BRANDS, INC (080184668)