

**LOW DOSE CHEWABLE ASPIRIN- aspirin tablet, chewable**  
**VALU MERCHANDISERS COMPANY**

-----  
**1105B-BST-2024-0613**

***Drug Facts***

**Active ingredient (in each tablet)**

Aspirin 81 mg (NSAID)\*

\* nonsteroidal anti-inflammatory drug

**Purpose**

Pain reliever

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

**Warnings**

**Reye's syndrome**

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert**

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning**

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)

- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

### **Do not use**

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor

### **Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

### **Ask a doctor or pharmacist before use if you are taking a prescription drug for:**

- gout
- diabetes
- arthritis

### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint
  - vomit blood
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

corn starch, FD&C yellow #6 aluminum lake, flavor, magnesium stearate, silicon dioxide, sucrose

**Questions or comments?**

1-844-292-1112

**PRINCIPAL DISPLAY PANEL**

Best Choice®

Compare to the active ingredient in Bayer® Chewable Aspirin†

LOW DOSE

Chewable Aspirin 81 mg

PAIN RELIEVER (NSAID)

FOR ADULTS

ASPIRIN REGIMEN\*\*

ORANGE FLAVORED

Talk to your doctor or other healthcare provider before using this product for your heart

Actual Size

81 mg each

36 TABLETS

INK AND COATING FREE  
FOR LOT AND  
EXPIRATION STAMPING

F1105801BST\_R1  
20166-0424-AAA  
BEST CHOICE



0 170082 20166 3

DO NOT USE IF IMPRINTED SEAL  
UNDER CAP IS BROKEN OR MISSING

This product is not manufactured or distributed by Bayer Healthcare LLC, distributor of Bayer® Chewable Aspirin.

**Drug Facts**

**Active ingredient (in each tablet)** **Purpose**  
Aspirin 81 mg (NSAID) .....Pain reliever  
nonsteroidal anti-inflammatory drug

**Uses**

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for this product

**Warnings**

- Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
- Allergy alert:** Aspirin may cause a severe allergic reaction which may include:
- hives
  - facial swelling
  - asthma (wheezing)
  - shock
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
- are age 60 or older
  - have had stomach ulcers or bleeding problems
  - take a blood thinning (anticoagulant) or steroid drug

**Drug Facts (continued)**

- Take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
  - have 3 or more alcoholic drinks every day while using this product
  - take more or for a longer time than directed
- Do not use**
- if you have ever had an allergic reaction to any other pain reliever/fever reducer
  - for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- Ask a doctor before use if**
- stomach bleeding warning applies to you
  - you have a history of stomach problems, such as heartburn
  - you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
  - you are taking a diuretic
  - you have asthma
- Ask a doctor or pharmacist before use if you are taking a prescription drug for:**
- gout
  - diabetes
  - arthritis

**Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - vomit blood
  - feel faint
  - have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days

**Drug Facts (continued)**

- fever gets worse or lasts more than 3 days
  - redness or swelling is present
  - new symptoms occur
  - ringing in the ears or a loss of hearing occurs
- These could be signs of a serious condition.
- If pregnant or breast-feeding, ask a health professional before use.** It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
- Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- store between 20°-25°C (68°-77°F) in a dry place
- retain carton for complete product information and warnings

**Inactive ingredients**

- corn starch, FD&C yellow #6 aluminum lake, titanium dioxide, silicon dioxide, sucrose

PROUDLY DISTRIBUTED BY:  
ASSOCIATED WHOLESALE GROCERS, INC.  
KANSAS CITY, KANSAS 66106

**Best Choice**  
100% Guaranteed

www.bestchoicebrand.com

Aspirin is not appropriate for everyone, so be sure to talk to your doctor before you begin an aspirin regimen.

Scan here for more product information.  
Call 1-844-292-1112 for more product information.

1-844-292-1112

**Questions or comments?**

**Drug Facts (continued)**

Compare to the active ingredient in Bayer® Chewable Aspirin†

**Best Choice®**

**LOW DOSE**

**Chewable Aspirin 81 mg**

**PAIN RELIEVER (NSAID)**

FOR ADULTS ASPIRIN REGIMEN\*\*

ORANGE FLAVORED

Talk to your doctor or other healthcare provider before using this product for your heart

**36 TABLETS**

Actual Size




LOW DOSE CHEWABLE ASPIRIN			
aspirin tablet, chewable			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63941-105
Route of Administration	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	81 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SUCROSE (UNII: C151H8M554)	

**Product Characteristics**

Color	orange	Score	no score
Shape	ROUND	Size	8mm
Flavor	ORANGE	Imprint Code	AP;009
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63941-105-01	1 in 1 CARTON	10/15/2021	06/30/2027
1		36 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:63941-105-03	3 in 1 CARTON	04/15/2024	03/31/2027
2		36 in 1 PACKAGE; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	10/15/2021	06/30/2027

**Labeler** - VALU MERCHANDISERS COMPANY (868703513)