CLARITIN REDITABS- loratadine tablet, orally disintegrating
Bayer HealthCare LLC

Claritin ®

RediTabs ®

Drug Facts

Active ingredient (in each tablet)

Loratadine 5 mg

Purpose

Antihistamine

Uses

temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25°C (68° to 77°F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or **www.claritin.com**

Distributed by MSD Consumer Care, Inc.,
PO Box 377, Memphis, TN 38151 USA,
a subsidiary of Merck & Co., Inc., Whitehouse Station, NJ USA.

PRINCIPAL DISPLAY PANEL - 10 Tablet Carton

NDC 11523-7202-3

Non-Drowsy*

Claritin ®

RediTabs ®

*loratadine **5 mg**/antihistamine*

Indoor & Outdoor

Allergies

****When taken as directed. See Drug Facts Panel.***

12

Hour

Relief of:

- *Sneezing*
- *Runny Nose*
- *Itchy, Watery Eyes*
- *Itchy Throat*

or Nose

No Water Needed

Melts in Your Mouth

10

ORALLY

DISINTEGRATING TABLETS

in each tablet) **Purpose**

.....Antihistamine

eves these symptoms due to hay fever or
lergies:
- itchy, watery eyes
- itching of the nose or throat

er had an allergic reaction to this product or

f you have liver or kidney disease. Your
f you need a different dose.

do not take more than directed. Taking more
lrowsiness.

r if an allergic reaction to this product occurs.
way.

ding, ask a health professional before use.
dren. In case of overdose, get medical help or
Center right away.

; tablet disintegrates, with or without water

s	1 tablet every 12 hours; not more than 2 tablets in 24 hours
age	ask a doctor
	ask a doctor

Drug Facts (continued)

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or **www.claritin.com**

Claritin®

RediTabs®

12 Hour

• No Water Needed • Melts in your mouth

Follow these directions carefully. Do not attempt to push the tablet through the foil.



1. Peel back outer cage.



2. Gently push tablet out.



3. Place the tablet on tongue and close mouth. The tablet will disintegrate.

CLARITIN REDITABS

loratadine tablet, orally disintegrating

Product Information			
Product Type	HUMAN OTC DRUG	**Item Code (Source)**	NDC:11523-7202
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	**Basis of Strength**	**Strength**
LORATADINE (UNII: 7AJO3BO7QN) (LORATADINE - UNII:7AJO3BO7QN)	LORATADINE	5 mg

Inactive Ingredients	
Ingredient Name	**Strength**
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GELATIN (UNII: 2G86QN327L)	
MANNITOL (UNII: 3OWL53L36A)	

Product Characteristics			
Color	white	**Score**	no score
Shape	ROUND (flat faced beveled edge)	**Size**	12mm
Flavor	MINT	**Imprint Code**	C5
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11523-7202-1	3 in 1 CARTON	12/12/2006	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11523-7202-3	1 in 1 CARTON	12/12/2006	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NDA	NDA021993	12/12/2006	

Labeler - Bayer HealthCare LLC (112117283)

Revised: 12/2025 Bayer HealthCare LLC