INTEGRITY PRE-POST PART B- sodium chlorite solution Premier Select Sires

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Integrity Pre-Post Part B

Integrity Pre-Post Part B
ClO2 Teat Dip Part B

WARNING

KEEP OUT OF REACH OF CHILDREN.

AVOID FREEZING.

KEEP OUT OF DIRECT SUNLIGHT.

CAUSES SERIOUS EYE IRRITATION.

FOR FARM AND INDUSTRIAL USE ONLY.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice/ attention.

STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal

plant .

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

NET CONTENTS: 55 Gallons

LOT:

EXP. DATE:

Use Premier Integrity Pre-Post effectively to reduce the spread of mastitis causing organisms.

Recommended for pre and post milking applications.

DIRECTIONS:

IMPORTANT: MUST BE MIXED PRIOR TO USE. Mix PART A & PART B in equal parts. Mix by volume.

CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing, and will continue to generate Chlorine Dioxide for up to 5-7 days. After 7 days, discard solution and mix a new solution.

General Use

Once mixed according to directions, do not mix product with any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

Pre Dipping Use

Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

Post Dipping Use

Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to

air dry. Do not wipe.

If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

Extended Use

When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking.

Dipping should be resumed at least twice daily one week before calving.

Manufactured For: Premier Select Sires 1 Stony Mountain Road Tunkhannock PA 18657 Phone (800)-227-6417



WARNING

KEEP OUT OF REACH OF CHILDREN. AVOID FREEZING. KEEP OUT OF DIRECT SUNLIGHT. Causes serious eye irritation. For farm and industrial use only.

PREVENTION: Wash skin thoroughly after handling. Wear eye protection/ face protection.

FIRST AID / RESPONSE:

IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical

advice/attention.STORAGE: Store locked up.

DISPOSAL: Dispose of contents/ container to an approved waste disposal plant.

EMERGENCY TELEPHONE NUMBER (Chemtrec) 800-424-9300 CCN205494

ACTIVE INGREDIENTS: Sodium Chlorite 0.45%

Net Contents: 55 Gallon

Lot:

Exp. Date:

Manufactured For: Premier Select Sires 1 Stony Mountain Road Tunkhannock, PA 18657 Phone (800) 227-6417



CIO₂ Teat dip Part B

Use Premier Integrity Pre-Post effectively to reduce the spread of mastitis causing organisms. Recommended for pre and post milking applications.

DIRECTIONS:

IMPORTANT: MUST BE MIXED PRIOR TO USE. Mix PART A & PART B in equal parts. Mix by volume. CREATES CHLORINE DIOXIDE. Reaches minimum Chlorine Dioxide levels within 15 minutes of mixing, and will continue to generate Chlorine Dioxide for up to 5-7 days. After 7 days, discard solution and mix a new solution.

GENERAL USE: Once mixed according to directions, do not mix product with any other teat dips, or other product.

Use a teat dip cup to apply. Clean dip cup just prior to use.

If cow's teats are sore or chapped, consult a veterinarian before starting or continuing to use teat dip.

If transferred from this container to any other, make sure that the other container is thoroughly pre-cleaned and bears the proper container labeling for this product.

If product in dip cup becomes visibly dirty, discard contents and replenish with undiluted product. Do not reuse or return any unused product to the original container.

This product can also be foamed or sprayed using recommended equipment.

PRE DIPPING USE: Perform forestripping and inspect teats for clinical mastitis.

Pre-wash teats thoroughly just prior milking with appropriate udder wash solution or pre-milking teat dip. Teats should then be dried with single-service towels.

Just before each cow is milked, dip, foam, or spray, each teat full-length into the teat dip or foam cup containing product.

Allow product to stand for 30 seconds.

Wipe teats dry after dipping, using single-service towels to avoid contamination of milk.

POST DIPPING USE: Immediately after each milking, use product.

Dip each teat full-length into the teat dip cup containing product. Allow to air dry. Do not wipe. If outside temperature is below freezing, allow to air dry on the teat before cow leaves the parlor.

EXTENDED USE: When a cow is being dried off, the teats should be dipped in product, once a day for 3-4 days after the last milking. Dipping should be resumed at least twice daily one week before calving.

INTEGRITY PRE-POST PART B

sodium chlorite solution

Product Information

Product Type OTC ANIMAL DRUG Item Code (Source) NDC:86012-3004

Route	of Administration	TOPICAL

ı	Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength	
	SODIUM CHLORITE (UNII: G538EBV4VF) (CHLORITE ION - UNII: Z63H374SB6)	SODIUM CHLORITE	0.014526 kg in 1 kg	

P	Packaging						
#	Item Code	Package Description	Marketing Start Date	Marketing End Date			
1	NDC:86012-3004-1	19 kg in 1 CONTAINER					
2	NDC:86012-3004-2	57 kg in 1 DRUM					
3	NDC:86012-3004-3	208 kg in 1 DRUM					
4	NDC:86012-3004-4	1041 kg in 1 CONTAINER					

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
unapproved drug other		11/21/2022		

Labeler - Premier Select Sires (065398711)

Revised: 11/2022 Premier Select Sires