

**LEADER KIDS SPF 50 SUNSCREEN- homosalate, octisalate, oxybenzone, octocrylene, avobenzene aerosol, spray  
Cardinal Health**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Leader Kids SPF 50 Continuous Spray Sunscreen**

**Active Ingredients**

Avobenzene 3.0%, Homosalate 10.0%, Octisalate 5.0%, Octocrylene 4.0%, Oxybenzone 5.0%

**Purpose**

Sunscreen

**Uses**

• helps prevent sunburn • if used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

**Warnings**

For external use only

**FLAMMABLE**

- do not use near heat, flame or while smoking
- avoid long term storage above 104 deg F (40 degC)

**Do not use**

on damaged or broken skin

**When using this product**

keep out of eyes. Rinse with water to remove. Do not puncture or incinerate. Contents under pressure. Do not store at temperatures above 120 deg F.

**Stop use and ask a doctor**

if rash occurs.

## **Keep out of reach of children.**

If product is swallowed, get medical help or contact a Poison Control Center right away.

## **Directions**

- Apply liberally 15 minutes before sun exposure
- Hold can 4-6 inches away from body, spray evenly to ensure complete coverage
- Do not spray into face. Spray into hand and apply to face.
- Use in well ventilated, but not windy areas
  
- Reapply:
  - After 80 minutes of swimming or sweating
  - Immediately after towel drying
  - At least every 2 hours
  
- Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 am - 2 pm
  - wear long sleeved shirts, pants, hats, and sunglasses

## **Inactive Ingredients**

Alcohol denat. (73.2% V/V), acrylates/octylacrylamide copolymer, stearytrimethylsilane, glycerin, diethylhexyl syringylidenemalonate, tocopherol (vitamin E), fragrance, retinyl palmitate (vitamin A palmitate), caprylic/capric triglyceride



Compare to Coppertone® Kids® SPF 50+

# Kids Sunscreen

## Clear Continuous Spray

### BROAD SPECTRUM SPF 50

# 50

Sprays at Any Angle  
Water Resistant  
(80 Minutes)



6 FL OZ (177mL)

#### Drug Facts

Active Ingredients	Purpose
Avobenzone 3.0% Homosalate 10.0% Octisalate 5.0% Octocrylene 4.0% Oxybenzone 5.0%	Sunscreen

**Uses** • helps prevent sunburn • if used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun

#### Warnings

For external use only

#### FLAMMABLE:

- do not use near heat, flame or while smoking
- avoid long term storage above 104°F (40°C)

Do not use on damaged or broken skin.

**When using this product** • keep out of eyes. Rinse with water to remove. • do not puncture or incinerate. Contents under pressure. • do not store at temperatures above 120° F.

**Stop use and ask a doctor** if rash occurs.

Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

- apply liberally 15 minutes before sun exposure
- hold can 4-6 inches away from body, spray evenly to ensure complete coverage
- do not spray into face. Spray into hand and apply to face.
- use in well ventilated, but not windy areas
- reapply:
  - after 80 minutes of swimming or sweating
  - immediately after towel drying
  - at least every 2 hours
- **Sun Protection Measures.** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad-spectrum SPF of 15 or higher and other sun protection measures including:
  - limit time in the sun, especially from 10 a.m. - 2 p.m.
  - wear long-sleeve shirts, pants, hats, and sunglasses
- children under 6 months: ask a doctor

#### Inactive Ingredients

alcohol denat. (73.2% V/V), acrylates/octylacrylamide copolymer, stearyltrimethylsilane, glycerin, diethyhexyl syringylidenemalonate, tocopherol (vitamin E), fragrance, retinyl palmitate (vitamin A palmitate), caprylic/capric triglyceride

#### Other information

- protect this product from excessive heat and direct sun
- may stain or damage some fabrics, materials or surfaces

**Questions or comments?**  
Call toll free 1-800-200-6313

\*This product is not manufactured or distributed by Schering-Plough HealthCare Products, Inc., distributor of Coppertone® brand products.

All Leader® Brand products are 100% satisfaction guaranteed or return to place of purchase for a full refund.

DISTRIBUTED BY  
CARDINAL HEALTH  
DUBLIN, OHIO 43017  
CIN 4246666  
www.myleader.com



DOT-SP 14786  
M5655



## LEADER KIDS SPF 50 SUNSCREEN

homosalate, octisalate, oxybenzone, octocrylene, avobenzone aerosol, spray

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37205-723
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	100 mg in 1 mL
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	50 mg in 1 mL
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6VM) (OCTOCRYLENE - UNII:5A68WGF6VM)	OCTOCRYLENE	40 mg in 1 mL
<b>OXYBENZONE</b> (UNII: 950OS7VE0Y) (OXYBENZONE - UNII:950OS7VE0Y)	OXYBENZONE	50 mg in 1 mL
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	30 mg in 1 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>STEAROXYTRIMETHYLSILANE</b> (UNII: 9862TW94B2)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>DIETHYLHEXYL SYRINGYLIDENEMALONATE</b> (UNII: 3V5U97P248)	
<b>TOCOPHEROL</b> (UNII: R0ZB2556P8)	
<b>VITAMIN A PALMITATE</b> (UNII: 1D1K0N0VVC)	
<b>MEDIUM-CHAIN TRIGLYCERIDES</b> (UNII: C9H2L21V7U)	
<b>ACRYLATE/ISOBUTYL METHACRYLATE/N-TERT-OCTYLACRYLAMIDE COPOLYMER (75000 MW)</b> (UNII: JU3XHR8VWK)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37205-723-30	177 mL in 1 CAN; Type 0: Not a Combination Product	01/15/2012	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	01/15/2012	

**Labeler** - Cardinal Health (097537435)

Revised: 1/2022

Cardinal Health